

Vv156 Lecture 27

Dr Jing Liu

UM-SJTU Joint Institute

December 8, 2016

- For the harmonic series, Bernoulli noticed that the subsequence diverges

$$s_2, \quad s_4, \quad s_8, \quad s_{16}, \quad s_{32}, \quad \dots$$

$$s_2 = 1 + \frac{1}{2}$$

$$s_4 = 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) > 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right) = 1 + 1$$

$$s_8 = 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right) > 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right) + \left(\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\right) = 1 + \frac{3}{2}$$

\vdots

- Hence he concluded that the harmonic series diverges; however, we might not be able to do the same to every other series, for example,

Q: Does the following series converge?

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots + \frac{1}{n^2} + \dots$$

- There is no closed formula for the partial sum of this series, so we cannot check

$$\lim_{n \rightarrow \infty} s_n$$


- And the test for divergence is inconclusive,

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{1}{n^2} = 0$$

- We can determine the convergence of $\sum_{n=1}^{\infty} \frac{1}{n^2}$ by comparing it with $\int_1^{\infty} \frac{1}{x^2} dx$.

- Let $f(x) = \frac{1}{x^2}$, then the partial sum is

$$\begin{aligned} s_n &= \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{n^2} \\ &= f(1) + f(2) + f(3) + \cdots + f(n) \\ &< f(1) + \int_1^n f(x) dx \end{aligned}$$



- Thus the limit of the partial sum is less than the following sum

$$s_n < f(1) + \int_1^n f(x) dx \implies \lim_{n \rightarrow \infty} s_n < f(1) + \lim_{n \rightarrow \infty} \int_1^n f(x) dx$$


- However, we have the following alternative inequality

$$s_n = f(1) + f(2) + \cdots + f(n) > \int_1^n f(x) dx \implies \lim_{n \rightarrow \infty} s_n > \lim_{n \rightarrow \infty} \int_1^n f(x) dx$$

- So the series will be convergent if the improper integral is convergent, and will diverge if the improper integral diverges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_1^n f(x) dx &= \lim_{n \rightarrow \infty} \int_1^n \frac{1}{x^2} dx \\ &= \lim_{n \rightarrow \infty} \left[-\frac{1}{x} \right]_1^n = 1 \end{aligned}$$

- Hence the series is convergent.



Integral test

Suppose $\{a_n\}$ is a sequence such that $a_n = f(n)$, where $f(x)$ is a

1. continuous,
2. positive,
3. decreasing function on $[1, \infty)$.

Then the series $\sum_{n=1}^{\infty} a_n$ and $\int_1^{\infty} f(x) dx$ both converge or both diverge.

Exercise

Test the series

$$\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$$

for convergence or divergence.

The Comparison Test

- Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are series with positive terms.

If

1. $\sum_{n=1}^{\infty} b_n$ is convergent and

2. $a_n \leq b_n$ for all n .

then the series $\sum a_n$ is convergent.

If

1. $\sum_{n=1}^{\infty} b_n$ is divergent and

2. $a_n \geq b_n$ for all n .

then the series $\sum a_n$ is divergent.

The limit Comparison Test

- Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are series with positive terms.

If

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c, \quad \text{where}$$

1. if c is a finite and $c > 0$, then either both series converge or both diverge.
2. if $c = 0$ and $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} a_n$ converges.
3. if $c = \infty$ and $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges.

Exercise

For what values of p is the series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ convergent?

Alternating Series test

If the alternating series $\sum_{n=1}^{\infty} (-1)^{n-1} b_n$, where $b_n > 0$, satisfies

$$1. b_{n+1} \leq b_n \quad \text{and} \quad 2. \lim_{n \rightarrow \infty} b_n = 0$$

then the series is convergent.

Proof

- We consider the even-numbered partial sum $s_2, s_4, s_6, \dots, s_{2n}$,

$$s_{2n} = b_1 - b_2 + b_3 - b_4 + b_5 - b_6 + \dots + b_{2n-1} - b_{2n}$$

- Because $b_n - b_{n+1} \geq 0$ for all n , the sequence of partial sums $\{s_{2n}\}$ is increasing.
- Also, because, for all n ,

$$s_{2n} = b_1 - (b_2 - b_3) - (b_4 - b_5) - \dots - (b_{2n-2} - b_{2n-1}) - b_{2n} \leq b_1$$

Proof

- Therefore the sequence $\{s_{2n}\}$ is bounded above as well as being increasing.
- Clearly an increasing sequence is bounded below, and thus the limit exists

$$s = \lim_{n \rightarrow \infty} s_{2n}$$

by the monotonic sequence theorem.

- It remains to show the odd-numbered partial sums s_{2n+1} also converges to s .

$$\lim_{n \rightarrow \infty} s_{2n+1} = \lim_{n \rightarrow \infty} s_{2n} + \lim_{n \rightarrow \infty} b_{2n+1} = s \quad \square$$

Exercise

Test the series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n^2}{n^3 + 1}$ for convergence or divergence.

Q: It is sufficient but not necessary. Can you think of a counterexample?

Definition

A series $\sum_n^{\infty} a_n$ is called **absolutely convergent** if the series $\sum_n^{\infty} |a_n|$ is convergent.

Q: Is alternating harmonic series absolutely convergent?

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n}$$

- It is **not absolutely convergent** because the corresponding series of absolute values is the harmonic series and is therefore divergent.

Q: Is it convergent?

- However, it can be shown that it **is convergent** by the alternating series test.

Definition

A series $\sum_n^{\infty} a_n$ is known as **conditionally convergent** if the series is convergent but **not** absolutely convergent.

Exercise

(a) Prove the following using mathematical induction.

$$\frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} \cdots + \frac{1}{2n-1} - \frac{1}{2n} = \frac{1}{n} \left[\frac{1}{1 + \frac{1}{n}} + \frac{1}{1 + \frac{2}{n}} + \cdots + \frac{1}{1 + \frac{n}{n}} \right]$$

(b) Find the value to which the alternating harmonic series converges to.

Riemann series theorem

If series $\sum_n a_n$ is a conditionally convergent and r is any real number, then there is a **rearrangement** of $\sum_n a_n$ that has a sum equal to r .

$$\begin{aligned} - \quad 0 &= (1 - 1) + (1 - 1) + (1 - 1) + \cdots \\ &\neq 1 - 1 + 1 - 1 + 1 - 1 + \cdots \neq 1 + (-1 + 1) + (-1 + 1) + (-1 + 1) + \cdots \\ &= 1 \end{aligned}$$

- If a series is an absolutely convergent with sum s , then any **rearrangement** of it has the same sum s . However, any series that is only **conditionally convergent** can be rearranged to give a different sum.
- If we halve the alternating harmonic series,

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots = \ln 2 \implies \frac{1}{2} - \frac{1}{4} + \frac{1}{6} - \frac{1}{8} + \cdots = \frac{1}{2} \ln 2$$

- Now if we introduce some zeros

$$\begin{array}{ccccccccc} \frac{1}{2} & & -\frac{1}{4} & & +\frac{1}{6} & & -\frac{1}{8} & & +\cdots & = \frac{1}{2} \ln 2 \\ 0 + \frac{1}{2} + 0 & & -\frac{1}{4} + 0 & & +\frac{1}{6} + 0 & & -\frac{1}{8} + 0 & & +\cdots & = \frac{1}{2} \ln 2 \end{array}$$

- If we add the alternating harmonic series and the last series together,

$$1 + \frac{1}{3} - \frac{1}{2} + \frac{1}{5} + \frac{1}{7} - \frac{1}{4} + \cdots = \frac{3}{2} \ln 2$$

- The last series contains the same terms as the original alternating harmonics series, but rearranged so that one negative term occurs after two positive terms.

Theorem

If a series is absolutely convergent, then it is convergent.

Proof

- Notice

$$0 \leq a_n + |a_n| \leq 2|a_n|$$

- Absolutely convergence means the series $\sum_{n=1}^{\infty} |a_n|$ is convergent, thus the series

$\sum_{n=1}^{\infty} 2|a_n|$ converges, so by the comparison test the following series converges

$$\sum_{n=1}^{\infty} (a_n + |a_n|)$$

- So $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (a_n + |a_n|) - \sum_{n=1}^{\infty} |a_n|$ converges since both series converge.

The Ratio Test

This test is useful in determining whether a given series is absolutely convergent.

$$\begin{aligned}\text{If } \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= L < 1 \\ &\implies \text{Absolutely convergent} \\ &\implies \text{Convergent}\end{aligned}$$

$$\begin{aligned}\text{If } \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= L > 1 \\ &\implies \text{divergent}\end{aligned}$$

$$\begin{aligned}\text{If } \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= 1 \\ &\implies \text{Inconclusive}\end{aligned}$$

Exercise

Test the series $\sum_{n=1}^{\infty} (-1)^n \frac{n^3}{3^n}$ for absolute convergence.

- If we have integer powers, it is more convenient to use the following test, e.g.

$$\sum_{n=1}^{\infty} \left(\frac{2n+3}{3n+2} \right)^n$$

The Root Test

$$\begin{aligned} \text{If } \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L < 1 \\ \implies \text{Absolutely convergent} \\ \implies \text{Convergent} \end{aligned}$$

$$\begin{aligned} \text{If } \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L > 1 \\ \implies \text{divergent} \end{aligned}$$

$$\begin{aligned} \text{If } \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = 1 \\ \implies \text{Inconclusive} \end{aligned}$$