

# VP390 Problem Set 5

Pan, Chongdan ID:516370910121

June 18, 2020

## 1 Problem 1

- (b) Assume  $\psi$  is the solution for  $-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$

Then  $-\frac{\hbar^2}{2m} \frac{d^2\psi(-x)}{dx^2} + V(-x)\psi(-x) = E\psi(-x)$

Since  $V(x) = V(-x)$

$-\frac{\hbar^2}{2m} \frac{d^2\psi(-x)}{dx^2} + V(x)\psi(-x) = E\psi(-x)$

$\psi(-x)$  also satisfies this stationery equation.

Therefore  $\psi(x) \pm \psi(-x)$  satisfies this equation as well.

## 2 Problem 2

- (a) When  $x < 0$ ,  $\Psi(x) = 0$

When  $x > a$ ,  $\Psi(x) = Ae^{-\kappa_1 x}$ , where  $\kappa_1^2 = -\frac{2mE}{\hbar^2}$

When  $0 \leq x \leq a$ ,  $V(x) = -V_0 \rightarrow \frac{\partial^2 \Psi(x)}{\partial x^2} + \frac{2m}{\hbar^2}(E + V_0)\Psi(x) = 0$

Let  $\kappa_2 = \sqrt{\frac{2m}{\hbar^2}(E + V_0)}$ ,  $\Psi(x) = C_1 \cos \kappa_2 x + C_2 \sin \kappa_2 x$

$\Psi(0) = 0 \rightarrow C_1 = 0$

$\Psi(a)' = -A\kappa_1 e^{-\kappa_1 a} = \kappa_2 C_2 \cos \kappa_2 a \rightarrow C_2 = \frac{A\kappa_1 e^{-\kappa_1 a}}{\kappa_2 \cos \kappa_2 a}$

$\int_{-\infty}^{\infty} |\Psi(x)|^2 dx = \int_0^a \left( \frac{A\kappa_1 e^{-\kappa_1 a}}{\kappa_2 \cos \kappa_2 a} \cos \kappa_2 x \right)^2 dx + \int_a^{\infty} A^2 e^{-2\kappa_1 x} dx = 1$

$\frac{A^2}{2\kappa_1} e^{-2\kappa_1 a} + \left( \frac{A\kappa_1 e^{-\kappa_1 a}}{\kappa_2 \cos \kappa_2 a} \right)^2 \frac{\sin(2a\kappa_2) + 2a\kappa_2}{4\kappa_2} = 1$

$A = \sqrt{\frac{8\kappa_1 \kappa_2 (\kappa_2 \cos \kappa_2 a)^2}{e^{-2\kappa_1 a} 4\kappa_2 (\kappa_2 \cos \kappa_2 a)^2 + 2\kappa_1 (\sin(2a\kappa_2) + 2a\kappa_2) (\kappa_1 e^{-\kappa_1 a})^2}}$

$\kappa_1^2 + \kappa_2^2 = \frac{2m}{\hbar^2} V_0$

Since the bound states only exists when  $x > 0$ , which has the similar form with finite well when  $\psi$  is odd, then the bound states are only the original odd wave bound states in finite well problem.

- (b) The solution when  $x > a$  has the same form of solution with same value of  $\kappa_1, \kappa_2$ , only  $C_2, A$  need recalculated. For  $0 \leq x \leq a$ , it has the same form as the odd solution with finite potential well.

When  $x < -a$ ,  $\Psi(x) = Be^{-\kappa_1 x}$ , where  $\kappa_1^2 = -\frac{2mE}{\hbar^2}$

When  $x > a$ ,  $\Psi(x) = Ae^{-\kappa_1 x}$

When  $0 \leq x \leq a$ , If the  $\psi$  is odd,  $\psi = C \sin \kappa_2 x$ , which has the solution of same form with this problem's.

- (c) According to Figure.1 shows that only first bound state is for even function.




Figure 1: Energy level for finite well

Therefore, it's possible to set the width and the depth of the well so that there is no bound state or only one bound state.

- (d) According to above graph, to have only one bound states in odd function, the radius should  $\frac{\pi}{2} \leq r < \frac{3\pi}{2}$

$$\text{Since } a = r \sqrt{\frac{\hbar^2}{2mV_0}}$$

$$\text{Then } 1.62 \times 10^{-10} m \leq r < 4.85 \times 10^{-10} m$$

### 3 Problem 3

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x)}{\partial x^2} - E\Psi(x) = V_0 \delta(x) \Psi(x)$$

Integrate the equation over an infinitesimal interval  $[-\varepsilon, \varepsilon]$ , we get:

$$-\frac{\hbar^2}{2m} (\Psi(\varepsilon)' - \Psi(-\varepsilon)') - E \int_{-\varepsilon}^{\varepsilon} \Psi(x) dx = V_0 \Psi(0)$$

$$\text{When } x \neq 0, \frac{\partial^2 \Psi(x)}{\partial x^2} + \frac{2m}{\hbar^2} E \Psi(x) = 0$$

$$\text{When } x < 0, \Psi(x) = A e^{\kappa x}, \text{ where } \kappa^2 = -\frac{2mE}{\hbar^2}$$

$$\text{When } x > 0, \Psi(x) = B e^{-\kappa x}$$

Since  $\Psi(x)$  should be continuous,  $\Psi(0) = A = B$ , hence  $\Psi(x)$  is even because  $V(x)$  is symmetric.

Then  $\Psi(\varepsilon)' = -\Psi(-\varepsilon)'$ , and the original integral equation becomes:

$$\Psi(\varepsilon)' = -\frac{m}{\hbar^2} (E \int_{-\varepsilon}^{\varepsilon} \Psi(x) dx + V_0 \Psi(0)) = \frac{m}{\hbar^2} [2E \frac{A}{\kappa} (e^{-\kappa \varepsilon} - 1) - V_0 \Psi(0)]$$

$$\lim_{\varepsilon \rightarrow 0} \Psi(\varepsilon)' = -\frac{m}{\hbar^2} V_0 A \neq 0, \text{ hence } \Psi(x)' \text{ is not continuous at } x = 0$$

$$\text{For normalization } \int_{-\infty}^0 |A e^{\kappa x}|^2 dx + \int_0^{\infty} |A e^{-\kappa x}|^2 dx = 1$$

$$\frac{A^2}{\kappa} = 1 \rightarrow A = \sqrt{\kappa}$$

$$\text{Plug it back, when } x > 0, \text{ we get } \Psi(x)' = \kappa^{\frac{3}{2}} e^{-\kappa x} = \frac{mA}{\hbar^2} [2E \frac{1}{\kappa} (e^{-\kappa x} - 1) - V_0]$$

$$\text{Then we get } -\frac{2E}{\kappa} = V_0 \rightarrow E = -\frac{V_0^2 m}{2\hbar^2}$$

## 4 Problem 4

1.  $\Psi(x, 0) = \sqrt{\frac{1}{3}}\psi_3 + \sqrt{\frac{2}{3}}\psi_5$

where  $\psi_n = C_n H_n(\frac{x}{x_0}) e^{-\frac{1}{2}(\frac{x}{x_0})^2}$ ,  $C_n = \frac{1}{\sqrt{2^n n! x_0 \sqrt{\pi}}}$ ,  $x_0 = \sqrt{\frac{\hbar}{m\omega}}$

$\langle \Psi_n, \Psi_m \rangle = \int_{-\infty}^{\infty} \Psi_n^* \Psi_m dx = \int_{-\infty}^{\infty} H_n(\frac{x}{x_0}) H_m(\frac{x}{x_0}) C_n C_m e^{-(\frac{x}{x_0})^2} dx$

Let  $y = \frac{x}{x_0}$ , then  $\langle \Psi_n, \Psi_m \rangle = \int_{-\infty}^{\infty} H_n(y) H_m(y) C_n C_m e^{-y^2} x_0 dy$

According to the property, when  $m \neq n$ ,  $\int_{-\infty}^{\infty} H_n(y) H_m(y) F(y) dy = 0$

Thus  $\langle \Psi_n, \Psi_m \rangle = 0$ , and the eigenfunction is orthogonal.

When  $n = m$ ,  $\langle \Psi_n, \Psi_m \rangle = \int_{-\infty}^{\infty} H_n(y)^2 C_n^2 e^{-y^2} x_0 dy = \frac{x_0}{2^n n! x_0 \sqrt{\pi}} \int_{-\infty}^{\infty} H_n(y)^2 e^{-y^2} dy$

Since  $\int_{-\infty}^{\infty} H_n(y)^2 e^{-y^2} dy = 2^n n! \sqrt{\pi} \delta_{nn} = 2^n n! \sqrt{\pi}$

Then  $\langle \Psi_n, \Psi_n \rangle = C_n^2 2^n n! \sqrt{\pi} = 1$ , so each eigenfunction is normalized.

$\langle \Psi, \Psi \rangle = \langle \sqrt{\frac{1}{3}}\psi_3, \sqrt{\frac{1}{3}}\psi_3 \rangle + \langle \sqrt{\frac{1}{3}}\psi_3, \sqrt{\frac{2}{3}}\psi_5 \rangle + \langle \sqrt{\frac{2}{3}}\psi_5, \sqrt{\frac{1}{3}}\psi_3 \rangle + \langle \sqrt{\frac{2}{3}}\psi_5, \sqrt{\frac{2}{3}}\psi_5 \rangle$   
 $= \frac{1}{3} \langle \psi_3, \psi_3 \rangle + 0 + 0 + \frac{2}{3} \langle \psi_5, \psi_5 \rangle = \frac{1}{3} + \frac{2}{3} = 1$

So it's normalized

2.  $\Psi(x, 0) = \sqrt{\frac{1}{144 x_0 \sqrt{\pi}}} [8(\frac{x}{x_0})^3 - 12\frac{x}{x_0}] e^{-\frac{1}{2}(\frac{x}{x_0})^2} + \sqrt{\frac{1}{5760 x_0 \sqrt{\pi}}} [32(\frac{x}{x_0})^5 - 160(\frac{x}{x_0})^3 + 120\frac{x}{x_0}] e^{-\frac{1}{2}(\frac{x}{x_0})^2}$

where  $x_0 = \sqrt{\frac{\hbar}{m\omega}}$

3. The possible outcome of energy is  $E_3 = \frac{7}{2}\hbar\omega$ ,  $E_5 = \frac{11}{2}\hbar\omega$   
 The probability to get  $E_3$  is  $\frac{1}{3}$ , for  $E_5$  is  $\frac{2}{3}$

4.  $E = \frac{1}{3}E_3 + \frac{2}{3}E_5 = \frac{29}{6}\hbar\omega$

5.  $\Psi(x, t) = \sqrt{\frac{1}{3}} e^{\frac{i}{\hbar} E_3 t} \psi_3(x) + \sqrt{\frac{2}{3}} e^{\frac{i}{\hbar} E_5 t} \psi_5(x)$

## 5 Problem 5

(a) Let  $y = \frac{x}{x_0}$

When  $n = m$ ,  $\langle \Psi_n, \Psi_m \rangle = \int_{-\infty}^{\infty} H_n(y)^2 C_n^2 e^{-y^2} x_0 dy = \frac{x_0}{2^n n! x_0 \sqrt{\pi}} \int_{-\infty}^{\infty} H_n(y)^2 e^{-y^2} dy$

Since  $\int_{-\infty}^{\infty} H_n(y)^2 e^{-y^2} dy = 2^n n! \sqrt{\pi} \delta_{nn} = 2^n n! \sqrt{\pi}$

Then  $\langle \Psi_n, \Psi_n \rangle = C_n^2 2^n n! \sqrt{\pi} = 1$ , so each eigenfunction is normalized.

(b) According to text book:  $\int_{-\infty}^{\infty} \psi_n^* x \psi_m dx = 0$  unless  $n = m \pm 1$

Then  $\langle x \rangle = \int_{-\infty}^{\infty} \psi_n^* x \psi_n dx = 0$ , and  $\langle p \rangle = 0$

Therefore  $\Delta x = \sqrt{\langle x^2 \rangle} = C_1^2 \int_{-\infty}^{\infty} H_1(\frac{x}{x_0})^2 x^2 e^{-(\frac{x}{x_0})^2} dx$

$\Delta p = \sqrt{\langle p^2 \rangle} = C_1^2 \int_{-\infty}^{\infty} -\hbar H_1(\frac{x}{x_0}) e^{-\frac{1}{2}(\frac{x}{x_0})^2} \frac{\partial^2 H_1(\frac{x}{x_0}) e^{-\frac{1}{2}(\frac{x}{x_0})^2}}{\partial x^2} dx$

Then for the ground state  $\Delta x \Delta p = \frac{\hbar}{2}$

```

1000 a0 = 0.5/m;
1001 h =
1002 2*1/2*(2*1/2*a0+0.5)*0.5;
1003 0.751126
1004 2*1/2*(2*1/2*a0+0.5)*0.5
1005 0.751126
1006 2*1/2*(2*1/2*a0+0.5)*0.5
1007 0.751126
1008 2*1/2*(2*1/2*a0+0.5)*0.5
1009 0.751126
1010 2*1/2*(2*1/2*a0+0.5)*0.5
1011 0.751126
1012 2*1/2*(2*1/2*a0+0.5)*0.5
1013 0.751126
1014 2*1/2*(2*1/2*a0+0.5)*0.5
1015 0.751126
1016 2*1/2*(2*1/2*a0+0.5)*0.5
1017 0.751126
1018 2*1/2*(2*1/2*a0+0.5)*0.5
1019 0.751126
1020 2*1/2*(2*1/2*a0+0.5)*0.5
1021 0.751126
1022 2*1/2*(2*1/2*a0+0.5)*0.5
1023 0.751126
1024 2*1/2*(2*1/2*a0+0.5)*0.5
1025 0.751126
1026 2*1/2*(2*1/2*a0+0.5)*0.5
1027 0.751126
1028 2*1/2*(2*1/2*a0+0.5)*0.5
1029 0.751126
1030 2*1/2*(2*1/2*a0+0.5)*0.5
1031 0.751126
1032 2*1/2*(2*1/2*a0+0.5)*0.5
1033 0.751126
1034 2*1/2*(2*1/2*a0+0.5)*0.5
1035 0.751126
1036 2*1/2*(2*1/2*a0+0.5)*0.5
1037 0.751126
1038 2*1/2*(2*1/2*a0+0.5)*0.5
1039 0.751126
1040 2*1/2*(2*1/2*a0+0.5)*0.5
1041 0.751126
1042 2*1/2*(2*1/2*a0+0.5)*0.5
1043 0.751126
1044 2*1/2*(2*1/2*a0+0.5)*0.5
1045 0.751126
1046 2*1/2*(2*1/2*a0+0.5)*0.5
1047 0.751126
1048 2*1/2*(2*1/2*a0+0.5)*0.5
1049 0.751126
1050 2*1/2*(2*1/2*a0+0.5)*0.5
1051 0.751126
1052 2*1/2*(2*1/2*a0+0.5)*0.5
1053 0.751126
1054 2*1/2*(2*1/2*a0+0.5)*0.5
1055 0.751126
1056 2*1/2*(2*1/2*a0+0.5)*0.5
1057 0.751126
1058 2*1/2*(2*1/2*a0+0.5)*0.5
1059 0.751126
1060 2*1/2*(2*1/2*a0+0.5)*0.5
1061 0.751126
1062 2*1/2*(2*1/2*a0+0.5)*0.5
1063 0.751126
1064 2*1/2*(2*1/2*a0+0.5)*0.5
1065 0.751126
1066 2*1/2*(2*1/2*a0+0.5)*0.5
1067 0.751126
1068 2*1/2*(2*1/2*a0+0.5)*0.5
1069 0.751126
1070 2*1/2*(2*1/2*a0+0.5)*0.5
1071 0.751126
1072 2*1/2*(2*1/2*a0+0.5)*0.5
1073 0.751126
1074 2*1/2*(2*1/2*a0+0.5)*0.5
1075 0.751126
1076 2*1/2*(2*1/2*a0+0.5)*0.5
1077 0.751126
1078 2*1/2*(2*1/2*a0+0.5)*0.5
1079 0.751126
1080 2*1/2*(2*1/2*a0+0.5)*0.5
1081 0.751126
1082 2*1/2*(2*1/2*a0+0.5)*0.5
1083 0.751126
1084 2*1/2*(2*1/2*a0+0.5)*0.5
1085 0.751126
1086 2*1/2*(2*1/2*a0+0.5)*0.5
1087 0.751126
1088 2*1/2*(2*1/2*a0+0.5)*0.5
1089 0.751126
1090 2*1/2*(2*1/2*a0+0.5)*0.5
1091 0.751126
1092 2*1/2*(2*1/2*a0+0.5)*0.5
1093 0.751126
1094 2*1/2*(2*1/2*a0+0.5)*0.5
1095 0.751126
1096 2*1/2*(2*1/2*a0+0.5)*0.5
1097 0.751126
1098 2*1/2*(2*1/2*a0+0.5)*0.5
1099 0.751126
1100 2*1/2*(2*1/2*a0+0.5)*0.5
1101 0.751126
1102 2*1/2*(2*1/2*a0+0.5)*0.5
1103 0.751126
1104 2*1/2*(2*1/2*a0+0.5)*0.5
1105 0.751126
1106 2*1/2*(2*1/2*a0+0.5)*0.5
1107 0.751126
1108 2*1/2*(2*1/2*a0+0.5)*0.5
1109 0.751126
1110 2*1/2*(2*1/2*a0+0.5)*0.5
1111 0.751126
1112 2*1/2*(2*1/2*a0+0.5)*0.5
1113 0.751126
1114 2*1/2*(2*1/2*a0+0.5)*0.5
1115 0.751126
1116 2*1/2*(2*1/2*a0+0.5)*0.5
1117 0.751126
1118 2*1/2*(2*1/2*a0+0.5)*0.5
1119 0.751126
1120 2*1/2*(2*1/2*a0+0.5)*0.5
1121 0.751126
1122 2*1/2*(2*1/2*a0+0.5)*0.5
1123 0.751126
1124 2*1/2*(2*1/2*a0+0.5)*0.5
1125 0.751126
1126 2*1/2*(2*1/2*a0+0.5)*0.5
1127 0.751126
1128 2*1/2*(2*1/2*a0+0.5)*0.5
1129 0.751126
1130 2*1/2*(2*1/2*a0+0.5)*0.5
1131 0.751126
1132 2*1/2*(2*1/2*a0+0.5)*0.5
1133 0.751126
1134 2*1/2*(2*1/2*a0+0.5)*0.5
1135 0.751126
1136 2*1/2*(2*1/2*a0+0.5)*0.5
1137 0.751126
1138 2*1/2*(2*1/2*a0+0.5)*0.5
1139 0.751126
1140 2*1/2*(2*1/2*a0+0.5)*0.5
1141 0.751126
1142 2*1/2*(2*1/2*a0+0.5)*0.5
1143 0.751126
1144 2*1/2*(2*1/2*a0+0.5)*0.5
1145 0.751126
1146 2*1/2*(2*1/2*a0+0.5)*0.5
1147 0.751126
1148 2*1/2*(2*1/2*a0+0.5)*0.5
1149 0.751126
1150 2*1/2*(2*1/2*a0+0.5)*0.5
1151 0.751126
1152 2*1/2*(2*1/2*a0+0.5)*0.5
1153 0.751126
1154 2*1/2*(2*1/2*a0+0.5)*0.5
1155 0.751126
1156 2*1/2*(2*1/2*a0+0.5)*0.5
1157 0.751126
1158 2*1/2*(2*1/2*a0+0.5)*0.5
1159 0.751126
1160 2*1/2*(2*1/2*a0+0.5)*0.5
1161 0.751126
1162 2*1/2*(2*1/2*a0+0.5)*0.5
1163 0.751126
1164 2*1/2*(2*1/2*a0+0.5)*0.5
1165 0.751126
1166 2*1/2*(2*1/2*a0+0.5)*0.5
1167 0.751126
1168 2*1/2*(2*1/2*a0+0.5)*0.5
1169 0.751126
1170 2*1/2*(2*1/2*a0+0.5)*0.5
1171 0.751126
1172 2*1/2*(2*1/2*a0+0.5)*0.5
1173 0.751126
1174 2*1/2*(2*1/2*a0+0.5)*0.5
1175 0.751126
1176 2*1/2*(2*1/2*a0+0.5)*0.5
1177 0.751126
1178 2*1/2*(2*1/2*a0+0.5)*0.5
1179 0.751126
1180 2*1/2*(2*1/2*a0+0.5)*0.5
1181 0.751126
1182 2*1/2*(2*1/2*a0+0.5)*0.5
1183 0.751126
1184 2*1/2*(2*1/2*a0+0.5)*0.5
1185 0.751126
1186 2*1/2*(2*1/2*a0+0.5)*0.5
1187 0.751126
1188 2*1/2*(2*1/2*a0+0.5)*0.5
1189 0.751126
1190 2*1/2*(2*1/2*a0+0.5)*0.5
1191 0.751126
1192 2*1/2*(2*1/2*a0+0.5)*0.5
1193 0.751126
1194 2*1/2*(2*1/2*a0+0.5)*0.5
1195 0.751126
1196 2*1/2*(2*1/2*a0+0.5)*0.5
1197 0.751126
1198 2*1/2*(2*1/2*a0+0.5)*0.5
1199 0.751126
1200 2*1/2*(2*1/2*a0+0.5)*0.5

```

Figure 2: Mathematica Code for Integral Calculation

According to the Mathematica code in Figure.2 and test result for larger  $n$ , we know that  $\Delta x = \sqrt{\frac{\hbar}{2m\omega}}$ ,  $\Delta p = \sqrt{\frac{m\omega\hbar}{2}}$  and  $\Delta x \Delta p = \frac{\hbar}{2}$  for ground state.

$$\begin{aligned}
& \{ \{ \{ \{ \{ \{ P[n], \frac{0.751126 e^{-\frac{0.5 x^2}{\hbar/m}} \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]}{(2^n (\frac{\hbar}{m})^{0.5} n! )^{0.5}} \}, \left\{ \frac{0.751126 e^{-\frac{0.5 x^2}{\hbar/m}} \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]}{(2^n (\frac{\hbar}{m})^{0.5} n! )^{0.5}} \right\}^2, \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1}, x^2 \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1} \right\} \\
& \frac{x^2 0.56419 e^{-\frac{1 x^2}{\hbar/m}} \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1}, \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1}, \{ \{ \{ \{ \{ \{ \infty, \infty \}, -\infty, -\infty \}, \{ \infty, \infty \}, \{ x, -\infty, \infty \} \}, \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1} dx, \\
& \{ \{ \text{FreeQ}\left[ \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1}, \text{HoldPattern}[\text{Root}[\text{Integrate}[\text{ImproperDump}[\text{aa}\_\_ /; \text{Internal}[\text{DependsOnQ}[\{ \text{Integrate}[\text{ImproperDump}[\text{aa}\_\_], x]]], \text{True}], \right. \\
& \left. \{ \text{True}, \text{False} \}], \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1} dx \right\}, \left\{ \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1} dx \right\}^{0.5} \right\}, \\
& X[n\_]= \left\{ \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1} dx \right\}^{0.5}, \left\{ \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{\hbar/m}} x^2 \text{HermiteH}\left[n, \frac{x}{\hbar/m}\right]^2}{(2^n (\frac{\hbar}{m})^{0.5} n! )^1} dx \right\}^{0.5} \right\}
\end{aligned}$$

Figure 3: Calculation procedure for Mathematica to find  $\Delta x$

$$\begin{aligned}
&= \left\{ \left\{ \left\{ \left\{ \left\{ P[n], \frac{0.751126 e^{-\frac{0.5 x^2}{m \omega}} \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^{0.5}} \right\}, \left( \frac{0.751126 e^{-\frac{0.5 x^2}{m \omega}} \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^{0.5}} \right)^2, \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1}, x^2 \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} \right\}, \right. \\
&\quad \frac{x^2 0.56419 e^{-\frac{1 x^2}{m \omega}} \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1}, \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} \right\}, \left\{ \{\omega, \infty\}, -\omega, -\infty, \{\omega, \infty\}, \{x, -\infty, \infty\} \right\}, \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} dx, \\
&\quad \left\{ \text{FreeQ}\left[\frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1}, \text{HoldPattern}[\text{Root}[\text{Integrate}[\text{ImproperDump}[\text{aa}\_\_]; \text{Internal}[\text{DependsOnQ}[(\text{Integrate}[\text{ImproperDump}[\text{aa}\_\_], x]]], \text{True}]\right], \text{True} \right\}, \\
&\quad \text{! True, False} \right\}, \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} dx \right\}, \left( \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} dx \right)^{0.5} \right\}, \\
&\quad x[n\_]= \left( \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} dx \right)^{0.5}, \left( \int_{-\infty}^{\infty} \frac{0.56419 e^{-\frac{1 x^2}{m \omega}} x^2 \text{HermiteH}\left[n, \frac{x}{\left(\frac{h}{m \omega}\right)^{0.5}}\right]^2}{\left(2^n \left(\frac{h}{m \omega}\right)^{0.5} n!\right)^1} dx \right)^{0.5} \right\}
\end{aligned}$$

Figure 4: Calculation procedure for Mathematica to find  $\Delta p$

According to Figure.3 and Figure.4, we know that the value of  $\Delta x$  and  $\Delta p$  increase with  $n$ , hence their product increases with  $n$ , and  $\Delta x \Delta p \geq \frac{\hbar}{2}$

- (c)  $\hat{K} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2}$ , hence  $\bar{K} = \frac{(\Delta p)^2}{2m}$   
 $\hat{V} = V(x) = \frac{1}{2} m \omega^2 x^2$ , hence  $\bar{V} = \frac{m \omega^2}{2} (\Delta x)^2$   
Therefore  $\bar{K} \bar{V} = \frac{\omega^2}{4} (\Delta x \Delta p)^2 \geq \frac{\hbar \omega^2}{16}$   
When it's ground state, we plug the value of  $\Delta x, \Delta p$ :  
We get  $\bar{K} = \frac{\omega \hbar}{4}, \bar{V} = \frac{\omega \hbar}{4}$ , then  $\bar{K} = \bar{V} = \frac{E_0}{2}$

$$\begin{aligned}
\text{In}[26] &= \mathbf{X[1]} \\
\text{In}[4] &= \text{ConditionalExpression}\left[1.224744871391589 \cdot \left(\frac{1}{\left(\frac{h}{m \omega}\right)^{1.5} \left(\frac{m \omega}{h}\right)^{5/2}}\right)^{0.5}, \text{Re}\left[\frac{m \omega}{h}\right] > 0\right] \\
&\quad \mathbf{p[1]} \\
\text{Out}[4] &= \text{ConditionalExpression}\left[1.22474 \cdot \left(\frac{1}{\left(\frac{h}{m \omega}\right)^{1.5} \left(\frac{m \omega}{h}\right)^{5/2}}\right)^{0.5}, \text{Re}\left[\frac{m \omega}{h}\right] > 0\right]
\end{aligned}$$

Figure 5: The coefficients are same

According to Mathematica result in Figure.5 in question (b), we can find that the coefficients before  $\Delta x, \Delta p$  are same.

Therefore, for all  $n$ -eigenstates,  $\bar{K} = \bar{V} = \frac{E_0}{2}$

## 6 Problem 6

- (a) Assume the center of the cube as the original point. If the electron displacement is  $x$ , then we can imagine it's in a ball with radius  $x$ , with bound charge  $Q' = \frac{Q}{L^3} \frac{4\pi x^3}{3}$ , since the electric field caused by outside potential and charge is symmetric, we only need to calculate the electric field caused by internal charge for the oscillation model. According to Gauss Law:

$$4\pi x^2 E = \frac{Q'}{\epsilon_0} \rightarrow E = \frac{Qx}{3\epsilon_0 L^3}$$

$$\text{Then the spring constant } A = \frac{qE}{x} = \frac{Qe}{3\epsilon_0 L^3}$$

- (b) The original harmonic oscillator equation is  $-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x)}{\partial x^2} + \frac{m\omega^2 x^2}{2} \psi(x) = E\psi(x)$ , where  $V(x) = \frac{1}{2}m\omega^2 x^2$

$$\text{In our equation } V(x) = \frac{1}{2}m\omega^2 x^2 + \frac{e\Phi_0}{L}x = \frac{1}{2}m\omega^2 \left(x + \frac{e\Phi_0}{mL\omega^2}\right)^2 - \frac{e^2\Phi_0^2}{2mL^2\omega^2}$$

$$\text{Then our equation becomes: } -\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x)}{\partial x^2} + \frac{1}{2}m\omega^2 \left(x + \frac{e\Phi_0}{mL\omega^2}\right)^2 \psi(x) = \left(E + \frac{e^2\Phi_0^2}{2mL^2\omega^2}\right) \psi(x)$$

Assume the eigenfunction and eigenvalue in normal quantum oscillation is  $\psi_n(x), E_n$

Then in our case, the new eigenfunction and eigenvalue is :

$$\psi_n\left(x + \frac{e\Phi_0}{mL\omega^2}\right), E_n - \frac{e^2\Phi_0^2}{2mL^2\omega^2}, \text{ where } \omega = \sqrt{\frac{A}{m}}, A = \frac{qE}{x} = \frac{Qe}{3\epsilon_0 L^3}$$

Actually, it's just like we change the position of the oscillator