




PART II



Q2



Q3

(a) $E_1 = \epsilon$ $E_2 = 8\epsilon$ $E_8 = 512\epsilon$


$$\Psi(x,t) = \frac{\sqrt{2}}{5} e^{-\frac{i}{\hbar} \epsilon t} \psi_1 + \frac{\sqrt{2}}{5} e^{-\frac{i}{\hbar} 8\epsilon t} \psi_2 + \frac{1}{5} e^{-\frac{i}{\hbar} 512\epsilon t} \psi_8$$

(b) $\Pr(E = 8\epsilon = E_2) = C_2^2 = 0.4$

Q4 Heisenberg uncertainty principle claims that for any particle, $\Delta x \Delta p \geq \frac{\hbar}{2}$ so there is a trade-off between the particle's position and momentum. The more accurate we know about the particle's position, the less we know about its momentum, and vice versa.




Q5.



According to the structure of NH_3 , we know the arrangement of H is equilateral triangle and symmetric, the N atom also oscillates up and down symmetric


In the y-axis, which is N's direction, the potential is:



Since the N-atom can move between left and right half even when $E < V$ in QM! — is

In a double-well

For $\psi_{\text{left}} = \frac{1}{\sqrt{2}}(\psi_1 + \psi_2)$ $\psi_{\text{right}} = \frac{1}{\sqrt{2}}(\psi_1 - \psi_2)$ where ψ_1 is even ψ_2 is odd



ψ_{left} will change to ψ_{right} and change back between two configurations
Hence it can be modeled as 1D oscillation potential well

Q6

a) $\Pr\left[\left[0, \frac{L}{2}\right]\right] = \Pr\left[\left[-\frac{L}{2}, 0\right]\right] < \Pr\left[\left[\frac{L}{4}, \frac{3}{4}L\right]\right]$

b) $P = \frac{30}{L^5} \hbar \int_0^L x(x-L) \frac{\partial x(x-L)}{\partial x} dx = 0$

Q7 The lowest levels are $E_0 = \frac{3}{2}\hbar\omega$ and $E_1 = \frac{7}{2}\hbar\omega$

because since the left part of $V(x)$ is ∞ , the $\psi(x)=0$ for $x \leq 0$


and the energy level should be odd function.

Hence it should be $E_1 = \frac{3}{2}\hbar\omega$, $E_3 = \frac{7}{2}\hbar\omega$ in original oscillator model.



1. a) we just replace all $c = 3 \times 10^8$ m/s with $u_0 = 250$ m/s, because they're the max velocity for all objects

b) Yes because simultaneity is relative
since Thomas is moving, he can see Manuel and David's clock with different time



$$\Delta u_0 t = 250 \times 5 \times 60 = 75000 \text{ (m)}$$

$$\Delta x = 60000 \text{ m}$$

Then $\Delta s^2 = (\Delta u_0 t)^2 - (\Delta x)^2 > 0$, so the two instances are time like
Manuel is reasonable

d) Yes, if they're all not moving

Q2. a) $U_0 q = \frac{1}{2} m c^2$
 $U_0 = \frac{m c^2}{2q}$

b) Before acceleration, the particle has energy $E = m c^2$

After, $E = m c^2 + q U_0$, so $K = q U_0 = \frac{m c^2}{\sqrt{1 - u^2/c^2}} - m c^2$

then $u = c \sqrt{1 - \frac{(m c^2)^2}{(q U_0 + m c^2)^2}}$

Q3 For the stationary equation: $\hat{H} \psi(x) = E \psi(x)$

For $x \leq -a$, $\psi_1(x) = A e^{K_1 x}$ where $K_1^2 = -\frac{2m(E - V_1)}{\hbar^2}$

For $x \geq a$, $\psi_3(x) = A e^{K_3 x}$ where $K_3^2 = -\frac{2m(E - V_2)}{\hbar^2}$

For $-a < x < a$, $\psi_2(x) = C_1 \cos K_2 x + C_2 \sin K_2 x$ where $K_2^2 = \frac{2m}{\hbar^2} (E + V_0)$

$\psi_1(-a) = \psi_2(-a)$

$\psi_1'(-a) = \psi_2'(-a)$

$\psi_3(a) = \psi_2(a)$

$\psi_3'(a) = \psi_2'(a)$

$\int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1$



Q4:

$$a) \phi = \begin{cases} \frac{1}{a} & |x| < \frac{a}{2} \\ 0 & |x| > \frac{a}{2} \end{cases}$$

$$\psi = \sum C_n \psi_n \text{ where } C_n = \langle \psi_n, \phi \rangle$$

$$b) i) p = -i\hbar \frac{d}{dx} \left(\frac{1}{a} \right) = 0$$

ii)

$$Q5 a) \Delta p \Delta x = \sqrt{\langle p^2 \rangle_\psi - \langle p \rangle_\psi^2} \sqrt{\langle x^2 \rangle_\psi - \langle x \rangle_\psi^2}$$

$$\langle p^2 \rangle_\psi = \langle -i\hbar \frac{d}{dx} [\langle p \rangle_\psi + i \epsilon (x - \langle x \rangle_\psi)] \psi, \psi \rangle$$

$$\text{Then } \Delta p \Delta x \geq \frac{\hbar}{2}$$

b)

Q6 According to the diagram

$$\text{Since } (ka)^2 + (ka)^2 = \frac{2ma^2 V_0}{\hbar^2}$$

we can see the nodes of different states have different radii.

Since $E = \frac{k^2 \hbar^2}{2m} - V_0$, plug it back, we can get difference E_n



扫描全能王 创建



JOINT INSTITUTE
交大密西根学院

Pledge of Honor for Online Examinations Summer Term of 2020

The University of Michigan – Shanghai Jiao Tong University Joint Institute trusts its students to participate in examinations in an honorable and respectful manner, following a spirit of fairness and equality. Cheating, seeking unfair advantage and disturbing the safe and harmonious environment of examinations are contrary to the ethical principles of students of the Joint Institute. The letter and spirit of the Honor Code shall guide the behavior of students, faculty and all members of the Joint Institute.

Therefore, I hereby declare that

1. I will neither give nor receive unauthorized aid during the present examination, nor will I conceal any violations of the Honor Code by others or myself.
2. I confirm that I have read and understood the rules and procedures for examination set out by SJTU. I will follow them to the best of my ability.
3. I understand that violating the rules and procedures for examinations or the Honor Code will lead to administrative and/or academic sanctions.

Signature: 潘崇聃

Student ID: 516370910121

Date: 2020.6.22