

Homework 4 Solution

Problems:

1. (a) [2!] $x(t) = 2\text{rect}\left(\frac{t-2}{4}\right)$ so $X(\omega) = 8e^{-2j\omega}\text{sinc}\left(\frac{2\omega}{\pi}\right)$ by time scaling and shifting property.

(b) [2!] $x(t) = e^{-3t}\text{rect}\left(\frac{t-2}{4}\right) = e^{-3t}u(t) - e^{-3(t-4)}u(t-4)e^{-12}$ so $X(\omega) = \frac{1}{3+j\omega} [1 - e^{-12}e^{-4j\omega}]$

(c) [3!] $x(t) = t\text{rect}\left(\frac{t-2}{4}\right) = ty(t)$ where from part (a): $Y(\omega) = 4e^{-2j\omega}\text{sinc}\left(\frac{2\omega}{\pi}\right)$.

Thus $X(\omega) = \frac{d}{d\omega}jY(\omega) = \frac{d}{d\omega}j2e^{-2j\omega}\frac{\sin(2\omega)}{\omega} = \frac{e^{-4j\omega}(1+4j\omega) - 1}{\omega^2}$

(d) [3!] $x(t) = \cos(4\pi t)y(t)$ where from part (a): $Y(\omega) = 4e^{-2j\omega}\text{sinc}\left(\frac{2\omega}{\pi}\right)$.

Hence $X(\omega) = \frac{1}{2}(Y(\omega - 4\pi) + Y(\omega + 4\pi)) = 2e^{-2j\omega} \left(\text{sinc}\left(\frac{2\omega - 8\pi}{\pi}\right) + \text{sinc}\left(\frac{2\omega + 8\pi}{\pi}\right) \right)$

2. (a) [3!]

$$[e^{-at} \cos \omega_0 t]u(t) = \frac{1}{2}e^{-at}e^{j\omega_0 t}u(t) + \frac{1}{2}e^{-at}e^{-j\omega_0 t}u(t)$$

$$X(j\omega) = \frac{1}{2(\alpha - j\omega_0 + j\omega)} + \frac{1}{2(\alpha + j\omega_0 + j\omega)} = \frac{\alpha + j\omega}{(\alpha + j\omega)^2 + \omega_0^2}$$

(b) [3!]

$$x(t) = x_1(t) + x_2(t)$$

where,

$$x_1(t) = e^{-3t}\sin(2t)u(t) \xrightarrow{\mathcal{F}} X_1(j\omega) = \frac{1/(2j)}{3-j2+j\omega} - \frac{1/(2j)}{3+j2+j\omega} = \frac{2}{(3+j\omega)^2 + 4}$$

$$x_2(t) = e^{3t}\sin(2t)u(-t) \xrightarrow{\mathcal{F}} X_2(j\omega) = -X_1(-j\omega) = -\frac{1/(2j)}{3-j2-j\omega} + \frac{1/(2j)}{3+j2-j\omega} = \frac{-2}{(3-j\omega)^2 + 4}$$

Thus,

$$X(j\omega) = X_1(j\omega) + X_2(j\omega) = \frac{3j}{9 + (\omega + 2)^2} - \frac{3j}{9 + (\omega - 2)^2}$$

(c) [4!]

According to the FT table,

$$e^{-t^2/4} \xleftrightarrow{\mathcal{F}} 2\sqrt{\pi}e^{-\omega^2} = F_1(\omega)$$

and also

$$(-jt)^2 e^{-t^2/4} \xleftrightarrow{\mathcal{F}} \frac{d^2}{d\omega^2} F_1(\omega) = -4\sqrt{\pi}e^{-\omega^2}(1 - 2\omega^2)$$

Thus $F(\omega) = -\frac{d^2}{d\omega^2} F_1(\omega) = 4\sqrt{\pi}e^{-\omega^2}(1 - 2\omega^2)$

3. [7!]

(i) [4!] From the FT table, $\frac{1}{\pi} \text{rect}\left(\frac{t}{\pi}\right) \xleftrightarrow{\mathcal{F}} \text{sinc}\left(\frac{\omega}{2}\right)$ and $\frac{1}{\pi} \text{tri}\left(\frac{t}{\pi}\right) \xleftrightarrow{\mathcal{F}} \text{sinc}^2\left(\frac{\omega}{2}\right)$.Hence $F(\omega) = \text{sinc}^3(\omega/2) \xleftrightarrow{\mathcal{F}} \frac{1}{\pi} \text{rect}\left(\frac{t}{\pi}\right) * \frac{1}{\pi} \text{tri}\left(\frac{t}{\pi}\right)$.

(ii) [3!] By using graphical convolution,

$$\text{rect}\left(\frac{t}{\pi}\right) * \text{tri}\left(\frac{t}{\pi}\right) = \begin{cases} \frac{1}{2\pi} \left(\frac{3}{2}\pi - t\right)^2 & \frac{1}{2}\pi < t < \frac{3}{2}\pi \\ \frac{1}{2\pi} \left(\frac{3}{2}\pi + t\right)^2 & -\frac{3}{2}\pi < t < -\frac{1}{2}\pi \\ \frac{3}{4}\pi - \frac{t^2}{\pi} & -\frac{1}{2}\pi < t < \frac{1}{2}\pi \\ 0 & \text{otherwise} \end{cases}$$

and it follows that

$$f(t) = \begin{cases} \frac{1}{2\pi^3} \left(\frac{3}{2}\pi - |t|\right)^2 & \frac{1}{2}\pi < |t| < \frac{3}{2}\pi \\ \frac{3}{4\pi} - \frac{t^2}{\pi^3} & -\frac{1}{2}\pi < t < \frac{1}{2}\pi \\ 0 & \text{otherwise} \end{cases}$$

4. [8!]

$$f_e(t) = \frac{1}{2}[f(t) + f(-t)] \xleftrightarrow{\mathcal{F}} F_e(\omega) = \frac{1}{2}[F(\omega) + F(-\omega)]$$

$$f_o(t) = \frac{1}{2}[f(t) - f(-t)] \xleftrightarrow{\mathcal{F}} F_o(\omega) = \frac{1}{2}[F(\omega) - F(-\omega)]$$

Since $f(t)$ is real, $F(-\omega) = F^*(-\omega)$, and it follows that

$$f_e(t) \xleftrightarrow{\mathcal{F}} F_e(\omega) = \frac{1}{2}[F(\omega) + F^*(\omega)] = \text{real}\{F(\omega)\}$$

$$f_o(t) \xleftrightarrow{\mathcal{F}} F_o(\omega) = \frac{1}{2}[F(\omega) - F^*(\omega)] = j\text{imag}\{F(\omega)\}$$

5. [10!]

i) Method 1. The generator signal(the time-limited signal when $n=0$) is

$$g(t) = x(t)\text{rect}(t/6) = 2\delta(t) - \delta(t-2) - \delta(t+2)$$

so its FT is

$$G(\omega) = 2 - e^{j2\omega} - e^{-j2\omega} = 2(1 - \cos 2\omega)$$

Thus the FT of the period signal $x(t)$ is

$$X(\omega) = \sum_{k=-\infty}^{\infty} c_k 2\pi \delta(\omega - k\omega_0)$$

$$\begin{aligned}
&= \sum_{k=-\infty}^{\infty} \frac{1}{T_0} G(\omega)|_{\omega=k\omega_0} 2\pi \delta(\omega - k\omega_0) \\
&= \sum_{k=-\infty}^{\infty} \omega_0 G(k\omega_0) \delta(\omega - k\omega_0) \\
&= \sum_{k=-\infty}^{\infty} (\pi/3) 2(1 - \cos 2k\pi/3) \delta(\omega - k\pi/3) \\
&= \sum_{k=-\infty}^{\infty} a_k \delta(\omega - k\pi/3)
\end{aligned}$$

where $a_0 = 0, a_{\pm 1} = \pi, a_{\pm 2} = \pi, a_{\pm 3} = 0 \dots$

ii) Method 2.

$$x(t) = 2x_1(t) - x_2(t) - x_3(t)$$

$$x_1(t) = \sum_{n=-\infty}^{\infty} \delta(t - T_0 n)$$

$$x_2(t) = \sum_{n=-\infty}^{\infty} \delta(t - T_0 n - 2)$$

$$x_3(t) = \sum_{n=-\infty}^{\infty} \delta(t - T_0 n + 2)$$

Where $T_0 = 6$. The FS of $x_1(t)$ is (See Lecture 11, p91-p94)

$$x_1(t) = \sum_{k=-\infty}^{\infty} \frac{1}{T_0} e^{jk\omega_0 t}$$

And by time shift property of FS we can get,

$$x_2(t) = \sum_{k=-\infty}^{\infty} \frac{1}{T_0} e^{-jk\omega_0 2} e^{jk\omega_0 t}$$

$$x_3(t) = \sum_{k=-\infty}^{\infty} \frac{1}{T_0} e^{jk\omega_0 2} e^{jk\omega_0 t}$$

By linear property of FS,

$$\begin{aligned}
x(t) &= 2 \sum_{k=-\infty}^{\infty} \frac{1}{T_0} e^{jk\omega_0 t} - \sum_{k=-\infty}^{\infty} \frac{1}{T_0} e^{-jk\omega_0 2} e^{jk\omega_0 t} - \sum_{k=-\infty}^{\infty} \frac{1}{T_0} e^{jk\omega_0 2} e^{jk\omega_0 t} \\
&= \sum_{k=-\infty}^{\infty} \frac{2}{T_0} (1 - \cos(k\omega_0 2)) e^{jk\omega_0 t}
\end{aligned}$$

The FT of $x(t)$ is

$$x(t) \xrightarrow{\mathcal{F}} X(\omega) = \sum_{k=-\infty}^{\infty} \frac{2}{T_0} (1 - \cos(k\omega_0 2)) \delta(\omega - k\omega_0)$$

$$= \sum_{k=-\infty}^{\infty} (\pi/3) 2(1 - \cos 2k\pi/3) \delta(\omega - k\pi/3)$$

$$= \sum_{k=-\infty}^{\infty} a_k \delta(\omega - k\pi/3)$$

where $a_0 = 0, a_{\pm 1} = \pi, a_{\pm 2} = \pi, a_{\pm 3} = 0 \dots$




Figure 1: Problem 5

6. [10!]

(a) [4!]

$$X(j\omega) = \frac{e^{(4\omega + \pi/3)j} + e^{-(4\omega + \pi/3)j}}{2}$$

$$x(t) = \frac{1}{2} e^{-j\pi t/12} \delta(t+4) + \frac{1}{2} e^{j\pi t/12} \delta(t-4)$$

(b) [6!] From the given figure we can get

$$X(j\omega) = \begin{cases} \omega e^{-3j\omega} & , |\omega| < 1 \\ 0 & , |\omega| > 1 \end{cases}$$

$$x(t) = \frac{\cos(t-3) - 1 - (t-3)\sin(t-3)}{\pi(t-3)^2}$$

7. [15!] From the figure we can get the frequency response of the lowpass differentiator is,

$$H(j\omega) = \begin{cases} \frac{j\omega}{3\pi} & , -3\pi \leq \omega \leq 3\pi \\ 0 & , \text{otherwise} \end{cases}$$

(a) Since $x(t) = \cos(2\pi t + \theta)$, $X(j\omega) = e^{-j\omega\theta/2\pi} [\delta(\omega - 2\pi) + \delta(\omega + 2\pi)]$. It is zero outside the region $-3\pi \leq \omega \leq 3\pi$. Thus,

$$Y(j\omega) = X(j\omega)H(j\omega) = \frac{j\omega}{3\pi} X(j\omega)$$

$$y(t) = \frac{1}{3\pi} \frac{dx(t)}{dt} = -\frac{2}{3} \sin(2\pi t + \theta)$$

- (b) Since $x(t) = \cos(4\pi t + \theta)$, $X(j\omega) = e^{-j\omega\theta/4\pi} \pi [\delta(\omega - 4\pi) + \delta(\omega + 4\pi)]$. The nonzero portions of $X(j\omega)$ lie outside the region $-3\pi \leq \omega \leq 3\pi$. This implies,

$$Y(j\omega) = X(j\omega)H(j\omega) = 0$$

$$y(t) = 0$$

- (c) The Fourier series coefficients of the signal $x(t)$ are given by

$$c_k = \frac{1}{1} \int_0^{0.5} \sin(2\pi t) e^{-jk2\pi t} dt$$

Also we have

$$X(j\omega) = 2\pi \sum_{k=-\infty}^{\infty} a_k \delta(\omega - k\omega_0)$$

Where $\omega_0 = 2\pi$. Thus, we can see that only the portions when $k = 0, \pm 1$ lie in the region $-3\pi \leq \omega \leq 3\pi$.

$$c_0 = \frac{1}{\pi}, c_1 = c_{-1}^* = \frac{-1}{4j}$$

The portion that can pass the filter is $x_{lp}(t) = \frac{1}{\pi} + \frac{1}{2} \sin(2\pi t)$. Finally, $y(t) = \frac{1}{3\pi} \frac{dx_{lp}(t)}{dt} = \frac{1}{3} \cos(2\pi t)$

8. [10!]

$$x(t) = t \operatorname{sinc}^2(t) \xleftrightarrow{\mathcal{F}} X(\omega) = j \frac{d}{d\omega} \operatorname{tri}\left(\frac{\omega}{2\pi}\right) = \pm j \frac{1}{2\pi} \operatorname{rect}\left(\frac{\omega}{4\pi}\right)$$

$$\text{So } E = \frac{1}{2\pi} \int_{-\infty}^{\infty} |X(\omega)|^2 d\omega = \frac{1}{2\pi} \int_{-2\pi}^{2\pi} \frac{1}{4\pi^2} d\omega = \boxed{\frac{1}{2\pi^2}}$$

9. (a) [10!]

$$H(s) = \frac{s^2 + s + 1}{(s^2 + 6s + 25)(s^2 + 2)} = \frac{s^2 + s + 1}{(s+2)(s+3-4j)(s+3+4j)} = \frac{r_1}{s+2} + \frac{r_2}{s+3-4j} + \frac{r_2^*}{s+3+4j}$$

$$\text{with } r_1 = \left. \frac{s^2 + s + 1}{s^2 + 6s + 25} \right|_{s=-2} = \frac{3}{17} \text{ and } r_2 = \left. \frac{s^2 + s + 1}{(s+3+4j)(s+2)} \right|_{s=-3+4j} = \frac{7}{17} + \frac{71}{136}j = 0.665e^{j0.903}.$$

Thus

$$H(j\omega) = \frac{3/17}{j\omega + 2} + \frac{0.665e^{j0.903}}{j\omega + 3 - 4j} + \frac{0.665e^{-j0.903}}{j\omega + 3 + 4j}$$

Taking the inverse FT yields

$$h(t) = \frac{3}{17} e^{-2t} u(t) + 1.33e^{-3t} \cos(4t + 0.903) u(t)$$

(b) [10!] Expanding the denominator polynomial:

$$\frac{Y(\omega)}{X(\omega)} = \frac{(j\omega)^2 + j\omega + 1}{(j\omega)^3 + 8(j\omega)^2 + 37j\omega + 50}$$

so by cross multiplying

$$50y(t) + 37\frac{d}{dt}y(t) + 8\frac{d^2}{dt^2}y(t) + \frac{d^3}{dt^3}y(t) = x(t) + \frac{d}{dt}x(t) + \frac{d^2}{dt^2}x(t)$$