

ALEXANDRIA UNIVERSITY

FACULTY OF BUSINESS

STATISTICAL PACKAGES 2026

Driving Data Analysis

TEAM: PEAKY BLINDERS CODE*Supervised by: Dr. Assem Fareed*★ **Mohamed Essam Mohamed Ibrahim**

• Aya Ahmed Mohamed Hussein

• Mohamed Ahmed Almaz

• Ahmed Samy Mohamed Abu Bakr

• Ziad Ateyya Ateyya

• Mohamed Alaa El-Din Hassan

• Malek Osama Mohamed Mahmoud

• Youssef Mostafa Mohamed Abd El-Wahab

• Mohamed Saad Abady Hamid

• Mazin Hassan Ali Hassan Ahmed

• Mark Mina Sidhom Farid

• Mahmoud Mohamed Mohamed
Ahmed■ **Dataset:** wooldridge::driving■ **Year:** 2026■ **Course:** Statistical Packages

TABLE OF CONTENTS

PART 1 — Descriptive Statistics

- 1.1 Central Tendency – Mean & Median
- 1.2 Variability – Standard Deviation, Range & IQR
- 1.3 Distribution Shape & Skewness
- 1.4 Outlier Detection (IQR Method)
- 1.5 Group Comparison: Seatbelt Usage
- 1.6 Visualizations: Histogram & Boxplot

PART 2 — One-Sample Hypothesis Test ($\mu = 2$)

- 2.1 Hypotheses Statement
- 2.2 Test Procedure & Results
- 2.3 Conclusion at 5% Significance

PART 3 — Two-Sample Hypothesis Test (Seatbelt Groups)

- 3.1 Hypotheses & Welch t-Test
- 3.2 Test Results & p-value
- 3.3 Practical Importance of Findings

PART 4 — Economic Categorical Analysis (ANOVA)

- 4.1 Income-Level Categorization
- 4.2 Group Means & Hypothesis Testing
- 4.3 Group Differences & Policy Implications

PART 5 — Correlation & Multiple Regression Analysis

- 5.1 Scatter Plots & Correlations
- 5.2 Simple Linear Regression
- 5.3 Multiple Regression Model
- 5.4 Model Diagnostics & Residual Analysis

FINAL — Conclusion & Policy Recommendations

DATASET OVERVIEW: wooldridge::driving

The **driving** dataset from the Wooldridge package is a rich, balanced panel dataset covering U.S. state-level traffic safety outcomes and policy variables. It spans multiple years and all 50 states, providing a powerful foundation for both cross-sectional and longitudinal analysis. This report focuses on the **totfatrte** variable (total fatality rate per 100 million vehicle miles driven) as the primary outcome variable of interest, while leveraging **seatbelt** usage, **unem** (unemployment rate), and **vehicmilespc** (vehicle miles per capita) as predictors.

Variable	Description	Role in Analysis
totfatrte	Total traffic fatality rate (per 100M VMT)	Primary outcome variable
seatbelt	Seatbelt usage indicator (0=no law, 1=secondary, 2=primary)	Key safety predictor / grouping variable
unem	State unemployment rate (%)	Economic control variable
vehicmilespc	Vehicle miles driven per capita	Economic activity proxy / income proxy

The dataset contains approximately **1,200 observations** (50 states × ~24 years), giving sufficient statistical power to detect meaningful relationships and validate hypothesis tests at the 5% significance level.

PART 1 DESCRIPTIVE STATISTICS

1.1 Central Tendency

Central tendency measures allow us to identify a 'typical' observation. For the **totfatrte** variable, R yields the following:

```
# Load data
library(wooldridge)
data('driving'); df <- driving

# Central tendency
avg_accidents <- mean(df$totfatrte, na.rm = TRUE)
med_accidents <- median(df$totfatrte, na.rm = TRUE)
```

```
Mean Accident Rate: 18.91856
Median Accident Rate: 18.43500
```

Statistic	Value
Mean Fatality Rate	18.919
Median Fatality Rate	18.435
Difference (Mean – Median)	+0.484

Which measure is more representative?

The **median (18.44)** is the more representative measure of central tendency for this dataset. The mean (18.92) is higher than the median — a diagnostic signal of **positive (right) skew**. A small number of states with catastrophically high fatality rates (some exceeding 50 per 100M VMT) pull the arithmetic mean upward. The median, being resistant to extreme values, gives a more accurate picture of what a 'typical' state experiences. This is a fundamental principle: in skewed distributions or in the presence of outliers, the median outperforms the mean as a representative summary.

1.2 Measures of Variability

Variability measures quantify how spread out the data is around its central value. Multiple dispersion measures are computed to capture both typical and extreme spread.

```
sd_accidents <- sd(df$totfatrte, na.rm = TRUE)
min_val <- min(df$totfatrte, na.rm = TRUE)
max_val <- max(df$totfatrte, na.rm = TRUE)
Q1 <- quantile(df$totfatrte, 0.25, na.rm = TRUE)
Q3 <- quantile(df$totfatrte, 0.75, na.rm = TRUE)
IQR_val <- Q3 - Q1
```

```
Standard Deviation: 6.367407
Range: 6.2 to 53.32 (Span = 47.12)
Q1 (25th pctl): ~14.35
Q3 (75th pctl): ~22.52
IQR: ~8.17
```

Measure of Dispersion	Value
Standard Deviation (σ)	6.367
Range (Max – Min)	47.12 (6.20 → 53.32)
Interquartile Range (IQR)	≈ 8.17
Coefficient of Variation (CV)	33.65%

The **standard deviation of 6.37** implies that fatality rates typically deviate by about 6.4 units from the mean — a substantial relative spread given that the mean is only ~19. The **range of 47.12** is enormous and signals the presence of extreme outliers. The **IQR of ~8.17** is a more robust measure of spread that is unaffected by outliers; it confirms that the middle 50% of states fall within an 8-unit band. The **Coefficient of Variation (CV = $\sigma/\mu \times 100 \approx 33.65\%$)** indicates high relative variability — nearly one-third of the mean — suggesting significant heterogeneity across states and time periods.

1.3 Distribution Shape & Skewness

```
# Mean > Median → Right (Positive) Skew
shape <- ifelse(avg_accidents > med_accidents,
  'Positively Skewed (Right-Skewed)',
  'Negatively Skewed (Left-Skewed)')
```

```
The distribution is: Positively Skewed (Right-Skewed)
```

Finding: The fatality rate distribution is **Positively Skewed (Right-Skewed)**.

Since Mean (18.92) > Median (18.44), the distribution has a longer tail on the right. This is consistent with real-world accident data: most states in most years have moderate fatality rates clustered between 10 and 25, but a small subset of states (particularly rural, low-density states in earlier years) experienced catastrophic fatality rates exceeding 35–50. These extreme values stretch the right tail. Policy implication: risk is not uniformly distributed — targeted interventions in high-fatality outlier states could disproportionately improve national averages.

1.4 Outlier Detection — The IQR Method

The IQR method identifies outliers as values lying more than $1.5 \times \text{IQR}$ below Q1 or above Q3. This is a non-parametric, robust method that does not assume normality.

```
lower_bound <- Q1 - 1.5 * IQR_val
upper_bound <- Q3 + 1.5 * IQR_val
outliers <- df$totfatrtte[df$totfatrtte < lower_bound | df$totfatrtte > upper_bound]
print(outliers)
```

```
Outliers detected (rates above 35.365 ):
[1] 41.31 42.07 43.22 46.51 42.31 38.08 37.20 38.42
[9] 35.77 52.18 53.32 39.69 38.43 37.67
```

Parameter	Value
Lower Fence ($Q1 - 1.5 \times IQR$)	≈ 6.19 (no lower outliers)
Upper Fence ($Q3 + 1.5 \times IQR$)	35.365
Number of Outliers Detected	14 observations
Most Extreme Value	53.32 ($>2.8 \times$ the median!)

Impact of Outliers: These 14 data points — representing state-years with fatality rates between 36 and 53 — have a compounding effect on statistical analysis: (1) They inflate the **mean** upward by ~ 0.5 units relative to the median. (2) They dramatically increase the **standard deviation** and range, overstating typical variability. (3) In regression models, these high-leverage points can **distort coefficient estimates**, particularly the intercept. (4) These outliers correspond primarily to rural states in the 1980s–early 1990s, before federal seatbelt mandates and highway safety improvements took effect — they are real phenomena, not data errors, and represent a historically critical era in American traffic safety.

1.5 Group Comparison: Higher vs. Lower Seatbelt Usage

The dataset is split into two groups based on whether a state's seatbelt usage measure exceeds the median. This creates a natural experiment to test whether compliance with seatbelt laws correlates with lower fatality rates.

```
med_seatbelt <- median(df$seatbelt, na.rm = TRUE)
df$seatbelt_group <- ifelse(df$seatbelt > med_seatbelt,
  'Higher Usage', 'Lower Usage')
comparison_mean <- aggregate(totfatrtte ~ seatbelt_group, data = df, mean)
comparison_sd <- aggregate(totfatrtte ~ seatbelt_group, data = df, sd)
```

```
seatbelt_group totfatrtte (mean)
1 Higher Usage 18.19397
2 Lower Usage 19.55683
seatbelt_group totfatrtte (sd)
1 Higher Usage 5.707669
2 Lower Usage 6.837558
```

Metric	Higher Seatbelt Usage	Lower Seatbelt Usage	Difference
Mean Fatality Rate	18.194	19.557	-1.363 ↓
Standard Deviation	5.708	6.838	-1.130 ↓
CV ($\sigma / \mu \times 100$)	31.37%	34.96%	-3.59 pp ↓

Interpretation: The Higher Seatbelt Usage group consistently outperforms the Lower Usage group across all three metrics. It has a lower mean fatality rate (18.19 vs 19.56), a smaller standard deviation (5.71 vs 6.84), and a lower Coefficient of Variation (31.4% vs 35.0%). The lower CV confirms that states with stronger seatbelt compliance are not only safer on average but also **more stable and predictable** in their safety performance — a crucial quality for long-term policy planning.

1.6 Visualizations: Histogram & Boxplot

The histogram and boxplot together provide complementary graphical evidence of the distribution's shape, central tendency, spread, and outliers.




Figure 1 — Left: Histogram of `totfatrte` showing right-skewed distribution with Mean (red solid) slightly above Median (yellow dashed). Right: Boxplot confirming heavy upper-tail outliers beyond the upper fence at 35.365.

Histogram Insights: The bars peak sharply between 10–20, confirming the dense cluster of typical observations. The elongated right tail (bars stretching to 50+) visually demonstrates positive skewness. The red mean line sits measurably to the right of the yellow median line — a direct graphical confirmation of skew.

Boxplot Insights: The green box (IQR) is compact relative to the whiskers, and the dots above the upper whisker (≈ 35.4) are the identified outliers. The median line within the box is roughly centered, but the upper whisker is notably longer than the lower one, again confirming asymmetry toward high values.

PART 2 ONE-SAMPLE HYPOTHESIS TEST (Claim: $\mu = 2$)

Objective & Context

The project states the claim that the average accident rate equals 2. Given our sample mean of ~18.92, this claim appears obviously false — but we must formally test it using the rigorous framework of inferential statistics to provide a defensible, quantitative conclusion.

2.1 Hypotheses Statement

Null Hypothesis (H_0):	$\mu = 2$ (The true population mean fatality rate equals 2)
Alt. Hypothesis (H_1):	$\mu \neq 2$ (The true population mean fatality rate is NOT 2)
Test Type:	Two-tailed one-sample t-test
Significance Level:	$\alpha = 0.05$ (5%)

2.2 Test Procedure & Results

Using the one-sample t-test formula: $t = (x_{\text{bar}} - \mu) / (s / \sqrt{n})$, with $x_{\text{bar}} = 18.919$, $\mu = 2$, $s = 6.367$, and $n \approx 1,200$:

```
# One-sample t-test against mu = 2
t.test(df$totfatrte, mu = 2)
```

Parameter	Value
Sample Mean ($\bar{x}$)	18.919
Hypothesized Mean (μ)	2.000
Test Statistic (t)	$\approx +92.0$
Degrees of Freedom	$\approx 1,199$
p-value	< 0.0001 (virtually zero)
95% CI for μ	approximately [18.56, 19.28]

CONCLUSION: At $\alpha = 0.05$, the p-value (< 0.0001) is vastly smaller than the significance level. We **STRONGLY REJECT H_0** . There is overwhelming statistical evidence that the true mean fatality rate is not 2 — it is approximately 18.9. A t-statistic of ~92 is extraordinary; the claimed value of 2 is nearly 17 standard errors below the observed mean.

PART 3 TWO-SAMPLE HYPOTHESIS TEST (Seatbelt Groups)

Objective

We formally test whether the difference in mean fatality rates between the 'Higher Seatbelt Usage' and 'Lower Seatbelt Usage' groups is statistically significant, or could plausibly be due to random sampling variation.

3.1 Hypotheses

H₀:	$\mu(\text{Higher}) = \mu(\text{Lower})$ — No difference in mean fatality rates
H_a:	$\mu(\text{Higher}) \neq \mu(\text{Lower})$ — A significant difference exists
Test:	Welch Two-Sample t-test (does not assume equal variances)

3.2 Test Results

```
t_test_results <- t.test(totfatrte ~ seatbelt_group, data = df)
print(t_test_results)
```

```
Welch Two Sample t-test
data: totfatrte by seatbelt_group
t = -3.7619, df = 1194.6, p-value = 0.0001768
alternative hypothesis: true difference in means != 0
95% CI: [-2.0736, -0.6521]
mean in group Higher Usage: 18.19397
mean in group Lower Usage: 19.55683
```

Result Parameter	Value
Test Statistic (t)	-3.7619
Degrees of Freedom	1,194.6 (Welch adjusted)
p-value	0.0001768
95% Confidence Interval	[-2.074, -0.652]
Mean Difference (Higher – Lower)	-1.363 fatalities / 100M VMT

3.3 Conclusion & Practical Importance

STATISTICAL CONCLUSION: $p = 0.000177 < \alpha = 0.05 \rightarrow \text{REJECT } H_0$. The difference in mean fatality rates is statistically significant. States/years with higher seatbelt compliance average 1.36 fewer fatalities per 100 million vehicle miles driven than those with lower compliance. PRACTICAL SIGNIFICANCE: With ~3 trillion vehicle miles driven in the U.S. annually, a reduction of 1.36 per 100M VMT translates to approximately 40,800 fewer fatalities per year — a finding of enormous public health and policy importance. This strongly validates the enforcement of mandatory primary seatbelt laws across all states.

PART 4 CATEGORICAL ECONOMIC ANALYSIS (ANOVA Framework)

A new categorical variable is constructed from **vehicmilespc** (vehicle miles per capita), used as a proxy for economic activity and income level. States are divided into terciles: Low, Medium, and High activity.

```
breaks <- unique(quantile(df$vehicmilespc,
  probs = c(0, 0.33, 0.66, 1), na.rm = TRUE))
df$activity_level <- cut(df$vehicmilespc,
  breaks = breaks,
  labels = c('Low', 'Medium', 'High'),
  include.lowest = TRUE)
econ_table <- table(df$activity_level, df$seatbelt_group)
income_mean <- aggregate(totfatrtte ~ activity_level, data = df, mean)
```

Contingency Table (Activity Level vs Seatbelt Group):

Higher Usage Lower Usage

Low Income/Activity 89 307

Medium Income/Activity 196 200

High Income/Activity 277 131

Mean Accident Rate by Economic Level:

Low Income/Activity 17.78008

Medium Income/Activity 18.29144

High Income/Activity 20.63223

4.2 Group Means & Hypothesis Testing (ANOVA)

Income/Activity Level	Mean Fatality Rate	Seatbelt Usage Pattern
Low Income/Activity	17.780	Predominantly Lower Usage (89 vs 307)
Medium Income/Activity	18.291	Roughly Equal (196 vs 200)
High Income/Activity	20.632 ■	Predominantly Higher Usage (277 vs 131)

H₀:	$\mu(\text{Low}) = \mu(\text{Medium}) = \mu(\text{High})$ — All group means are equal
H_a:	At least one group mean is significantly different from the others

The mean differences (Low: 17.78, Medium: 18.29, High: 20.63) are substantial, especially for the High group, which averages **2.85 more fatalities** than the Low group. An ANOVA test on this large sample ($n \approx 1,200$) confirms these differences are statistically significant (F-statistic $\gg 3.0$, $p < 0.05$). We **reject H₀**.

4.3 Visualization: Boxplot by Economic Activity Level




Figure 2 — Boxplot of fatality rates by economic activity level. The High activity group shows a visibly elevated median and wider upper quartile. Note the presence of extreme outliers in all three groups.

Policy Implication (Paradox of Economic Activity): The High Economic Activity group paradoxically has the *highest* fatality rate despite also having the most seatbelt users. This is a classic **exposure effect**: more miles driven = more opportunities for fatal accidents, even if individual trips are safer. Additionally, the contingency table reveals that low-activity states historically had very low seatbelt compliance, which, combined with their rural, high-speed road networks, may have compressed their fatality averages differently. This highlights the need for multivariate analysis — no single variable tells the full story.

PART 5 CORRELATION & REGRESSION ANALYSIS

5.1 Scatter Plots & Correlation Analysis

```
cor_sb <- cor(df$seatbelt, df$totfatrte, use = 'complete.obs')
cor_unem <- cor(df$unem, df$totfatrte, use = 'complete.obs')
```

```
Correlation (Seatbelt, Fatality): -0.1935476
Correlation (Unemployment, Fatality): +0.2067008
```

Variable Pair	Pearson r	Strength	Direction
Seatbelt ↔ Fatality Rate	-0.194	Weak–Moderate	Negative (inverse)
Unemployment ↔ Fatality Rate	+0.207	Weak–Moderate	Positive (direct)




Figure 3 — Scatterplot: Seatbelt Usage Measure vs. Fatality Rate. The seatbelt variable is coded 0/1/2 (no law / secondary / primary enforcement). The red regression line confirms a downward trend — higher enforcement is associated with lower fatality rates. Wide vertical spread reflects within-level variability across states and years.

Interpreting the Scatterplot: The discrete nature of the seatbelt variable (values 0, 1, 2) creates three vertical clusters rather than a classic scatter. Within each cluster, enormous vertical spread indicates that other factors beyond seatbelt law type heavily influence fatality rates. Nevertheless, the regression line's negative slope visually confirms the negative correlation: as seatbelt enforcement intensity increases from 0 (no law) to 2 (primary enforcement), the expected fatality rate decreases. The 95% prediction band (implied by the spread) is wide, foreshadowing the modest R^2 in the regression models.

5.2 Simple Linear Regression: Seatbelt → Fatality Rate

```
model_simple <- lm(totfatrte ~ seatbelt, data = df)
summary(model_simple)$coefficients
```

```
Estimate Std. Error t value Pr(>|t|)
(Intercept) 20.448307 0.2876440 71.088936 0.000000e+00
seatbelt -1.370947 0.2007769 -6.828212 1.363501e-11
```

Simple Regression Parameter	Value / Interpretation
Intercept (β_0)	20.448 ($p < 0.0001$)
Seatbelt Coefficient (β_1)	-1.371 ($p < 0.0001$)
t-statistic for seatbelt	-6.83
Interpretation	Each 1-level increase in seatbelt law → -1.37 fatalities

The simple model tells us: when a state has no seatbelt law (seatbelt = 0), the baseline predicted fatality rate is **20.45**. Each step up in seatbelt law strength (from no law → secondary law → primary law) reduces the predicted rate by **1.37 per 100M VMT**. This effect is **highly statistically significant** ($t = -6.83$, $p < 0.0001$). However, the simple model omits critical confounders such as economic conditions, which necessitates the multiple regression model.

5.3 Multiple Regression: Seatbelt + Unemployment

```
model_multi <- lm(totfatrte ~ seatbelt + unem, data = df)
summary(model_multi)$coefficients
```

```
Estimate Std. Error t value Pr(>|t|)
(Intercept) 17.1549331 0.71880498 23.865907 2.739498e-103
seatbelt -0.9289674 0.21764058 -4.268356 2.124765e-05
unem 0.4705227 0.09428954 4.990189 6.923785e-07
```

Coefficient	Estimate	Std. Error	t-value	p-value	Significant?
Intercept (β_0)	17.155	0.719	23.87	< 0.0001	✓ YES
Seatbelt (β_1)	-0.929	0.218	-4.27	0.0000212	✓ YES
Unemployment (β_2)	+0.471	0.094	+4.99	0.00000069	✓ YES

$$\text{Estimated Model: Fatality Rate} = 17.155 - 0.929 \times \text{Seatbelt} + 0.471 \times \text{Unemployment}$$

Coefficient Interpretation:

Intercept (17.155): The predicted fatality rate when seatbelt = 0 (no law) and unemployment = 0. Statistically this is a baseline anchor.

Seatbelt (−0.929): Holding unemployment constant, each one-unit increase in seatbelt law stringency (from no law to secondary, or secondary to primary) is associated with a decrease of **0.929 fatalities per 100M VMT**. This confirms seatbelt laws save lives even after controlling for economic conditions.

Unemployment (+0.471): Holding seatbelt usage constant, each 1-percentage-point increase in the unemployment rate is associated with an increase of **0.471 fatalities per 100M VMT**. This counterintuitive positive association reflects that unemployment captures macro-economic stress, reduced vehicle maintenance, and potentially riskier driving behavior. Alternatively, states with historically higher unemployment often coincide with more rural, high-speed environments.

5.4 Model Diagnostics & Residual Analysis

```
par(mfrow = c(1, 2))
plot(model_multi, which = 1:2) # Residuals vs Fitted + Q-Q Plot
```




Figure 4 — Diagnostic plots for the multiple regression model. Left: Residuals vs. Fitted (checks linearity & homoscedasticity). Right: Normal Q-Q plot (checks normality of residuals). Observations 1175, 1176, and 701 are flagged as potential influential outliers.

Diagnostic	Observation	Verdict
------------	-------------	---------

Residuals vs. Fitted (Linearity)	Residuals scattered around zero with no strong curved pattern; the pink smoother is roughly flat.	✓ Linearity assumption broadly satisfied
Homoscedasticity (Equal Variance)	Some fan-like spread at higher fitted values (16–22 range) and extreme outliers visible at top of residual plot.	■ Mild heteroscedasticity present
Q-Q Plot (Normality)	Most points follow the diagonal closely; upper tail deviates substantially due to the high-outlier state-years.	■ Heavy right tail; near-normal in middle
Influential Points (obs. 1175, 1176, 701)	Labeled in both plots; represent catastrophic outlier years. Their standardized residuals exceed 4–6 standard deviations.	■ High-leverage outliers; model robust for main conclusions

FINAL CONCLUSION & POLICY RECOMMENDATIONS

This comprehensive analysis of the Wooldridge *driving* dataset — conducted by **Team Peaky Blinders Code** — reveals a clear and coherent narrative about the determinants of traffic fatality rates across U.S. states. Below we synthesize the key findings from all five analytical parts.

1

DISTRIBUTION

Fatality rates are right-skewed (Mean 18.92 > Median 18.44) with 14 severe outliers above 35.37. The median is the more representative central measure. Outliers largely represent pre-regulation state-years.

2

HYPOTHESIS ($\mu=2$)

The claim that the average fatality rate equals 2 is categorically rejected ($t \approx +92$, $p < 0.0001$). The true mean is nearly 10× higher than the claimed value.

3

SEATBELT EFFECT

States with higher seatbelt compliance average 1.36 fewer deaths per 100M VMT ($p = 0.000177$). Over national driving volumes, this represents tens of thousands of lives annually.

4

ECONOMIC ACTIVITY

Higher economic activity is paradoxically associated with higher fatality rates (20.63 vs 17.78), driven by increased road exposure rather than decreased safety per trip.

5

REGRESSION MODEL

The multiple regression model confirms both seatbelt ($\beta = -0.929$, $p < 0.0001$) and unemployment ($\beta = +0.471$, $p < 0.0001$) as significant predictors. Both coefficients are theoretically consistent and statistically robust.

EVIDENCE-BASED POLICY RECOMMENDATIONS

A. Mandate Primary Seatbelt Laws Nationally

The data is unambiguous: primary seatbelt enforcement laws (coded as 2) are associated with significantly lower fatality rates. States still operating under secondary enforcement laws should be urged to upgrade immediately.

B. Target High-Activity Corridors

High economic activity states drive more miles per capita and bear the highest fatality burden. Infrastructure investment, speed enforcement, and median barriers on high-volume interstates should be prioritized in these regions.

C. Economic Safety Nets During Recessions

The positive unemployment–fatality correlation suggests that economic stress increases driving risk. Road safety programs, vehicle inspection subsidies, and drunk-driving enforcement should be intensified during economic downturns.

D. Address Outlier States Directly

The 14 high-fatality outlier observations correspond to specific, identifiable state-years. Studying what drove catastrophic rates in those cases — and what subsequently reduced them — offers the richest policy lessons.

Summary Statement: While macroeconomic conditions set the baseline level of road exposure and risk, **behavioral compliance — specifically seatbelt usage** — remains the most controllable, directly actionable variable in reducing fatal outcomes. The statistical evidence from this analysis is clear, consistent, and practically significant across all five analytical dimensions. Every level of seatbelt law stringency measurably saves lives.

Peaky Blinders Code · Alexandria University · Faculty of Business · Statistical Packages 2026 · Dr. Assem Fareed