

Lab Group Presentation

Peter Clark

February 1, 2017






Image courtesy of Scott St George, U. MN

- Large and small scale disturbance
- Recruitment and declines
- Changes in demography, composition and importance

February 1999

COMPARING TREE-RING AND INVENTORY DATA

223






FIG. 6. Tree-ring chronologies for the whole period of available records, computed as the average of ring-area series. Standard error bars (± 1 SE) are plotted to show variability of annual values. It is evident that pines with 1990 dbh > 50 cm (LARGE PINES) were much older and experienced a greater growth decline in the last century than pines with 1990 dbh < 50 cm (SMALL PINES). Tree-ring patterns over the last 400 yr reveal that the 20th century has been unique and suggest that the extreme growth decline experienced by large pines is partly a return to normal growth rates after the growth surge of the early 1900s.

Biondi 1998. Comparing tree-ring chronologies and repeated forest inventories as forest monitoring tools. Ecological Applications



DENDROPYROLOGY Fire Scarred trees

- Scale (spatial and temporal)
- Frequency, Intensity, Seasonality
- Drivers (climate, fuels)



Figures (top) fire history from Sky Islands (AZ), (bottom) superposed epoch analysis showing relationship between drought conditions and fire








Image courtesy of Daniel Bishop, Harvard Forest / Columbia University

DENDROENTOMOLOGY Trees limited by insects

- Defoliators (spruce budworm, forest tent caterpillar, gypsy moth)
- Cambium feeder (spruce beetle, mountain pine beetle)
- Root parasite (periodical cicadas)



Swetnam and Lynch. 1993 Multicentury, Regional-Scale Patterns of Western Spruce Budworm Outbreaks. Ecological Monographs

Snow and Insect Outbreaks in Central Oregon

Collaborators: Dr. Jim Speer (*Indiana State University*),
Dr. Lawrence Winship (*Hampshire College*)

Clark, P.W. , J. H. Speer, L.W. Winship. 2017
Extracting Climate and Pandora Moth Outbreaks
from A 1,500-Year Long Ponderosa Pine Chronology
from Central Oregon. Tree-Ring Research

Related Works: Speer, J. H., T. W. Swetnam, B. E. Wickman, and A. Youngblood. 2001. Changes in Pandora moth outbreak dynamics during the past 622 years. Ecology

Speer, J. H. and R. L. Holmes. 2004. Effects of Pandora moth outbreaks on ponderosa pine wood volume.

Tree-Ring Research





Inner Ring: 435CE
Outer Ring 985CE


A photograph showing two people in a volcanic landscape. The foreground and middle ground are covered in dark, jagged lava rocks. A person in a red jacket and hat is standing on the left, holding a blue pole. Another person in a green jacket and backpack is standing in the center, holding a large, light-colored, circular object. The background is a dense forest of tall evergreen trees under a blue sky with some clouds.

PANDORA MOTH




PANDORA MOTH OUTBREAK SIGNAL IN TREE RINGS

- Year 1: Reduced latewood
- Year 2: Fifty percent reduction in ring width compared to Y-1
- Years 3 through 4-20: gradually resume to normal growth



Bark ← → Inside




PANDROA MOTH AND SNOWPACK RECONSTRUCTED

****A**** The 1,376 year reconstruction of Fall to Spring Drought

****B**** A 5-year running average of the final chronology with pandora moth outbreaks shaded in black.

****C**** The number of trees recording pandora moth outbreaks (bars) as compared to total sample depth. The gray bars show the timing of pandora moth outbreaks across all three graphs



IMPLICATIONS

Ponderosa pine extremely long lived and preserved on MBF

Contributes to natural range of outbreak variability of endemic insect

Methods for decoupling limit factors in tree growth



QUESTIONS?

