

# Tables

2.1 Transformations of Dirac spinor and adjoint spinors under various symmetry operations.	92
2.2 Transformations of Dirac bilinear covariants under various symmetry operations.	93
5.1 $ft$ values from super-allowed Fermi decays.	175
5.2 The columns list the parent nucleus in the transition, the initial, and final spins, and the transition type. F stands for a pure Fermi transition and GT stands for a Gamow–Teller transition.	176
5.3 Nucleon operators for the decay $n \rightarrow e^- \bar{\nu}_e$ in the relativistic and non-relativistic limits, where $\sigma_{\mu\nu} = \frac{i}{2}[\gamma_\mu, \gamma_\nu]$ , and $\psi$ s are the nuclear wave functions.	178
5.4 Values of the coefficients $A$ and $B$ appearing in Eq. (5.67).	187
5.5 Electron and muon families with their respective lepton numbers.	189
5.6 Michel parameters in $V - A$ theory and their values determined in the experiments.	197
5.7 Decay modes of tau lepton.	212
5.8 Michel parameters of the $\tau$ lepton.	214
6.1 Quarks are spin $\frac{1}{2}$ fermions with baryon number $\frac{1}{3}$ ; they have been assigned positive parity. $Q( e )$ is the quark's charge in the units of electronic charge, $I$ is the isospin and its 3rd component is $I_3$ . $S$ , $C$ , $B$ , and $T$ stand for the strangeness, charm, bottom, and top quantum numbers, respectively.	215
6.2 Mesons ( $J^P = 0^-$ ) and their properties: quark content, $J$ represents angular momentum, $P$ the parity, $I$ the isospin, $I_3$ the 3rd component of $I$ , $S$ the strangeness, $M$ the mass and $\tau$ the lifetime of a given meson.	216
6.3 Baryons ( $J^P = \frac{1}{2}^+$ ) and their properties: quark content, isospin( $I$ ) and its 3rd component ( $I_3$ ), total angular momentum ( $J$ ) and parity ( $P$ ), strangeness ( $S$ ), mass ( $M$ ), and lifetime( $\tau$ ).	216
6.4 Decay modes with the corresponding branching ratios of charged as well as neutral pions and kaons.	218

6.5 Decay modes of hyperons with branching ratios. Neutron decays to $pe^- \bar{\nu}_e$ with 100% branching ratio.	218
6.6 Production modes of kaons.	234
6.7 Cabibbo model predictions for the $\beta$ -decays of baryons. Here, for simplicity, we have dropped the superscript $A$ from $F$ , $D$ in $g_1(0)$ and $g_1/f_1$ . The variables used in the table are defined as: $V_{ud} = \cos \theta_C$ , $V_{us} = \sin \theta_C$ , $\mu_p = 1.7928\mu_N$ , and $\mu_n = -1.9130\mu_N$ .	244
6.8 Results from $\sin \theta_C$ analysis using measured $g_1/f_1$ values.	247
6.9 Nonleptonic decays of hyperons.	251
6.10 Predicted $c\bar{c}$ and $b\bar{b}$ states with principal quantum numbers $n=1$ and 2, and radial quantum number $n_r = n - L$ , compared with experimentally observed states. Masses are given in $MeV/c^2$ .	265
6.11 Some of the observed parity violating effects in nucleons and nuclei. $A_L$ represents asymmetry and $P_L$ represents the polarization of the hadron(photon) in the final state.	271
8.1 Weak isospin ( $I$ ), its third component ( $I_3$ ), charge ( $Q( e )$ ) and hypercharge ( $Y = 2(Q - I_3)$ ) of leptons and scalar mesons in the W-S model with $l = e, \mu, \tau$ .	320
8.2 Couplings of the leptons to $Z_\mu$ field.	335
8.3 Isospin ( $I$ ), its third component ( $I_3$ ), charge ( $Q( e )$ ), and hypercharge ( $Y = 2(Q - I_3)$ ) of the first and second generation of quarks, that is, $(u, d)$ and $(c, s)$ quarks.	336
8.4 Couplings of the quarks $(u, d)$ to $Z_\mu$ field	337
8.5 Values of $\sin^2 \theta_W$ calculated from various data sets from different experiments.	341
8.6 Results from selected parity violating (PV) experiments. Asymmetries are given in parts per million (ppm).	343
8.7 Results from selected PV experiments. Asymmetries are given in ppm.	346
8.8 Experimental determinations of the ratios $g_l/g_v$ .	355
8.9 Branching ratios of the Higgs boson decays.	360
9.1 Values of $\alpha$ , $\beta$ , and $\gamma$ for $\nu_\mu e^-$ , $\bar{\nu}_\mu e^-$ , $\nu_e e^-$ , and $\bar{\nu}_e e^-$ scattering.	373
9.2 Measured values of the total cross section and $\sin^2 \theta_W$ from different experiments for $\nu_\mu e^-$ and $\bar{\nu}_\mu e^-$ scattering at 90% confidence level.	375
9.3 Cross section of neutrino–electron $\nu_e e$ ( $\bar{\nu}_e e$ ) scattering processes from different experiments. <sup>a</sup> Region in visible energy [1.5–3.0] MeV. <sup>b</sup> Region in visible energy [3.0–4.5] MeV.	376
9.4 Magnetic moment of neutrinos as measured by the different experiments.	377
10.1 Recent measurements of the axial dipole mass ( $M_A$ ).	410
11.1 Charged and neutral current induced inelastic processes. $N, N'$ represents protons and neutrons, $Y = \Lambda, \Sigma$ represents hyperons, $K = K^+, K^0$ represents kaons, $\bar{K} = K^-, \bar{K}^0$ represents antikaons, and $l = e, \mu, \tau$ represents leptons.	426
11.2 Properties of the spin 1/2 resonances available in the Particle Data Group (PDG) Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , parity $P$ , and the branching ratio of different meson–baryon interactions like $N\pi$ , $N\eta$ , $K\Lambda$ , and $K\Sigma$ .	427

11.3 Properties of the spin 3/2 resonances available in the PDG with Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , parity $P$ , and the branching ratio of different meson–baryon interactions like $N\pi$ , $N\eta$ , $K\Lambda$ , and $K\Sigma$ .	428
11.4 Properties of the meson (pion and kaon) resonances available in PDG with Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , spin $J$ , and parity $P$ .	428
11.5 Properties of the spin 1/2 and 3/2 hyperon resonances available in PDG with Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , spin $J$ , and parity $P$ .	429
11.6 MAID parameterization of the transition form factors for the spin 1/2 resonance on a proton target. $\mathcal{A}_\alpha(0)$ is given in units of $10^{-3} \text{ GeV}^{-\frac{1}{2}}$ and the coefficients $a_1$ , $a_2$ , $a_4$ , $b_1$ in units of $\text{GeV}^{-2}$ , $\text{GeV}^{-4}$ , $\text{GeV}^{-8}$ , $\text{GeV}^{-2}$ , respectively. For all fits, $a_3 = 0$ . For $S_{11}(1535)$ , $S_{31}(1620)$ , and $S_{11}(1650)$ , resonance $a_2$ and $a_4$ are taken to be 0.	436
11.7 MAID parameterization for a neutron target ( $a_{2,3,4} = 0$ ) for spin 1/2 resonances.	437
11.8 MAID parameterization of the transition form factors for the spin 3/2 resonance on a proton target. Different components of this table have the same meaning as in Table 11.6.	449
11.9 MAID parameterization for neutron target ( $a_{2,3,4} = 0$ ) for spin 3/2 resonances.	449
11.10 Expressions for $O_i^\mu(V)$ , $i = 1 - 6$ and $O_i^\mu(A)$ , $i = 1 - 8$ , where $P = p + p'$ .	465
12.1 The values of constant term ( $\mathcal{A}^i$ ) appearing in Eq. (12.6), where $i$ corresponds to the nucleon pole (NP), cross nucleon pole (CP), contact term (CT), pion pole (PP), and pion in flight (PF) terms. $f_\pi$ is the pion weak decay constant and $g_1$ is nucleon axial vector coupling.	482
12.2 Coupling constant ( $\mathcal{C}^{\mathcal{R}}$ ) for spin $\frac{1}{2}$ and spin $\frac{3}{2}$ resonances in the case of charged current induced processes. Here, $f^*$ stands for $R_{1/2} \rightarrow N\pi$ coupling which for $\Delta(1232)$ resonance is $f_{N\Delta\pi}$ and is $f_{NR_{1/2}\pi}(f_{NR_{3/2}\pi})$ for spin $\frac{1}{2}(\frac{3}{2})$ resonances.	485
12.3 The standard form factors for weak CC transitions of the $\text{SU}(3)$ baryon octet.	496
12.4 Constant factors appearing in the hadronic current. The upper sign corresponds to the processes with antineutrinos and the lower sign to that with neutrinos.	496
12.5 Values of the parameters appearing in the hadronic currents.	500
12.6 The physical states of the decuplet.	503
12.7 Constant factors appearing in the hadronic current.	505
13.1 Kinematical quantities for hadron and parton.	518
14.1 Parity of vector and axial vector multipoles.	573
14.2 Binding energy, and $Q$ value of the reaction for various nuclei. The last three columns are the parameters for MHO and 2pF densities. * is dimensionless for the MHO density.	584
15.1 Coefficients used in Eq. (15.24) for the calculation of $\text{Im}\Sigma_\Delta$ as a function of energy in the case of pion nuclear scattering.	612
15.2 Coefficients used for an analytical interpolation of $C(T_\pi)$ of Eq. (15.25).	613

16.1	Some of the experiments which studied (anti)neutrino induced DIS cross section and structure functions from nuclear targets.	638
16.2	The developments in the global DGLAP analysis of nPDFs since 1998. DIS: deep inelastic scattering; DY: Drell–Yan di-lepton production; nFFs: nuclear fragmentation functions.	641
17.1	Solar neutrino detectors and their main characteristics. LS stands for liquid scintillator detectors.	676
17.2	Atmospheric neutrino detectors.	684
17.3	Reactor (short baseline) neutrino detectors.	687
17.4	Reactor (long baseline) neutrino detectors.	687
17.5	Supernova neutrino detectors.	689
17.6	Geoneutrino detectors.	694
17.7	List of accelerators.	700
17.8	Accelerator (short baseline) neutrino detectors.	701
17.9	Accelerator (long baseline) neutrino detectors.	701
18.1	The impact of matter potential in various medium.	736
18.2	$\Delta m^2$ reach of experiments using various neutrino sources. These experiments have different $L$ and $E$ range.	742
18.3	Neutrino oscillation experiments sensitive to different $\theta_{ij}$ and $\Delta m_{ij}^2$ .	742
18.4	The best fit values of the three neutrino oscillation parameters from a global fit of the current neutrino oscillation data. The values (in brackets) correspond to $m_1 < m_2 < m_3$ ( $m_3 < m_1 < m_2$ ). The definition of $\Delta m^2$ used is: $\Delta m^2 = m_3^2 - \frac{m_2^2 + m_1^2}{2}$ . Thus, $\Delta m^2 = \Delta m_{31}^2 - \frac{\Delta m_{21}^2}{2} > 0$ , if $m_1 < m_2 < m_3$ and $\Delta m^2 = \Delta m_{32}^2 + \frac{\Delta m_{21}^2}{2} < 0$ , for $m_3 < m_1 < m_2$ .	743
19.1	Different neutrino processes in supernova and proto-neutron star matter.	752
20.1	Some of the measured values of $T_{1/2}^{2\nu}$ in various nuclei.	776
20.2	$T_{1/2}^{0\nu}$ and $\langle m_{\beta\beta} \rangle$ limits (90% C.L.) from the most recent experiments on $0\nu\beta\beta$ in various nuclei.	778
20.3	Table for rare leptonic processes with lepton flavour violation (LFV).	788
20.4	Rare hadronic processes with lepton flavor violation (LFV).	789
20.5	Recent experimental limits of FCNC kaon decays.	790
20.6	Recent values of various couplings $C_i$ ( $i = V, A, S, T$ ) determined from $\beta$ decays.	793
B.1	Values of the structure constants $f_{ijk}$ and $d_{ijk}$ .	810
B.2	Values of the coefficients $a$ and $b$ given in Eqs. (B.23) and (B.24).	815