

Questão 01

> restart

> f:=x->(3*exp(x)-1)/(x-2)*ln((x^2+2*x+3)/(x^2-3))

$$f := x \mapsto \frac{(3 \cdot e^x - 1) \cdot \ln\left(\frac{x^2 + 2 \cdot x + 3}{x^2 - 3}\right)}{x - 2} \quad (1)$$

Item (a)

> solve((x^2+2*x+3)/(x^2-3)>0)

$$(K - \infty, K \sqrt{3}), (\sqrt{3}, \infty) \quad (2)$$

> x<>2

$$x \neq 2 \quad (3)$$

> solve(x^2-3<>0)

$$\{x \neq \sqrt{3}, x \neq K \sqrt{3}\} \quad (4)$$

$$\text{Dom}(f) = (K - \infty, K \sqrt{3}), (\sqrt{3}, \infty), x \neq 2$$

Item (b)

> limit(f(x), x=2, left)

$$K - \infty \quad (5)$$

> limit(f(x), x=2, right)

$$\infty \quad (6)$$

> limit(f(x), x=-sqrt(3), left)

$$\infty \quad (7)$$

> limit(f(x), x=sqrt(3), right)

$$K - \infty \quad (8)$$

Assíntotas verticais:

> x=2;x=sqrt(3);x=-sqrt(3)


$$x = 2$$

$$x = \sqrt{3}$$

$$x = K \sqrt{3} \quad (9)$$

Item (c)

> plot([D(f)(x), 3], x=2..5)




```
> x0:=fsolve(D(f)(x)=3,x=3..4)
```

```
x0 := 3.708059748
```

(10)

Item (d)

```
> plot([D(f)(x), -8], x=-5..5)
```



```
> x0:=fsolve(D(f)(x)=-8)
```

(11)

```
> f(x0)
```

(12)

```
> y=D(f)(x0)*(x-x0)+f(x0)
```

(13)

Item (e)

```
> p:=x->a2*(x-3)^2+a1*(x-3)+a0
```

(14)

```
> s:=fsolve([(D@@0)(f)(3)=(D@@0)(p)(3), (D@@1)(f)(3)=(D@@1)(p)(3),
(D@@2)(f)(3)=(D@@2)(p)(3)])
```

(15)

```
> evalf(subs(s,p(3.1)));f(3.1);evalf(subs(s,p(3.1))-f(3.1))
```

(16)

Falso, pois $0.04207373 > 10^{-2}$

Questão 02

Item (a)

```
> restart
```

```
> V:=r->Pi*r^2*h
```

$$V := r \mapsto \pi \cdot r^2 \cdot h \quad (17)$$

```
> s:=solve(V(r)=150,h)
```

$$s := \frac{150}{\pi r^2} \quad (18)$$

```
> h:=unapply(s,r)
```

$$h := r \mapsto \frac{150}{\pi \cdot r^2} \quad (19)$$

```
> C:=r->Pi*r^2*3+2*Pi*r*h(r)*2
```


$$C := r \mapsto 3 \cdot \pi \cdot r^2 + 4 \cdot \pi \cdot r \cdot h(r) \quad (20)$$

```
> C(r)
```

$$3 \pi r^2 + \frac{600}{r} \quad (21)$$

Item (b)

```
> plot(C(r),r=0..10)
```



```
> rmin:=fsolve(D(C)(r)=0,r=2..4)
```

