

New Progress in Classic Area: Polynomial Root-squaring and Root-finding II

Soo Go¹, Victor Pan², and Pedro Soto³[0000–0002–7120–7362]

¹ The Graduate Center, CUNY New York, NY, USA sgo@gradcenter.cuny.edu

² Lehman College, CUNY New York, NY, USA victor.pan@lehman.cuny.edu

³ The Graduate Center, CUNY New York, NY, USA psoto@gradcenter.cuny.edu
<https://pedrojuansoto.github.io/>

Abstract. Motivated by the extensive work in the companion paper [5] submitted to CASC, we devised, analyzed and tested numerically an algorithm for computing a k th order derivative of the ratio $p'x/p(x)$ for a black box univariate polynomial $p(x)$ – given by a black box for its evaluation. This is an alternative to the algorithm in [5], which also works for a black box univariate polynomial $p(x)$.

Keywords: symbolic-numeric computing · root finding · polynomial algorithms · computer algebra.

1 Introduction

Our subject is a novel efficient algorithm for the computation of ratio $p'x/p(x)$ for a black box univariate polynomial $p(x)$ – given by a black box for its evaluation. This goal was motivated by the discovery in our companion submission to CASC [5] of the reduction to this problem of the classic root-squaring DLG iterations by Dandelin 1826, Lobachevsky 1834, Gräffe 1837(cf. [10]). For the sake of completeness of our exposition we reproduce the relevant material from [5] in the Appendix.

2 Related Works

The main method for root-finding by root-squaring was independently discovered by Dandelin, Lobachevsky, and Gräffe in the 19th century over the course of 10 years (See [10]). Two of the first works to consider algorithms based on DLG formulae for modern computers were [6] and [7]; the latter of which gave explicit pseudocode for the a root-finding algorithm that uses the DLG root squaring (Eq. 7). The work [12] also considers root-finding using Eq. 7 and further states that the DLG method becomes very unstable for more than 2 iterations; by contrast, as we will see in Sec. 5, our algorithm performs reasonably well for as many iterations as 12. The same authors went on to design a variation of the DLG root-finding algorithm that makes use of different renormalizations and

other preprocessing transformations for DLG root finding in [13], and the work also proves convergence results for their DLG iterative algorithm.

The authors of [3] consider DLG based algorithms for the solutions of fractional-order polynomials, *i.e.*, generalized polynomials that have rational exponents. In [9] the authors apply the DLG formulae to root counting. The authors of [8] apply the DLG formulae to solving polynomials over finite fields. The authors of [2] apply DLG iterations to the benchmark problem (*i.e.*, the isolation of the roots of a polynomials, See Sec. A) to improve upon the record bound in [14] which at the time was unbroken for 14 years.

This short survey illustrates that the application of the DLG to root-finding, while more than a century old, is still an interesting research topic. One of the novelties and improvements of our algorithms is to consider the identities given by Eq. 8, instead of the usual Eq. 7, to approximate the root radius. Furthermore, we show that, contrary to intuition, our algorithm is numerically stable and well-behaved when taking the limit of Eq. 8 to 0.

3 Our Algorithm for Evaluating $\left(\frac{p'(x)}{p(x)}\right)^{(\ell)}$




Fig. 1. The upper tree depicts the steps of `CIRCLE_ROOTS_RATIONAL_FORM`(p, q, l) in Alg.1 for $l = 2$, $p = 1$, and $q = 1$. The lower tree depicts the steps of `ROOTS`(r, t, u, l) in Alg.2 for $r = 2$, $l = 2$, $p = 1$, and $q = 1$

In this section we present the design of the general algorithm for approximating the root radius given by Eq. 9. There are two main steps: 1) going




Fig. 2. The steps of $\text{DLG_RATIONAL_FORM}(p, p', r, t, u, l)$ in Alg.3 for $r = 2$, $l = 2$, $t = 1$, and $u = 1$.

down the rational root tree, *i.e.*, performing Alg. 1, and 2) going up the rational root tree, *i.e.*, performing Alg. 3. The going-down is depicted in Fig. 1 and the going-up is depicted in Fig. 2. The other procedures are simple bookkeeping/preprocessing steps in between the two main going-down/going-up steps. In particular we 1) first convert a complex number x into polar coordinates r, θ where $\theta \approx \frac{p}{q} = \frac{p}{2^e}$, *i.e.*, perform Alg. 4, 2) perform the first going-down pass which gives the rational angles for the roots of x in fraction form, *i.e.*, perform Alg. 1, 3) perform the second pass of the going-down algorithm where we compute the values $|x|^{\frac{1}{2^m}} \exp(2\pi i \frac{p}{q})$ at the m^{th} level, and 4) finally compute the values given by Eq. 8 going back up the rational root tree.

The intuition behind Alg. 1 is that the square root operation satisfies

$$p \% q \neq 0 \implies \sqrt{\exp\left(2\pi i \frac{p}{q}\right)} = \exp\left(2\pi i \frac{p}{2q}\right) \quad (1)$$

Algorithm 1 CIRCLE_ROOTS_RATIONAL_FORM(p, q, l)

```

if  $p \% q == 0$  then
   $r, s := (1, 1)$ 
else
   $r, s := (p, 2q)$ 
end if
if  $r \% s == 0$  then
   $t, u := (1, 2)$ 
else
   $t, u := (2r + s, 2s)$ 
end if
if  $l == 1$  then
  return  $[(r, s), (t, u)]$ 
else if  $l != 0$  then
  left := CIRCLE_ROOTS_RATIONAL_FORM( $r, s, l - 1$ )
  right := CIRCLE_ROOTS_RATIONAL_FORM( $t, u, l - 1$ )
  return left  $\cup$  right
else
  return  $[(p, q)]$ 
end if

```

and

$$p \% q = 0 \implies \sqrt{\exp\left(2\pi i \frac{1}{1}\right)} = \exp\left(2\pi i \frac{1}{1}\right) = 1, \quad (2)$$

and the negation operation satisfies

$$p \% q \neq 0 \implies -\exp\left(2\pi i \frac{r}{s}\right) = \exp\left(2\pi i \frac{2r + s}{2s}\right) \quad (3)$$

and

$$p \% q = 0 \implies \sqrt{\exp\left(2\pi i \frac{1}{1}\right)} = \exp\left(2\pi i \frac{1}{2}\right) = -1; \quad (4)$$

therefore, the first four lines of Alg. 1 compute the (angle of the) positive square root, \sqrt{x} , of complex number on the unit circle and the next four lines Alg. 1 computes the (angle of the) negative square root, $-\sqrt{x}$. Therefore, we have:

Theorem 1. *For a complex number x with a rational angle $\frac{p}{q}$, i.e., $x = |x| \exp\left(2\pi i \frac{p}{q}\right)$, Alg. 1 correctly computes the roots in Eq. 8. In particular, for a rational angle in fractional form, it does so with exact precision.*

Proof. Equations 1, 2, 3, and 4 give the base case and the theorem follows by a straightforward induction.

Alg. 2 and Alg. 4 are both straightforward, so for the rest of this section we focus on the intuition behind Alg. 3.

Algorithm 2 ROOTS(r, t, u, l)

```

root_tree = CIRCLE_ROOTS_RATIONAL_FORM( $p, q, l$ )
circ_root = [exp( $2 \cdot \pi \cdot i \cdot \frac{r}{s}$ ) for  $r, s$  in root_tree]
roots = [  $\sqrt[l]{r \cdot \text{root}}$  for root in circ_root]
return roots

```

Algorithm 3 DLG_RATIONAL_FORM(p, p', r, t, u, l)

```

root := ROOTS( $r, t, u, l$ )
for  $r_i \in \text{root}$  do
    base_step[ $i$ ] :=  $\frac{p'(r_i)}{p(r_i)}$ 
end for
diff[0] := base_step
for  $i \leq l$  do
    for  $j \leq 2^{l-i-1}$  do
        diff[ $i+1$ ][ $j$ ] :=  $\frac{1}{2} \frac{\text{diff}[i][2j] - \text{diff}[i][2j+1]}{\text{root}[2j]}$ 
        root = roots( $r, t, u, l-1-i$ )
    end for
end for
return diff[ $l$ ][0]

```

Alg. 3 is a dynamic programming on Equation 8. Therefore, by Thm. 1, we have that Alg. 3 correctly computes Eq. 8. Specifically, Alg. 3 does the following: 1) it computes the last layer of the recursion Eq. 8 (*i.e.*, it computes $p'(x^{\frac{1}{2^l}})/p(x^{\frac{1}{2^l}})$) and then 2) it recursively applies Eq. 8 via dynamic programming until it finally computes $\frac{p'_\ell(x)}{p_\ell(x)}$ which is the desired quantity. The former is given by the line “base_step[i] := $\frac{p'(r_i)}{p(r_i)}$ ” and the latter is given by the line “diff[$i+1$][j] := $\frac{1}{2} \frac{\text{diff}[i][2j] - \text{diff}[i][2j+1]}{\text{root}[2j]}$ ” in Alg. 3; the desired output (*i.e.*, $p'(x^{\frac{1}{2^l}})/p(x^{\frac{1}{2^l}})$) is given by the line “diff[l][0]”.

Algorithm 4 DLG(p, p', l, x, ϵ)

```

angle :=  $\frac{1}{2\pi i} \log(x)$ 
 $u := 2^\epsilon$ 
 $t := (\text{angle} \cdot u) \% 1$ 
 $r := |x|$ 
return DLG_RATIONAL_FORM( $p, p', r, t, u, l$ )

```

The final step in the algorithm is to use Alg. 4 to compute root radius approximations r_d and r_1 . The procedure is given by Alg. 5. The rationale for generating a random x is that there may be roots close to 0 and thus, by taking the limit in certain directions, we avoid these possible poles; in particular, we

Algorithm 5 DLG_ROOT_RADIUS($p, p', p_{\text{rev}}, p'_{\text{rev}}, l, \epsilon, \epsilon'$)

 Uniformly Randomly Generate x in the unit circle
 $d := \deg(p)$ $r_{\min} := d/\text{DLG}(p, p', l, x \cdot 2^{-\epsilon}, \epsilon')$ $r_{\max} := \text{DLG}(p_{\text{rev}}, p'_{\text{rev}}, l, x \cdot 2^{-\epsilon}, \epsilon')/d$

have that

$$\lim_{\epsilon', \epsilon \rightarrow \infty} \text{DLG}(p, p', l, x \cdot 2^{-\epsilon}, \epsilon) = \frac{p'_\ell(0)}{p_\ell(0)} = \left(\frac{p'_{\ell-1}(x)}{p_{\ell-1}(x)} \right)'_{x=0} = \left(\frac{p'(x)}{p(x)} \right)^{(\ell)}_{x=0}, \quad (5)$$

for any x , if $p(0) \neq 0$ (See Thm 2).

4 Theoretical Analysis

We now give some theoretical guarantees: Thm. 2 proves the correctness of Alg. 5, and Thm. 3 and Thm. 4 give computational complexity bounds.

4.1 Correctness of the output

Theorem 2. *If $p(0) \neq 0$, then Alg. 5 computes the bounds given by Eq. 10 with probability 1.*

Proof. By Lem. 1 we have that the limit in Eq. 1 is well defined. Since there are at most finitely many roots for p_ℓ with a high probability Alg. 5 computes the correct approximation to the bounds in Eq. 10.

4.2 Complexity

Theorem 3. *Alg. 3 performs q floating point subtractions, divisions, and multiplications and q applications of \sin and \cos , where $q = 2^l$; furthermore, Alg. 3 performs at most Cq integer additions, “multiplications-by-2”, and $\%2^\epsilon$ (i.e., $\text{mod } 2^\epsilon$) operations, where $C = 1, 3, 2$ respectively.*

Proof. Looking at Fig. 1 and Fig. 2, we can see that the computational tree for the Alg. 3 is a binary tree with $2^l = q$ nodes; the proof for the constants $C = 1, 3, 2$ follows similarly follows from the inspection of the operations performed in Alg. 1.

Theorem 4. *The Cq integer additions, “multiplications-by-2”, and $\%2^\epsilon$ (i.e., $\text{mod } 2^\epsilon$) operations in Alg. 3 have negligible overhead. More precisely, integer additions are always additions of $2 \in \log \ell$ -bit integers and “multiplications-by-2” and $\%2^\epsilon$ (i.e., $\text{mod } 2^\epsilon$) operations have constant time overhead.*

Proof. Since Alg. 4 always passes in a denominator which is a power of two all of the integer $\%$ and \cdot operations are in fact “multiplications-by-2”, and $\%2^\epsilon$ (i.e., mod 2^ϵ) operations by a straightforward proof similar to the one in Alg. 3. Thus these operation are essentially constant overhead bit shift operations on a computing machine with binary words. Since $p\%r$ always reduces $p \mapsto 1$, we have that whenever an overflow of more than $\log r$ -bits happens in Alg. 1 it gets converted to an $\log r$ -bit integer; therefore, it suffices to prove that $\log r$ is bounded by $\epsilon \log \ell$. However, this once again follows by induction on the binary computation tree: since this tree has depth ℓ we see that any denominator is bounded by $\epsilon \log \ell$ by a simple induction.

In order to prove Thm. 2 we must first prove Lem. 1.

Lemma 1. *If $p(0) \neq 0$, then the limit $\lim_{x \rightarrow 0} \text{DLG}(p, p', l, x)$ is always well-defined.*

Proof. By applying induction on Eq. 7, we have that $p_l(0) \neq 0$ if $p(0) \neq 0$, and thus it suffices to consider the behavior of the numerator in Eq. 8. The case $\ell = 1$ follows from an application of L'Hopital's rule. For $\ell \neq 1$, there are two cases: either \sqrt{x} divides the numerator of $\frac{p'_\ell(\sqrt{x})}{p_\ell(\sqrt{x})}$ or it does not. If it does, then we are done since we can once again apply L'Hopital's rule. Otherwise, we get that $p'_\ell(\sqrt{x}) = c_0 + c_1\sqrt{x} + \dots + c_k(\sqrt{x})^k$ for some c_i with $c_0 \neq 0$. But then we have

$$\begin{aligned} \frac{p'_{\ell+1}(\sqrt{x})}{p_{\ell+1}(\sqrt{x})} &= \frac{1}{2\sqrt{x}} \left(\frac{p'_\ell(\sqrt{x})}{p_\ell(\sqrt{x})} - \frac{p'_\ell(-\sqrt{x})}{p_\ell(-\sqrt{x})} \right) \\ &= \frac{1}{2\sqrt{x}} \frac{b_0 c_0 + \sqrt{x} \cdot N_1(\sqrt{x}) - b_0 c_0 - \sqrt{x} \cdot N_2(\sqrt{x})}{p'_\ell(\sqrt{x}) p'_\ell(-\sqrt{x})} \\ &= \frac{1}{2} \frac{N_1(\sqrt{x}) - N_2(\sqrt{x})}{p'_\ell(\sqrt{x}) p'_\ell(-\sqrt{x})}, \end{aligned}$$

for some polynomials N_1 and N_2 with $b_0 = p'_\ell(0)$. Therefore the limit at zero is once again well-defined.

4.3 Stability

Definition 1. *If \tilde{f} is an algorithm for computing f we let $\delta f(x) = f(x) - \tilde{f}(x)$ then the condition number of $\{f\}(x)$ at x , $\kappa\{f\}(x)$, is given by*

$$\kappa\{f\}(x) = \lim_{\epsilon \rightarrow 0} \sup_{\|\delta x\| \leq \epsilon} \frac{\|\delta f(x)\|}{\|\delta x\|}.$$

Lemma 2. *The (relative) condition number operator κ satisfies the following properties:*

1. $\kappa\{f\}(x) = |x \log'(f(x))|$
2. $\kappa\left\{\frac{f}{g}\right\}(x) = |\kappa\{f\}(x)| - |\kappa\{g\}(x)|$

$$3. \kappa\{x^d\}(x) = d$$

Proof. This follows from the definition of the condition number.

Theorem 5. *If $p_\ell(0) \neq 0$, then $\frac{p'_\ell(x)}{p_\ell(x)}$ is well-conditioned at any point sufficiently close to 0.*

Proof. The proof of Lem. 1 gives us that $\frac{p'_\ell(x)}{p_\ell(x)}$ is a well behaved rational function at any point close to zero and thus the condition number $\kappa\{\frac{p'_\ell}{p_\ell}\}(x)$ is well defined at this point since the condition number of arithmetic operations of functions are themselves arithmetic operations in those same functions and the conditions number for a polynomial of degree $d \in \mathbb{R}$ is exactly d ; therefore, $\kappa\{\frac{p'_\ell}{p_\ell}\}(x)$ has a well defined/bounded condition number in the limit to zero.

Remark 1. Even though Thm. 5 states that $\frac{p'_\ell(x)}{p_\ell(x)}$ is highly stable, in practice, computing this function requires the use of trigonometric functions (*i.e.*, the roots in Alg. 2); therefore, our added precision in Alg. 1 helps with the instability associated with Alg. 2. Intuitively this helps the instability because the trigonometric functions are the only subroutines in the algorithm that have a non constant condition number and thus, special care must be taken with them; in particular, our algorithm gives a precise value (up to user specified precision) of the angle arguments to these trigonometric functions.

5 Experimental Results

5.1 Setup

We now present the results of our experiments in which we compute the bounds on the extremal root radii $|x_1|$ and $|x_d|$ given by 10 for the polynomials in the test suite of MPSolve. The test suite covers a number of univariate polynomial families over a range of degrees. (A fuller description of the test suite can be found in <https://numpi.dm.unipi.it/mpsolve-2.2/mpsolve.pdf>.)

Given a polynomial $p(x)$ of degree d , we use an implementation of DLG iterations that incorporates our algorithm from Section 3 for computing $(p'(x)/p(x))^{(\ell)}$ and in turn compute bounds, or estimates, for root radii, $r_1 = r_{\max}$ and $r_d = r_{\min}$ given by (10) on extremal root radii $|x_1|$ and $|x_d|$, respectively. The performance of these bounds are evaluated on the *relative error* in comparison to the corresponding root radius computed by MPSolve. That is,

$$\text{relative error}(r_i) = \frac{|r_i - |x_i||}{|x_i|}, \quad i = 1, d,$$

where $|x_i|$ is the minimal or maximal root radius found using MPSolve and r_i is the corresponding estimate for the root radii we compute.

For the parameters, we use $\ell = \lfloor \log_2 d \rfloor$ and for the precision we empirically found that $e = 2^{\ell/3} + 330$ bits of precision behaved pretty well; however, deducing the theoretically correct values of e would be an interesting future research direction.

By choosing $\mathcal{O}(\log d)$ iterations, we keep the number of iterations relatively small even when the degree of the polynomial increases. For instance, $\ell = 6$ for $d = \deg(p(x)) = 100$, and when d grows to 6400, we still have $\ell = 12$; likewise, since the number of bits of precision used where $e = 2^{\mathcal{O}(\ell)} = \mathcal{O}(d)$, we have that the precision did not grow too large neither.

All experiments were performed using Python 3.7.7 and MPSolve 3.2.1 on MacOS 11.6.1 with 2.8 GHz Dual-Core Intel Core i5 with 8 GB memory.

5.2 Observations

The overall results support our claims that our root radii approximations perform well when $p(x)$ has no roots extremely close to zero whereas the estimates are poor otherwise. The test results for the `chebyshev` family of polynomials in Table 1 demonstrate this trend quite clearly.

In general, the relative errors for the minimal root radius $|x_d|$ is 1.0 or less for roots away from the origin by more than 0.01, with some notable exceptions such as $p(x) = x^d - h^d$. That is, the difference between the minimal root radius bound we compute and the absolute value of the smallest root x_d tends to be less than $|x_d|$, i.e., $|r_d - |x_d|| \leq |x_d|$. Combined with the bound given in (12) gives us a rough heuristic expectation that $|x_d| \leq r_d \leq 2|x_d|$ if $|x_d| > 0.01$.

The relative errors for the maximal root radius $|x_1|$ for the lower bound r_1 reflects a similar trend: The errors are close to 1 when $|x_1|$ is large, showing that in relation to the root radius, our estimate is close to 0, the worst lower bound possible. Since $1/x_1$ the smallest root of p_{rev} , this is essentially the same situation as the root x_d of $p(x)$ being near 0.

Finally, our results demonstrate the consequence of Thm. 13 in Table 26 showing the figures for polynomial family `nroots` of the form $p(x) = x^d - 1$. The ratio of $|p(0)/p'(0)|$ is infinite, so our algorithm estimates the minimal root radius to be much larger than the actual root radius. On the other hand, the algorithm estimates the maximal root radius to be close to 0 since $|p_{\text{rev}}(0)/p'_{\text{rev}}(0)| = 0$.

5.3 Tables

The columns of the tables, in order, are respectively: d : degree of the input polynomial, ℓ : number of iterations, e : $-\log(|x|)$, `mp.dps`: the `mpmath` precision level used, the relative errors for the minimum and maximum root radii, total runtime, the extremal root radii as computed by MPSolve for the particular given polynomial.

The entries ‘-’ in the tables indicate the test was terminated before completion.

Table 1. Experimental Data for `chebyshev`

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR		r_1	TIME
20	4	616	332	0.155		0.0939	0.27	[0.0785, 0.997]	
40	5	616	332	0.0981		0.0589	1.13	[0.0393, 0.999]	
80	6	617	334	0.0593		0.0353	5.26	[0.0196, 1.0]	
160	7	617	334	2.71		0.0205	20.88	[0.00982, 1.0]	
320	8	617	334	37.6		0.465	88.4	[0.00491, 1.0]	

Table 2. Experimental Data for `chrma`

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN- TIME	MPSOLVE	
			DPS	ERROR	r_d	ERROR		r_1	ROOT
21	4	616	332	0.215		0.139	0.22	[1.0, 3.17]	
85	6	617	334	0.0369		0.0452	5.63	[1.0, 3.25]	
341	8	617	334	0.717		0.547	105.48	[0.884, 3.41]	

Table 3. Experimental Data for `chrma_d`

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR		r_1	TIME
20	4	616	332	0.16		0.175	0.26	[1.3, 3.01]	
84	6	617	334	0.0548		0.00507	5.53	[1.1, 3.06]	
340	8	617	334	0.658		0.699	93.82	[0.741, 3.11]	

Table 4. Experimental Data for `chrmc`

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d		ERROR
22	4	616	332	0.211	0.138	0.3	[1.0, 3.03]
342	8	617	334	0.718	0.279	94.64	[0.897, 4.13]

Table 5. Experimental Data for `chrmc_d`

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d	ERROR	r_1	TIME
11	3	616	332	0.251		0.242	0.05	[1.27, 2.8]
43	5	616	332	0.101		0.0996	1.23	[1.02, 2.97]
171	7	617	334	0.268		1.13	22.05	[0.715, 3.07]
683	9	619	338	0.164		0.257	374.69	[0.519, 3.1]

Table 6. Experimental Data for `curz`

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d	ERROR	r_1	TIME
20	4	616	332	0.156		0.854	0.21	[0.452, 1.15]
40	5	616	332	0.199		0.776	1.13	[0.379, 1.26]
80	6	617	334	0.086		0.227	5.49	[0.318, 1.34]
160	7	617	334	0.0385		0.509	20.79	[0.271, 1.38]

Table 7. Experimental Data for **easy**

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d ERROR	r_1 TIME	
100	6	617	334	0.12	0.0809	6.35	[0.949, 0.98]
200	7	617	334	0.068	0.047	25.57	[0.971, 0.99]
400	8	617	334	0.0381	0.0265	104.47	[0.983, 0.995]
1600	10	619	338	0.01	0.00	2228.22	[0.995, 0.999]
3200	11	619	338	0.00	-	-	[0.997, 0.999]

Table 8. Experimental Data for **exp**

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d ERROR	r_1 TIME	
50	5	616	332	3.76	0.126	1.48	[14.9, 39.4]
100	6	617	334	0.367	0.0598	7.37	[28.9, 83.9]
200	7	617	334	0.714	0.839	28.22	[56.8, 176.0]
400	8	617	334	0.965	0.985	107.96	[113.0, 365.0]

Table 9. Experimental Data for **geom1**

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d ERROR	r_1 TIME	
10	3	616	332	0.334	0.25	0.05	[1.0, 1.0E+18]
15	3	616	332	0.403	1.0	0.08	[1.0, 1.0E+28]
20	4	616	332	0.206	1.0	0.28	[1.0, 1.0E+38]
40	5	616	332	0.122	1.0	1.18	[1.0, 1.0E+78]

Table 10. Experimental Data for **geom2**

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d ERROR	r_1 TIME	
10	3	616	332	0.334	0.25	0.06	[1.0E-18, 1.0]
15	3	616	332	8.09E+4	0.287	0.06	[1.0E-28, 1.0]
20	4	616	332	2.63E+26	0.171	0.23	[1.0E-38, 1.0]
40	5	616	332	1.61E+72	0.109	1.14	[1.0E-78, 1.0]

Table 11. Experimental Data for **geom3**

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d ERROR	r_1 TIME	
10	3	616	332	0.334	0.25	0.05	[9.54E-7, 0.25]
20	4	616	332	2.41	0.171	0.24	[9.09E-13, 0.25]
40	5	616	332	1.95E+18	0.109	1.15	[8.27E-25, 0.25]
80	6	617	334	1.83E+45	0.0662	5.73	[6.84E-49, 0.25]

Table 12. Experimental Data for **geom4**

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT RADIUS
10	3	616	332	0.334	0.25	0.05	[4.0, 1.05E+6]		
20	4	616	332	0.206	0.707	0.27	[4.0, 1.1E+12]		
40	5	616	332	0.122	1.0	1.15	[4.0, 1.21E+24]		
80	6	617	334	0.0709	1.0	5.28	[4.0, 1.46E+48]		

Table 13. Experimental Data for **hermite**

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT RADIUS
20	4	616	332	0.155	0.129	0.28	[0.245, 5.39]		
40	5	616	332	0.0981	0.0876	1.31	[0.175, 8.1]		
80	6	617	334	0.0593	0.0555	5.97	[0.124, 11.9]		
160	7	617	334	0.0348	0.0335	23.56	[0.0877, 17.2]		
320	8	617	334	2.08	0.785	88.88	[0.062, 24.7]		

Table 14. Experimental Data for **kam1**

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT RADIUS
7	2	615	331	0.368	1.0	0.01	[3.0E-12, 15.8]		
7	2	615	331	0.368	1.0	0.01	[3.0E-40, 1.0E+4]		
7	2	615	331	2.37E+93	1.0	0.01	[3.0E-140, 1.0E+14]		

Table 15. Experimental Data for **kam2**

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT RADIUS
9	3	616	332	0.107	1.0	0.03	[1.73E-6, 251.0]		
9	3	616	332	0.107	1.0	0.03	[1.73E-20, 1.0E+8]		
9	3	616	332	4.23E+46	1.0	0.03	[1.73E-70, 1.0E+28]		

Table 16. Experimental Data for **kam3**

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT RADIUS
9	3	616	332	0.107	1.0	0.03	[1.73E-6, 251.0]		
9	3	616	332	0.107	1.0	0.03	[1.73E-20, 1.0E+8]		
9	3	616	332	4.23E+46	1.0	0.03	[1.73E-70, 1.0E+28]		

Table 17. Experimental Data for `kir1`

d	ℓ	e	MP.- DPS	RELATIVE ERROR	RELATIVE r_d ERROR	RUN- r_1 TIME	MPSOLVE ROOT RADIUS
8	3	616	332	0.000244	0.000244	0.02	[0.5, 0.5]
44	5	616	332	4.41E-5	0.000443	1.28	[0.5, 0.5]
84	6	617	334	2.29E-5	0.000464	2.9	[0.5, 0.5]
164	7	617	334	1.16E-5	0.000476	9.79	[0.5, 0.5]

Table 18. Experimental Data for `kir1_mod`

d	ℓ	e	MP.- DPS	RELATIVE ERROR	RELATIVE r_d ERROR	RUN- r_1 TIME	MPSOLVE ROOT RADIUS
44	5	616	332	0.000983	0.00095	1.26	[0.5, 0.5]
84	6	617	334	0.00364	0.00364	2.99	[0.498, 0.502]
164	7	617	334	0.00734	0.00749	10.35	[0.496, 0.504]

Table 19. Experimental Data for `lagurerre`

d	ℓ	e	MP.- DPS	RELATIVE ERROR	RELATIVE r_d ERROR	RUN- r_1 TIME	MPSOLVE ROOT RADIUS
20	4	616	332	0.206	0.167	0.22	[0.0705, 66.5]
40	5	616	332	0.122	0.108	1.28	[0.0357, 142.0]
80	6	617	334	0.0709	0.0659	5.63	[0.018, 297.0]
160	7	617	334	3.09	0.954	22.21	[0.00901, 610.0]
320	8	617	334	41.4	0.996	89.64	[0.00451, 1.24E+3]

Table 20. Experimental Data for `lar1`

d	ℓ	e	MP.- DPS	RELATIVE ERROR	RELATIVE r_d ERROR	RUN- r_1 TIME	MPSOLVE ROOT RADIUS
20	4	616	332	7.06E+9	1.0	0.1	[3.73E-22, 1.0E+50]
200	7	617	334	9.69E+19	0.311	0.83	[3.73E-22, 41.0]

Table 21. Experimental Data for `legendre`

d	ℓ	e	MP.- DPS	RELATIVE ERROR	RELATIVE r_d ERROR	RUN- r_1 TIME	MPSOLVE ROOT RADIUS
20	4	616	332	0.155	0.0969	0.25	[0.0765, 0.993]
40	5	616	332	0.0981	0.0604	1.15	[0.0388, 0.998]
80	6	617	334	0.0593	0.0359	6.08	[0.0195, 1.0]
160	7	617	334	2.72	0.0637	19.97	[0.00979, 1.0]
320	8	617	334	37.7	0.456	81.12	[0.0049, 1.0]

Table 22. Experimental Data for `lsr`

d	ℓ	e	MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d	ERROR	r_1
24	4	616	332	2.88E+8	1.0	0.29	[1.0E-20, 1.0E+20]
52	5	616	332	1.81E+14	1.0	0.25	[1.0E-20, 1.0E+10]
52	5	616	332	1.81E+34	1.0	0.22	[1.0E-40, 1.0E+20]
52	5	616	332	1.81E+74	1.0	0.2	[1.0E-80, 1.0E+40]
224	7	617	334	3.62E+18	1.0	9.8	[1.0E-20, 1.0E+20]
500	8	617	334	1.92E+3	1.0	6.79	[0.0001, 2.0E+4]
500	8	617	334	5.77E+3	0.995	3.27	[3.33E-5, 1.0E+3]
500	8	617	334	1.05	1.0	3.05	[0.0916, 1.0E+200]

Table 23. Experimental Data for `mand`

			MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE	
d	ℓ	e	DPS	ERROR	r_d ERROR	r_1	TIME	ROOT RADIUS
31	4	616	332	0.295	0.144		0.4	[0.445, 2.0]
63	5	616	332	0.127	0.0854		2.16	[0.403, 2.0]
127	6	617	334	0.0898	0.0488		8.58	[0.373, 2.0]
255	7	617	334	0.0609	0.701		37.41	[0.351, 2.0]
511	8	617	334	0.0234	1.63		174.71	[0.334, 2.0]
1023	9	619	338	0.357	0.142		569.08	[0.321, 2.0]
2047	10	619	338	1.12	0.241		2372.57	[0.311, 2.0]
4095	11	619	338	1.68	0.38		9209.19	[0.303, 2.0]

Table 24. Experimental Data for `mig1`

MP.- RELATIVE RELATIVE RUN-							MPSOLVE
d	ℓ	e	DPS	ERROR	r_d ERROR	r_1 TIME	ROOT RADIUS
20	4	616	332	0.126	1.0	0.08	[0.01, 2.26]
50	5	616	332	0.0157	0.987	1.8	[0.00999, 1.83E+3]
100	6	617	334	0.0563	1.0	0.5	[0.01, 1.15]
100	6	617	334	0.0183	1.0	4.05	[0.01, 7.92]
200	7	617	334	2.66	1.0	0.8	[0.01, 1.07]
200	7	617	334	2.59	0.999	9.37	[0.01, 2.33]
500	8	617	334	18.2	0.99	1.59	[0.01, 1.03]
500	8	617	334	18.0	0.984	15.65	[0.01, 1.36]

Table 25. Experimental Data for **mult**

d	ℓ	e	MP.-		RELATIVE	RELATIVE	RUN-	MPSOLVE
			DPS	ERROR	r_d	ERROR	r_1	TIME
15	3	616	332	0.29		0.189	0.12	[0.869, 1.07]
20	4	616	332	0.0782		0.475	0.24	[0.01, 2.68]
22	4	616	332	0.213		0.105	0.24	[1.0, 20.0]
68	6	617	334	0.0566		0.0556	4.59	[0.25, 2.24]

Table 26. Experimental Data for **nroots**

			MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE			
d	ℓ	e	DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT	RADIUS
50	5	616	332	5.18E+13		1.0		0.12		[1.0, 1.0]
100	6	617	334	7.06E+6		1.0		0.19		[1.0, 1.0]
200	7	617	334	2.69E+3		1.0		0.39		[1.0, 1.0]
400	8	617	334	50.5		0.981		0.79		[1.0, 1.0]
800	9	619	338	6.34		0.864		1.73		[1.0, 1.0]
1600	10	619	338	1.71		0.631		3.19		[1.0, 1.0]
3200	11	619	338	0.645		0.392		6.79		[1.0, 1.0]
6400	12	623	346	0.289		0.224		14.55		[1.0, 1.0]

Table 27. Experimental Data for **nrooti**

			MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE			
d	ℓ	e	DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT	RADIUS
50	5	616	332	4.94E+13	1.0			0.1	[1.0, 1.0]	
100	6	617	334	7.07E+6	1.0			0.32	[1.0, 1.0]	
200	7	617	334	2.69E+3	1.0			0.46	[1.0, 1.0]	
400	8	617	334	50.5	0.981			0.85	[1.0, 1.0]	
800	9	619	338	6.34	0.864			1.85	[1.0, 1.0]	
1600	10	619	338	1.71	0.631			3.23	[1.0, 1.0]	
3200	11	619	338	0.645	0.392			7.02	[1.0, 1.0]	
6400	12	623	346	0.289	0.224			13.31	[1.0, 1.0]	

Table 28. Experimental Data for **sendra**

			MP.-	RELATIVE	RELATIVE	RUN-	MPSOLVE
d	ℓ	e	DPS	ERROR	r_d ERROR	r_1 TIME	ROOT RADIUS
20	4	616	332	0.283	0.158	0.25	[0.9, 2.05]
40	5	616	332	0.159	0.101	1.25	[0.95, 2.02]
80	6	617	334	0.287	0.573	5.2	[0.975, 2.01]
160	7	617	334	0.658	2.14	24.34	[0.987, 2.01]
320	8	617	334	0.809	1.64	88.08	[0.994, 2.0]

Table 29. Experimental Data for **sparse**

d	ℓ	e	MP.-		RELATIVE		RELATIVE		RUN-		MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT	RADIUS		
100	6	617	334	0.11		1.0		0.26		[0.968, 1.01]		
200	7	617	334	0.0361		0.995		1.16		[0.969, 1.0]		
400	8	617	334	0.0211		0.929		2.19		[0.969, 1.0]		
800	9	619	338	0.0118		0.739		6.56		[0.969, 1.0]		
6400	12	623	346	0.00202		0.158		72.81		[0.969, 1.0]		

Table 30. Experimental Data for **spiral**

d	ℓ	e	MP.-		RELATIVE		RELATIVE		RUN-		MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT	RADIUS		
10	3	616	332	5.1E-7		1.21E-6		0.05		[1.0, 1.0]		
15	3	616	332	3.49E-7		1.15E-6		0.07		[1.0, 1.0]		
20	4	616	332	4.55E-7		1.3E-6		0.23		[1.0, 1.0]		
25	4	616	332	3.67E-7		1.25E-6		0.29		[1.0, 1.0]		
30	4	616	332	3.08E-7		1.21E-6		0.41		[1.0, 1.0]		

Table 31. Experimental Data for **toep**

d	ℓ	e	MP.-		RELATIVE		RELATIVE		RUN-		MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT	RADIUS		
128	7	617	334	0.0386		0.562		18.71		[1.31, 64.4]		
256	8	617	334	0.0219		0.918		73.35		[1.34, 64.4]		
128	7	617	334	0.0386		0.0272		17.68		[0.4, 13.2]		
256	8	617	334	0.0225		0.599		72.66		[0.383, 13.2]		

Table 32. Experimental Data for **wilk**

d	ℓ	e	MP.-		RELATIVE		RELATIVE		RUN-		MPSOLVE	
			DPS	ERROR	r_d	ERROR	r_1	TIME	ROOT	RADIUS		
20	4	616	332	0.206		0.141		0.22		[1.0, 20.0]		
30	4	616	332	0.237		0.0859		0.33		[1.0, 319.0]		
40	5	616	332	0.122		0.0927		1.27		[1.0, 40.0]		
80	6	617	334	0.0709		0.121		5.59		[1.0, 80.0]		
160	7	617	334	0.0404		0.824		21.82		[1.0, 160.0]		
320	8	617	334	0.0228		0.983		89.77		[1.0, 320.0]		

References

1. Baur, W., Strassen, V.: The complexity of partial derivatives. Theoretical computer science **22**(3), 317–330 (1983)
2. Becker, R., Sagraloff, M., Sharma, V., Yap, C.K.: A near-optimal subdivision algorithm for complex root isolation based on the pellet test and newton iteration. J. Symb. Comput. **86**, 51–96 (2018)
3. Bialas, S., Górecki, H.: Generalization of vieta’s formulae to the fractional polynomials, and generalizations the method of graeffe-lobachevsky. Bulletin of The Polish Academy of Sciences-technical Sciences **58**, 624–629 (2010)
4. de Boor, C.: Divided differences (1995)
5. Go, S, P.V.S.P.: New progress in classic area: Polynomial root-squaring and root-finding i. Submitted to CASC 2022
6. Grau, A.A.: On the reduction of number range in the use of the graeffe process. J. ACM **10**(4), 538–544 (oct 1963). <https://doi.org/10.1145/321186.321198>, <https://doi-org.ezproxy.gc.cuny.edu/10.1145/321186.321198>
7. Grau, A.A.: Algorithm 256: Modified graeffe method [c2]. Commun. ACM **8**(6), 379–380 (jun 1965). <https://doi.org/10.1145/364955.364974>, <https://doi-org.ezproxy.gc.cuny.edu/10.1145/364955.364974>
8. Grenet, B., van der Hoeven, J., Lecerf, G.: Deterministic root finding over finite fields using graeffe transforms. Applicable Algebra in Engineering, Communication and Computing **27**, 237–257 (2015)
9. van der Hoeven, J.: Efficient root counting for analytic functions on a disk (2011)
10. Householder, A.S.: Dandelin, lobacevskii, or graeffe. The American Mathematical Monthly **66**(6), 464–466 (1959), <http://www.jstor.org/stable/2310626>
11. Linnainmaa, S.: Taylor expansion of the accumulated rounding error. BIT Numerical Mathematics 1976 16:2 **16**, 146–160 (1976). <https://doi.org/10.1007/BF01931367>, <https://link.springer.com/article/10.1007/BF01931367>
12. Malajovich, G., Zubelli, J.P.: On the geometry of graeffe iteration. J. Complex. **17**, 541–573 (2001)
13. Malajovich, G., Zubelli, J.P.: Tangent graeffe iteration. Numerische Mathematik **89**, 749–782 (2001)
14. Pan, V.Y.: Univariate polynomials: nearly optimal algorithms for numerical factorization and root-finding. Journal of Symbolic Computation **33**(5), 701–733 (2002)

A Background and Motivation

We set up some basic notation and background: we discuss Dandelin, Lobachevsky, and Gräffe’s Formulae in Sec. A.1 and recall properties of the extremal root radii in Sec. A.2. The reverse of a polynomial $p := p_0 + p_1x^1 + \dots + p_dx^d$ is defined as $p_{\text{rev}} := p_d + p_{d-1}x^1 + \dots + p_0x^d$ and for a polynomial $p(x) = \prod_{i \in [d]}(x - x_i)$ where

$$|x_1| \geq |x_2| \geq \dots \geq |x_d|. \quad (6)$$

we define $r_i(c, p) = |x_i - c|$ as the root radii p about c . We are particular interested in the case where $c = 0$ since we can always reduce to this case (see Sec. A.2).

A.1 Dandelin, Lobachevsky, and Gräffe's Formulae

The common procedure for root radii approximation based on the classical technique of recursive root-squaring is to first make an input polynomial $p(x)$ monic by scaling it and/or the variable x and then apply k DLG (that is, Dandelin's aka Lobachevsky's or Gräffe's) root-squaring iterations (cf. [10]),

$$p_0(x) = \frac{1}{p_d} p(x), \quad p_{i+1}(x) = (-1)^d p_i(\sqrt{x}) p_i(-\sqrt{x}), \quad i = 0, 1, \dots, \ell \quad (7)$$

for a fixed positive integer ℓ (see Remark 2 below). The i th iteration squares the roots of $p_i(x)$ and consequently the root radii from the origin, as well as the isolation of the unit disc $D(0, 1)$. Then one approximates the ratio ρ_+/ρ_- , the new scaled ratio of the root radii, for the polynomial $p_\ell(x)$ within a factor of γ and readily recovers the approximation of this ratio for $p_0(x)$ and $p(x)$ within a factor of $\gamma^{1/2^\ell}$.

Given the coefficients of $p_i(x)$ we can reduce the i th root-squaring iteration, that is, the computation of the coefficients of $p_{i+1}(x)$, to polynomial multiplication and perform it in $\mathcal{O}(d \log(d))$ arithmetic operations. Unless the positive integer ℓ is small, the absolute values of the coefficients of $p_\ell(x)$ vary dramatically, and realistically one should either stop because of severe problems of numerical stability or apply the stable algorithm by Gregorio Malajovich and Jorge P. Zubelli [12], which performs a single root-squaring at arithmetic cost of order d^2 .

For a black box polynomial $p(x)$, however, we apply DLG iterations without computing the coefficients, and the algorithm turns out to be quite efficient: for ℓ iterations evaluate $p(x)$ at 2^ℓ equally spaced points on a circle and obtain the values of the polynomial $p_\ell(x) = \prod (x - x_j^{2^\ell})$ at these 2^ℓ points.

Furthermore, we evaluate the ratio $p'(x)/p(x) = p'_0(x)/p_0(x)$ at these points by applying the recurrence

$$\frac{p'_{i+1}(x)}{p_{i+1}(x)} = \frac{1}{2\sqrt{x}} \left(\frac{p'_i(\sqrt{x})}{p_i(\sqrt{x})} - \frac{p'_i(-\sqrt{x})}{p_i(-\sqrt{x})} \right), \quad i = 0, 1, \dots \quad (8)$$

Recurrences (7) and (8) reduce the evaluation of $p_\ell(c)$ to the evaluation of $p(c)$ at $q = 2^\ell$ points $c^{1/q}$ and for $c \neq 0$ reduce the evaluation of the ratio $p'_\ell(x)/p_\ell(x)$ at $x = c$ to the evaluation of the ratio $p'(x)/p(x)$ at the latter $q = 2^\ell$ points $x = c^{1/q}$. We will see that we can apply recurrence (8) to support fast convergence to the convex hull of the roots.

For $x = 0$, the recurrence (8) can be specialized as follows:

$$\frac{p'_\ell(0)}{p_\ell(0)} = \left(\frac{p'_{\ell-1}(x)}{p_{\ell-1}(x)} \right)'_{x=0} = \left(\frac{p'(x)}{p(x)} \right)^{(\ell)}_{x=0}, \quad \ell = 1, 2, \dots \quad (9)$$

Eq. (9) enables us to strengthen upper estimates in Eq. 10 for root radii $r_j(0, p)$ at the origin because $r_j(0, p_\ell) = r_j(0, p)^{2^\ell}$ for $j = 1, \dots, d$ (see Eq. 10); we can

approximate the higher order derivatives $\left(\frac{p^{(g)}(x)}{p^{(g-1)}(x)}\right)^{(\ell)}$ at $x = 0$ by following Remark. 3. Besides the listed applications of root-squaring, one can apply DLG iterations to randomized exclusion tests for sparse polynomials. One can apply root-squaring $p(x) \mapsto p_\ell(x)$ to improve the error bound for the approximation of the power sums of the roots of $p(x)$ in the unit disc $D(0, 1)$ by Cauchy sums, but the improvement is about as much as the additional cost incurred by increasing the number q of points of evaluation of the ratio $\frac{p'}{p}$.

Remark 2. One can approximate the leading coefficient p_d of a black box polynomial $p(x)$. This coefficient is not involved in recurrence (8), and one can apply recurrence (7) by using a crude approximation to p_d and if needed can scale polynomials $p_i(x)$ for some i .

We can treat $\frac{p'(x)}{p(x)}$ as another blackbox that runs the same order of cost since the same blackbox oracle that evaluates $p(x)$ can be used to evaluate $p'(x)$ as well, based on the following theorem:

Theorem 6. *Given an algorithm that evaluates a black box polynomial $p(x)$ at a point x over a field \mathcal{K} of constants by using A additions and subtractions, S scalar multiplications (that is, multiplications by elements of the field \mathcal{K}), and M other multiplications and divisions, one can extend this algorithm to the evaluation at x of both $p(x)$ and $p'(x)$ by using $2A + M$ additions and subtractions, $2S$ scalar multiplications, and $3M$ other multiplications and divisions.*

Proof. [11] and [1] prove the theorem for any function $f(x_1, \dots, x_s)$ that has partial derivatives in all its s variables x_1, \dots, x_s .

Remark 3. Given a complex c and a positive integer ℓ one can approximate the values at $x = c$ of $(p(x)/p'(x))^{(\ell)}$ for a black box polynomial $p(x)$ by using divided differences, by extending expressions $p^{(i+1)}(c) = \lim_{x \rightarrow c} \frac{p^{(i)}(x) - p^{(i)}(c)}{x - c}$ for $i = 0, 1, \dots, \ell - 1$ and based on the mean value theorem [4].

A.2 Estimation of extremal root radii

If we do not know whether assumptions (6) holds, we can apply the following well-known bounds on the extremal root radii:

$$|x_d| \leq d \left| \frac{p(0)}{p'(0)} \right| \text{ and } |x_1| \geq \left| \frac{p'_{\text{rev}}(0)}{d p_{\text{rev}}(0)} \right|. \quad (10)$$

We can deduce these bounds from the well-known expression

$$\frac{p'(x)}{p(x)} = \sum_{j=1}^d \frac{1}{x - x_j}, \quad (11)$$

which we can obtain by differentiating the equation $p(x) = p_d \prod_{j=1}^d (x - x_j)$. By extending these bounds to the polynomial $p_k(x)$ of Eq. (7) we obtain that

$$|x_d|^{2^k} \leq d / \left| \left(\frac{p'(x)}{p(x)} \right)_{x=0}^{(k)} \right| \text{ and } |x_1|^{2^k} \geq \frac{1}{d} \left| \left(\frac{p'_{\text{rev}}(x)}{p_{\text{rev}}(x)} \right)_{x=0}^{(k)} \right|. \quad (12)$$

Under the assumptions (6) for $i = 2$ for $p(x)$ and for $p_{\text{rev}}(x)$, respectively, the latter two bounds become sharp as k increases, by virtue of (10), and next we argue informally that it tends to be sharp with a high probability under random root models. Indeed,

$$\frac{1}{|x_d|} \leq \frac{1}{d} \left| \frac{p'(c)}{p(c)} \right| = \frac{1}{d} \left| \sum_{j=1}^d \frac{1}{c - x_j} \right| \quad (13)$$

by virtue of (10), and so the approximation to the root radius $|x_d|$ is poor if and only if severe cancellation occurs in the summation of the d roots, and similarly for the approximation of $r_1(c, p)$. Such a cancellation only occurs for a narrow class of polynomials $p(x)$, with a low probability if we assume a random root model.

Next we prove, however, that estimates (10) and (12) are extremely poor for worst case inputs.

Theorem 7. *The ratios $\left| \frac{p(0)}{p'(0)} \right|$ and $\left| \frac{p_{\text{rev}}(0)}{p'_{\text{rev}}(0)} \right|$ are infinite for $p(x) = x^d - h^d$ and $h \neq 0$, while $|x_d| = |x_1| = |h|$.*

Proof. Observe that the roots $x_j = h \exp(\frac{(j-1)\mathbf{i}}{2\pi d})$ of $p(x) = x^d - h^d$ for $j = 1, 2, \dots, d$ are the d th roots of unity up to scaling by h .

The problem persists for the root radius $r_d(w, p)$ where $p'(w)$ and $p'_{\text{rev}}(w)$ vanish; rotation of the variable $p(x) \leftarrow t(x) = p(ax)$ for $|a| = 1$ does not fix it but shifts $p(x) \leftarrow t(x) = p(x - c)$ for $c \neq 0$ can fix it, thus *enhancing the power of estimates (10) and (12)*.