

Supporting Information

Cecala, J.M. and Wilson Rankin, E.E. Wild bee functional diversity and plant associations in native and conventional plant nurseries.

Contents

Table S1. The 13 plant nurseries sampled and their physical characteristics.

Table S2. Sampling schedule for the 13 nurseries in the study from March 2016 to May 2018.

Table S3. List of bee species and morphotaxa collected in this study.

Table S4. Results of significance tests for fixed effects from mixed models.

Table S5. Model selection for mixed models.

Table S6. Numbers of bee species detected, categorized by three functional traits of interest.

Table S7. List of flowering plant species off which bees were collected.

Fig. S1. View of a sampling plot with blue vane trap deployed at a plant nursery.

Fig. S2. Diet breadth of bee individuals collected by sweep netting and vane trapping.

Fig. S3. Nesting location of bee individuals collected by sweep netting and vane trapping.

Fig. S4. Social organization of bee individuals collected by sweep netting and vane trapping.

Table S1. The 13 plant nurseries sampled and their physical characteristics. Also included are the total numbers of individuals and species of bees collected by sweep netting and vane trapping, respectively, at each nursery.

Nursery	Type	County (California, USA)	Area (ha)	Number of plots	Proportion natural area (1-km buffer)	Elevation (m)	Sweep netted bees	Vane trapped bees
01	conventional	Riverside	4.00	4	0.257	316	61 bees in 19 spp.	1,437 bees in 28 spp.
02	conventional	Riverside	2.90	3	0.210	303	44 bees in 18 spp.	524 bees in 26 spp.
03	conventional	Riverside	3.89	4	0.289	580	7 bees in 5 spp.	245 bees in 22 spp.
04	conventional	Riverside	3.80	4	0.081	523	0 bees	17 bees in 8 spp.
05	conventional	Riverside	30.00	12	0.562	471	21 bees in 10 spp.	1,098 bees in 41 spp.
06	native	San Diego	0.50	2	0.711	105	5 bees in 4 spp.	5 bees in 5 spp.
07	native	San Diego	4.83	4	0.915	471	50 bees in 23 spp.	386 bees in 37 spp.
08	native	Orange	7.30	7	0.793	114	42 bees in 23 spp.	751 bees in 53 spp.
09	conventional	Los Angeles	4.58	4	0.106	151	44 bees in 21 spp.	1,004 bees in 37 spp.
10	conventional	Los Angeles	4.29	4	0.151	104	10 bees in 8 spp.	147 bees in 20 spp.
11	native	Los Angeles	3.10	3	0.301	303	52 bees in 29 spp.	185 bees in 29 spp.
12	native	San Diego	7.21	7	0.300	12	22 bees in 12 spp.	319 bees in 22 spp.
13	conventional	Riverside	3.86	4	0.614	775	17 bees in 12 spp.	194 bees in 30 spp.
TOTAL							375 bees in 92 spp.	6,312 bees in 102 spp.

Table S2. Sampling schedule for the 13 nurseries in the study from March 2016 to May 2018. A “1” indicates that nursery was sampled, while a “0” indicates it was not sampled due to logistical reasons. Blacked-out cells indicate seasons when sampling was not possible due to closure of nursery property. Columns sums (bottom row) indicate the total number of nurseries sampled in that season, while row sums (rightmost column) indicate the total number of times that nursery was sampled in the study.

Nursery	2016			2017			2018	samplings per nursery
	Spring	Summer	Autumn	Spring	Summer	Autumn	Spring	
01	1	1	1	1	1	1	1	7
02	1	1	1	1	1	1	1	7
03	1	1						2
04	1	0	0	1				2
05	1	1	1	1	1	0	0	5
06	1							1
07	1	1	1	1	1	0	1	6
08	0	1	1	1	1	1	1	6
09		1	1	1	1	1	1	6
10	0	1	1	1	1	1	1	6
11	0	1	1	1	1	0	1	5
12	0	0	0	1	1	0	1	3
13	0	0	0	0	1	0	1	2
nurseries sampled	7	9	8	10	10	5	9	58

Table S3. List of bee species and morphotaxa, ordered by family and genus, collected in this study via sweep netting and blue vane traps. Morphotaxa are indicated by a species epithet consisting of “sp.” followed by a unique identifier. Each species’ diet breadth, nesting location, and social organization is designated along with the corresponding literature reference as a bracketed number. Species-level information on functional traits was not available for all species in the study, or, of course, for morphotaxa. In such cases, functional traits were assigned based on inference from their taxonomy at the genus level (e.g., Michener (2007) notes that “all *Andrena* species nest in the ground”). Instances where functional traits were assigned in this manner are indicated by table cells merged across species within the column designating the trait. Lastly, the numbers of individual bees (by sex and collection method) collected in the study are listed at the right. In the “subgenus” column: [unk] = taxon was not identifiable to a described subgenus; [NA] = subgenera are not currently broadly established for this genus.

Taxonomy				Functional traits			sweep netting		vane trapping	
family	genus	subgenus	species	diet breadth	nesting location	social organization	females	males	females	males
Andrenidae	Andrena	Plastandrena	Andrena prunorum	generalist[1]	below ground[2]	solitary[2]	0	1	0	5
		Simandrena	Andrena angustitarsata	generalist[3]			1	0	1	0
		[unk]	Andrena sp. 2c-A	specialist[4]			0	0	5	1
		[unk]	Andrena sp. 2c-B				0	0	2	0
		[unk]	Andrena sp. 3c-B				0	0	2	0
		[unk]	Andrena sp. 3c-C				0	0	1	2
		[unk]	Andrena sp. 3c-D				0	10	0	2
	Calliopsis	Calliopsis	Calliopsis rhodophila	generalist[5]	below ground[2]	solitary[6]	4	1	11	1
		Nomadopsis	Calliopsis scutellaris	generalist[3]			2	1	0	0
	Panurginus	[NA]	Panurginus bilobatus	specialist[3]	below ground[2]	solitary[6]	2	1	5	3
			Panurginus sp. B				0	0	1	0
	Perdita	Perdita	Perdita rhois	generalist[3]	below ground[2]	solitary[6]	1	5	0	0
		Pygoperdita	Perdita interrupta	specialist[3]			1	4	0	0
		Pygoperdita	Perdita micheneri	specialist[2]			3	1	1	2
		Pygoperdita	Perdita mojavensis	specialist[3]			0	0	1	1

		[unk]	<i>Perdita</i> sp. B	specialist[2]			0	0	2	3
		[unk]	<i>Perdita</i> sp. D				0	0	4	0
		[unk]	<i>Perdita</i> sp. E				0	0	2	0
		[unk]	<i>Perdita</i> sp. F				2	0	5	3
Apidae	<i>Anthophora</i>	<i>Anthophoroides</i>	<i>Anthophora californica</i>	generalist [7]	below ground[7]	solitary[2]	9	7	4	7
		<i>Micranthophora</i>	<i>Anthophora curta</i>	generalist[3]	below ground[2]	solitary[2]	1	0	0	0
		<i>Mystacanthophora</i>	<i>Anthophora urbana</i>	generalist[3]			6	0	126	10
		[unk]	<i>Anthophora</i> sp. A	generalist[3]			1	0	0	0
	<i>Anthophorula</i>	<i>Anthophorisca</i>	<i>Anthophorula nitens</i>	generalist[8]	below ground[8]	solitary[8]	0	0	0	1
		<i>Anthophorula</i>	<i>Anthophorula albicans</i>	generalist[3]	below ground[2]	solitary[2]	0	0	1	0
		<i>Anthophorula</i>	<i>Anthophorula cerei</i>	generalist[3]			0	0	1	0
		<i>Anthophorula</i>	<i>Anthophorula torticornis</i>	generalist[3]			0	0	1	1
	<i>Apis</i>	<i>Apis</i>	<i>Apis mellifera</i>	generalist[2]	cavity[2]	eusocial[6]	10	0	633	0
	<i>Bombus</i>	<i>Bombias</i>	<i>Bombus nevadensis</i>	generalist[3]	below ground[3]	eusocial[3]	0	1	0	0
		<i>Cullumanobombus</i>	<i>Bombus crotchii</i>				0	0	3	0
		<i>Pyrobombus</i>	<i>Bombus melanopygus</i>				2	0	1	0
		<i>Pyrobombus</i>	<i>Bombus vandykei</i>				0	0	1	0
		<i>Pyrobombus</i>	<i>Bombus vosnesenskii</i>				1	0	10	1
		<i>Thoracobombus</i>	<i>Bombus californicus</i>				1	3	4	4
		<i>Thoracobombus</i>	<i>Bombus sonorus</i>				2	0	12	1
	<i>Ceratina</i>	<i>Ceratinula</i>	<i>Ceratina arizonensis</i>	generalist[3]	cavity[2]	solitary[2]	0	0	2	1
		<i>Zadontomerus</i>	<i>Ceratina acantha</i>	generalist[3]			9	3	40	0
		<i>Zadontomerus</i>	<i>Ceratina micheneri</i>	generalist[3]			0	0	1	0
		<i>Zadontomerus</i>	<i>Ceratina nanula</i>	generalist[3]			1	0	9	0
		<i>Zadontomerus</i>	<i>Ceratina sequoiae</i>	specialist[3]			0	0	1	0

	<i>Diadasia</i>	<i>Coquillettapis</i>	<i>Diadasia angusticeps</i>	specialist[9]	below ground[2]	solitary[9]	0	0	1	0
		<i>Coquillettapis</i>	<i>Diadasia australis</i>				1	1	20	26
		<i>Coquillettapis</i>	<i>Diadasia bituberculata</i>				1	0	9	6
		<i>Coquillettapis</i>	<i>Diadasia diminuta</i>				0	5	3	3
		<i>Coquillettapis</i>	<i>Diadasia laticauda</i>				0	0	13	4
		<i>Coquillettapis</i>	<i>Diadasia lutzi</i>				0	0	1	0
		<i>Coquillettapis</i>	<i>Diadasia nitidifrons</i>				0	4	8	2
		<i>Coquillettapis</i>	<i>Diadasia rinconis</i>				0	0	125	42
		<i>Coquillettapis</i>	<i>Diadasia vallicola</i>				2	0	0	0
		<i>Dasiapis</i>	<i>Diadasia ochracea</i>				1	0	22	0
		<i>Diadasia</i>	<i>Diadasia enavata</i>				0	0	2	0
	<i>Eucera</i>	<i>Synhalonia</i>	<i>Eucera cordleyi</i>	generalist[3]	below ground[2]	solitary[2]	0	0	1	3
		<i>Synhalonia</i>	<i>Eucera dorsata</i>	generalist[3]			0	0	1	0
		<i>Synhalonia</i>	<i>Eucera lunata</i>	generalist[3]			0	0	1	0
	<i>Habropoda</i>	[NA]	<i>Habropoda depressa</i>	generalist[10]	below ground[10]	solitary[10]	2	0	0	0
			<i>Habropoda tristissima</i>	generalist[10]	below ground[10]	solitary[10]	0	0	8	0
	<i>Melecta</i>	<i>Melecta</i>	<i>Melecta pacifica</i>	cleptoparasitic[2]]	cleptoparasitic[2]]	cleptoparasitic[2]]	0	0	1	0
	<i>Melissodes</i>	[unk]	<i>Melissodes</i> sp. 4	generalist[2]	below ground[2]	solitary[11]	0	0	1	0
		[unk]	<i>Melissodes</i> sp. A				3	12	202	133
		[unk]	<i>Melissodes</i> sp. B				1	3	103	17
		[unk]	<i>Melissodes</i> sp. C				1	4	78	107
		[unk]	<i>Melissodes</i> sp. D				4	4	31	62
	<i>Nomada</i>	[NA]	<i>Nomada</i> sp. A	cleptoparasitic[2]	cleptoparasitic[2]	cleptoparasitic[2]	1	1	0	0

				]	]	]				
	<i>Oreopasites</i>	[unk]	<i>Oreopasites</i> sp. 1	cleptoparasitic[2]]	cleptoparasitic[2]]	cleptoparasitic[2]]	0	0	0	1
		[unk]	<i>Oreopasites</i> sp. A				0	0	3	0
	<i>Peponapis</i> *	<i>Peponapis</i>	<i>Peponapis pruinosa</i>	specialist[2]	below ground[7]	solitary[12]	0	0	53	116
	<i>Svastra</i>	<i>Epimelissodes</i>	<i>Svastra sabinensis</i>	generalist[13]	below ground[14]	solitary[13]	0	2	0	0
	<i>Triepeolus</i>	[NA]	<i>Triepeolus concavus</i>	cleptoparasitic[2]]	cleptoparasitic[2]]	cleptoparasitic[2]]	0	1	0	0
			<i>Triepeolus</i> sp. B				2	1	0	0
	<i>Xenoglossa</i>	<i>Eoxenoglossa</i>	<i>Xenoglossa strenua</i>	specialist[2]	below ground[2]	solitary[2]	0	0	72	103
		<i>Xenoglossa</i>	<i>Xenoglossa patricia</i>				0	0	1	3
	<i>Xeromelecta</i>	<i>Melectomorpha</i>	<i>Xeromelecta californica</i>	cleptoparasitic[2]]	cleptoparasitic[2]]	cleptoparasitic[2]]	0	0	0	1
Colletidae	<i>Colletes</i>	[NA]	<i>Colletes</i> sp. A1	generalist[3]	below ground[2]	solitary[2]	1	0	0	0
			<i>Colletes</i> sp. A2				1	0	0	0
			<i>Colletes</i> sp. B				3	0	0	0
	<i>Hylaeus</i>	<i>Hylaeus</i>	<i>Hylaeus mesillae</i>	generalist[2]	cavity[2]	solitary[2]	2	12	2	1
Halictidae	<i>Agapostemon</i>	<i>Agapostemon</i>	<i>Agapostemon femoratus</i>	generalist[16]	below ground[2]	solitary[17]	0	0	28	1
		<i>Agapostemon</i>	<i>Agapostemon melliventris</i>	generalist[16]		solitary[17]	1	1	274	16
		<i>Agapostemon</i>	<i>Agapostemon texanus</i>	generalist[16]		solitary[18]	12	17	1692	112
	<i>Augochlorella</i>	<i>Augochlorella</i>	<i>Augochlorella pomoniella</i>	generalist[19]	below ground[20]	eusocial[21]	1	2	22	3
	<i>Conanthalictus</i>	<i>Phaceliapis</i>	<i>Conanthalictus cockerelli</i>	specialist[2]	below ground[22]	solitary[23]	0	0	0	1
	<i>Dieunomia</i>	<i>Epinomia</i>	<i>Dieunomia nevadensis</i>	generalist[24]	below ground[24]	solitary[24]	1	1	0	0
	<i>Dufourea</i>	[NA]	<i>Dufourea</i> sp. 2	specialist[25]	below	solitary[23]	1	0	8	2

			<i>Dufourea trochantera</i>		ground[23]		0	1	308	39
	<i>Halictus</i>	<i>Nealictus</i>	<i>Halictus farinosus</i>	generalist[26]	below ground[26]	eusocial[27]	0	0	36	3
		<i>Odontalictus</i>	<i>Halictus ligatus</i>	generalist[28]	below ground[28]	eusocial[28]	40	15	40	2
		<i>Seladonia</i>	<i>Halictus tripartitus</i>	generalist[29]	below ground[30]	eusocial[30]	9	0	198	0
	<i>Lasioglossum</i>	<i>Dialictus</i>	<i>Lasioglossum</i> sp. dia1	generalist[2]	below ground[2]	eusocial[2]	1	0	0	0
			<i>Lasioglossum</i> sp. dia10				1	0	0	0
			<i>Lasioglossum</i> sp. dia11				1	0	0	0
			<i>Lasioglossum</i> sp. dia12				1	0	0	0
			<i>Lasioglossum</i> sp. dia2				11	0	0	0
			<i>Lasioglossum</i> sp. dia3				0	4	0	0
			<i>Lasioglossum</i> sp. dia4				4	0	0	0
			<i>Lasioglossum</i> sp. dia5				1	0	0	0
			<i>Lasioglossum</i> sp. dia6				1	0	0	0
			<i>Lasioglossum</i> sp. dia7				2	1	0	0
			<i>Lasioglossum</i> sp. dia8				5	1	0	0
			<i>Lasioglossum</i> sp. dia9				1	0	0	0
			<i>Lasioglossum</i> sp. diaA				0	8	0	0
			<i>Lasioglossum</i> sp. diaD				0	1	0	0
			<i>Lasioglossum</i> sp. diaE				0	1	0	0
			<i>Lasioglossum</i> sp. disA				0	0	980	10
			<i>Lasioglossum</i> sp. disB				0	0	66	0
			<i>Lasioglossum</i> sp. disD				0	0	5	0
			<i>Lasioglossum</i> sp. disE				0	0	333	0
		<i>Evyllaenus</i>	<i>Lasioglossum</i> sp. EV01	generalist[2]	below ground[2]	eusocial[2]	0	0	208	3

			<i>Lasioglossum</i> sp. EV02				5	0	19	0
			<i>Lasioglossum</i> sp. EV03				1	0	42	1
			<i>Lasioglossum</i> sp. EV04				0	0	3	0
			<i>Lasioglossum</i> sp. EV05				0	0	1	0
			<i>Lasioglossum</i> sp. EV06				1	0	3	0
			<i>Lasioglossum</i> sp. EV07				0	0	2	0
	<i>Lasioglossum</i>		<i>Lasioglossum</i> sp. LA02	generalist[2]	below ground[2]	eusocial[2]	0	0	3	0
			<i>Lasioglossum sisymbrii</i>	generalist[31]	below ground[32]	solitary[33]	4	0	49	0
	<i>Micralictoides</i>	[NA]	<i>Micralictoides ruficaudus</i>	specialist[34]	below ground[35]	solitary[35]	4	1	4	1
	<i>Sphecodes</i>	[NA]	<i>Sphecodes</i> sp. A	cleptoparasitic[2]	cleptoparasitic[2]	cleptoparasitic[2]	1	1	0	1
Megachilidae	<i>Anthidium</i>	<i>Anthidium</i>	<i>Anthidium placitum</i>	specialist[36]	below ground[36]	solitary[36]	0	0	1	0
	<i>Ashmeadiella</i>	<i>Ashmeadiella</i>	<i>Ashmeadiella buconis</i>	generalist[37]	cavity[37]	solitary[6]	9	2	0	0
		<i>Ashmeadiella</i>	<i>Ashmeadiella cubiceps</i>	generalist[3]			1	0	0	0
		<i>Ashmeadiella</i>	<i>Ashmeadiella difugita</i>	generalist[3]			2	0	0	0
		<i>Ashmeadiella</i>	<i>Ashmeadiella meliloti</i>	generalist[3]			1	0	0	0
		<i>Ashmeadiella</i>	<i>Ashmeadiella</i> sp. 1	generalist[3]			0	0	0	1
		<i>Cubitognatha</i>	<i>Ashmeadiella salviae</i>	specialist[37]			2	0	0	0
	<i>Atoposmia</i>	<i>Eremosmia</i>	<i>Atoposmia</i> sp. A	specialist[3]	cavity[3]	solitary[6]	0	0	1	0
	<i>Dianthidium</i>	<i>Dianthidium</i>	<i>Dianthidium parvum</i>	generalist[3]	cavity[3]	solitary[6]	0	0	1	0
		<i>Dianthidium</i>	<i>Dianthidium pudicum</i>	generalist[3]	cavity[3]		1	0	1	0
	<i>Hoplitis</i>	[unk]	<i>Hoplitis</i> sp. 1	generalist[3]	cavity[2]	solitary[6]	0	0	1	0
		[unk]	<i>Hoplitis</i> sp. 2				0	0	4	2
	<i>Megachile</i>	<i>Litomegachile</i>	<i>Megachile brevis</i>	generalist[38]	cavity[38]	solitary[38]	1	0	0	0
		<i>Sayapis</i>	<i>Megachile inimca</i>	generalist[3]	cavity[2]	solitary[2]	2	0	0	0

				generalist[3]	cavity[2]	solitary[2]				
		[unk]	<i>Megachile</i> sp. 2				0	1	0	0
		[unk]	<i>Megachile</i> sp. 3a				0	1	0	0
		[unk]	<i>Megachile</i> sp. 3b				0	1	0	0
		[unk]	<i>Megachile</i> sp. 4				0	0	0	1
		[unk]	<i>Megachile</i> sp. 5				0	0	0	1
		[unk]	<i>Megachile</i> sp. D				1	0	0	0
		[unk]	<i>Megachile</i> sp. E				1	0	2	0
		[unk]	<i>Megachile</i> sp. F				1	0	0	0
		[unk]	<i>Megachile</i> sp. G				0	0	2	0
		[unk]	<i>Megachile</i> sp. H				0	0	3	0
		[unk]	<i>Megachile</i> sp. I				0	0	1	0
		[unk]	<i>Megachile</i> sp. J				0	0	1	0
		[unk]	<i>Megachile</i> sp. X				1	4	0	0
	<i>Osmia</i>	[unk]	<i>Osmia</i> sp. 1	generalist[3]	cavity[2]	solitary[2]	0	1	0	0
		[unk]	<i>Osmia</i> sp. A				3	0	0	0
		[unk]	<i>Osmia</i> sp. C				1	0	0	0
		[unk]	<i>Osmia</i> sp. D				1	0	0	0
		[unk]	<i>Osmia</i> sp. E				1	0	0	0
		[unk]	<i>Osmia</i> sp. F				2	0	0	0
		[unk]	<i>Osmia</i> sp. G				0	0	2	0
		[unk]	<i>Osmia</i> sp. H				0	0	1	0

References for Table S3

1. Tepedino, V. J., Alston, D. G., Bradley, B. A., Toler, T. R. & Griswold, T. L. 2007 Orchard pollination in Capitol Reef National Park, Utah, USA. Honey bees or native bees? *Biodiversity and Conservation* **16**, 3083–3094. (doi:10.1007/s10531-007-9164-8)
2. Michener, C. D. 2007 *The Bees of the World*. 2nd edn. Baltimore, MD: The Johns Hopkins University Press.
3. Krombein, K. V., Hurd, P. D., Smith, D. R. & Burks, B. D. 1979 *Catalog of Hymenoptera in America North of Mexico*. Washington, D.C.: Smithsonian Press.
4. Larkin, L. L., Neff, J. L. & Simpson, B. B. 2008 The evolution of a pollen diet: Host choice and diet breadth of *Andrena* bees (Hymenoptera: Andrenidae). *Apidologie* **39**, 133–145. (doi:10.1051/apido:2007064)
5. Wyatt, R. 1986 Ecology and evolution of self-pollination in *Arenaria uniflora* (Caryophyllaceae). *Journal of the Kansas Entomological Society* **74**, 403. (doi:10.2307/2260264)
6. Danforth, B. N., Cardinal, S., Praz, C., Almeida, E. A. B. & Michez, D. 2013 The Impact of Molecular Data on Our Understanding of Bee Phylogeny and Evolution. *Annual Review of Entomology* **58**, 57–78. (doi:10.1146/annurev-ento-120811-153633)
7. Roulston, T. H. & Cane, J. H. 2000 The Effect of Diet Breadth and Nesting Ecology on Body Size Variation in Bees (Apiformes). *Journal of the Kansas Entomological Society* **73**, 129–142. (doi:10.2307/25085957?refreqid=search-gateway:87a9a826f22082f7c787cafla81af99c)
8. Rozen, J. G. & Snelling, R. R. 1986 Ethology of the Bee *Exomalopsis nitens* and Its Cleptoparasite (Hymenoptera: Anthophoridae). *Journal of the Kansas Entomological Society* **94**, 480–488. (doi:10.2307/25009567?refreqid=search-gateway:88d7988eb7ff3a52c8c43d2a2606a493)
9. Sipes, S. D. & Tepedino, V. J. 2005 Pollen-host specificity and evolutionary patterns of host switching in a clade of specialist bees (Apoidea: *Diadasia*). *Journal of Economic Entomology* **86**, 487–505.
10. Barthell, J. F., Bromberger, D. M., Daly, H. V. & Thorp, R. W. 1998 Nesting biology of the solitary digger bee *Habropoda depressa* (Hymenoptera: Anthophoridae) in urban and island environments. *Journal of the Kansas Entomological Society*, 116–136.
11. Estravis Barcala, M. C., Palottini, F. & Farina, W. M. 2019 Honey bee and native solitary bee foraging behavior in a crop with dimorphic parental lines. *PLoS ONE* **14**, e0223865–14. (doi:10.1371/journal.pone.0223865)
12. Willis, D. S. & Kevan, P. G. 1995 Foraging dynamics of *Peponapis pruinosa* (Hymenoptera: Anthophoridae) on pumpkin (*Cucurbita pepo*) in southern Ontario. *The Canadian Entomologist* **127**, 167–175.
13. Cane, J. H. 1995 Notes on the Nesting Biology of *Svastra atripes atrimitra* (LaBerge) (Hymenoptera: Apidae). *Journal of the Kansas Entomological Society* **68**, 238–240. (doi:10.2307/25085587?refreqid=search-gateway:bc055d60770d07f71f322c5df4fc571c)
14. Rozen, J. G. 1983 Nesting Biology of the Bee *Svastra sabinensis* (Hymenoptera, Anthophoridae). *Journal of the Kansas Entomological Society* **91**, 264–268.
15. Waller, G. D., Vaissière, B. E., Moffett, J. O. & Martin, J. H. 1985 Comparison of Carpenter Bees (*Xylocopa varipuncta* Patton) (Hymenoptera: Anthophoridae) and Honey Bees (*Apis mellifera* L.) (Hymenoptera: Apidae) as Pollinators of Male-sterile Cotton in Cages1. *Journal of Economic Entomology* **78**, 558–561. (doi:10.1093/jee/78.3.558)
16. Porter, C. C. 1983 Ecological Notes on Lower Rio Grande Valley *Augochloropsis* and *Agapostemon* (Hymenoptera: Halictidae). *The Florida Entomologist* **66**, 344–353. (doi:10.2307/3494129)
17. Janjic, J. & Packer, L. 2003 Phylogeny of the bee genus *Agapostemon* (Hymenoptera: Halictidae). *Systematic Entomology* **28**, 101–124. (doi:10.1046/j.1365-3113.2003.00204.x)
18. Eickwort, G. C. 1981 Aspects of the Nesting Biology of Five Nearctic Species of *Agapostemon* (Hymenoptera: Halictidae). *Journal of the Kansas Entomological Society* **54**, 337–351.
19. Barthell, J. F., Randall, J. M., Thorp, R. W. & Wenner, A. M. 2001 Promotion of seed set in yellow star-thistle by honey bees: evidence of an invasive mutualism. *Ecological Applications* **11**, 1870–1883.
20. Ordway, E. 1964 *Sphecodes pimpinellae* and Other Enemies of *Augochlorella* (Hymenoptera: Halictidae). *Journal of the Kansas Entomological Society* **37**, 139–152. (doi:10.2307/25083373?refreqid=search-gateway:dce19fe54452f20071bf1537412db0be)
21. Coelho, B. 2004 A review of the bee genus *Augochlorella* (Hymenoptera : Halictidae : Augochlorini). *Systematic Entomology* **29**, 282–323. (doi:10.1111/j.0307-6970.2004.00243.x)

22. Rozen, J. G., Jr & McGinley, R. J. 1976 Biology of the bee genus *Conanthalictus* (Halictidae, Dufoureaeinae). *American Museum Novitates* **2602**, 1–6.
23. Rozen, J. G., Jr 1993 Nesting biologies and immature stages of the rophitine bees (Halictidae): with notes on the cleptoparasite *Biastes* (Anthophoridae)(Hymenoptera, Apoidea). *American Museum Novitates*
24. Wcislo, W. T. & Engel, M. S. 1996 Social behavior and nest architecture of nomiine bees (Hymenoptera: Halictidae; Nomiinae). *Journal of the Kansas Entomological Society* **69**, 158–167.
25. Patiny, S., Michez, D. & Danforth, B. N. 2008 Phylogenetic relationships and host-plant evolution within the basal clade of Halictidae (Hymenoptera, Apoidea). *Cladistics* (doi:10.1111/j.1096-0031.2007.00182.x)
26. Nye, W. P. 1980 Notes on the Biology of *Halictus* (*Halictus*) *farinosus* Smith (Hymenoptera: Halictidae). *Agricultural Research Results, Western Series* **11**.
27. Albert, J. & Packer, L. 2013 Nesting biology and phenology of a population of *Halictus farinosus* Smith (Hymenoptera, Halictidae) in northern Utah. *Journal of Hymenoptera Research* **32**, 55–73. (doi:10.3897/jhr.32.4646)
28. Richards, M. H. 2004 Annual and Social Variation in Foraging Effort of the Obligately Eusocial Sweat Bee, *Halictus ligatus* (Hymenoptera: Halictidae). *Journal of the Kansas Entomological Society* **77**, 484–502. (doi:10.2317/E37.1)
29. Pendleton, R. L., Pendleton, B. K., Wetherill, K. R. & Griswold, T. 2008 Reproductive Biology of *Larrea tridentata*: A Preliminary Comparison Between Core Shrubland and Isolated Grassland Plants at the Sevilleta National Wildlife Refuge, New Mexico. *USDA Forest Service Proceedings*, 131–135.
30. Packer, L., Gravel, A.-I. D. & LeBuhn, G. 2007 Phenology and Social Organization of *Halictus* (*Seladonia*) *tripartitus* (Hymenoptera: Halictidae). *Journal of Hymenoptera Research* **16**, 281–292.
31. Cane, J. H., Minckley, R. L., Kervin, L. J., Roulston, T. H. & Williams, N. M. 2006 Complex responses within a desert bee guild (Hymenoptera: Apiformes) to urban habitat fragmentation. *Ecological Applications* **16**, 632–644.
32. Cane, J. H. 1991 Soils of Ground-Nesting Bees (Hymenoptera, Apoidea) - Texture, Moisture, Cell Depth and Climate. *Journal of the Kansas Entomological Society* **64**, 406–413.
33. Danforth, B. N., Conway, L. & Ji, S. 2003 Phylogeny of Eusocial *Lasioglossum* Reveals Multiple Losses of Eusociality within a Primitively Eusocial Clade of Bees (Hymenoptera: Halictidae). *Systematic Biol.* **52**, 23–36. (doi:10.1080/10635150390132687)
34. Bohart, G. E. & Griswold, T. L. 1987 A revision of the Dufoureaeinae genus *Micralictoides* Timberlake (Hymenoptera: Halictidae). *Pan. Pac. Ent*
35. Pesenko, Y. A. & Astafurova, Y. V. 2006 Contributions to the halictid fauna of the Eastern Palaearctic Region: subfamily Rophitinae. *Entomofauna* **27**.
36. Pow, C. J. 2019 Host-plant specialization and nesting biology of *Anthidium placitum* (Megachilidae) in Northwest California.
37. Michener, C. D. 1939 A Revision of the Genus *Ashmeadiella* (Hymen., Megachilidae). *The American Midland Naturalist* **22**, 1–84.
38. Michener, C. D. 1953 The Biology of a Leafcutter Bee (*Megachile brevis*) and Its Associates. *Univeristy of Kansas Science Bulletin* **35**.

Table S4. Results of significance tests for fixed effects, including interaction terms, from mixed models with nursery and year as random effects. Symbols in cells indicate the results of that hypothesis test: ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$; ns: $P > 0.05$. Parentheses around “ns” indicate that, in the identical model with *Apis mellifera* included, the P -value for that hypothesis test was ≤ 0.05 . Functional dispersion (FDis) accounts for multiple functional traits, so the two models for this variable lacked a functional trait effect and associated interactions (“N/A”).

bee collection method	independent variable	numerator degrees of freedom	proportions of individuals			proportions of species			functional dispersion (FDis)
			diet breadth	nesting location	social organization	diet breadth	nesting location	social organization	
sweep netting	functional trait	2	***	**	***	***	**	*	N/A
	season	2	ns	*	*	ns	ns	ns	ns
	nursery type	1	**	ns	ns	*	ns	ns	ns
	prop. natural area at 1 km	1	ns	ns	ns	ns	ns	ns	ns
	nursery area	1	ns	ns	ns	ns	ns	ns	ns
	trait x season	4	ns	**	*	ns	ns	ns	N/A
	trait x nursery type	2	***	ns	ns	**	ns	ns	N/A
	season x nursery type	2	ns	ns	ns	ns	ns	ns	ns
	trait x season x nursery type	4	ns	ns	ns	ns	ns	ns	N/A
vane trapping	functional trait	2	***	***	***	***	***	***	N/A
	season	2	***	(ns)	***	ns	ns	ns	ns
	nursery type	1	ns	***	***	ns	ns	ns	ns
	prop. natural area at 1 km	1	ns	ns	ns	ns	ns	ns	ns
	nursery area	1	ns	ns	ns	ns	ns	ns	ns
	trait x season	4	***	(ns)	***	*	ns	ns	N/A
	trait x nursery type	2	(ns)	***	***	*	*	ns	N/A
	season x nursery type	2	***	*	***	ns	ns	ns	*
	trait x season x nursery type	4	***	**	***	ns	ns	ns	N/A

Table S6. Numbers of bee species detected via sweep netting and vane trapping, categorized by three functional traits of interest: diet breadth, nesting location, and social organization. Cleptoparasites are tallied separately.

Sweep netting (92 spp.)		eusocial	solitary
generalist	below ground	26	22
	cavity	0	25
specialist	below ground	0	14
	cavity	0	1
cleptoparasites: 4			
Vane trapping (102 spp.)		eusocial	solitary
generalist	below ground	22	22
	cavity	0	20
specialist	below ground	0	31
	cavity	0	2
cleptoparasites: 5			

Table S7. List of flowering plant species off which bees were collected, organized by their native status ($N = 375$ bees). Shown are the numbers of bees collected off each plant species at native and at conventional plant nurseries, respectively (note that a “0” does *not* necessarily imply that plant species was present at that type of nursery).

plant species name	California native status	native nurseries	conventional nurseries
<i>Salvia clevelandii</i>	native	5	25
<i>Encelia californica</i>	native	12	8
<i>Baileya multiradiata</i>	native	14	0
<i>Erigeron glaucus</i>	native	10	3
<i>Eschscholzia californica</i>	native	13	0
<i>Abutilon palmeri</i>	native	10	1
<i>Eriophyllum confertiflorum</i>	native	2	7
<i>Salvia mellifera</i>	native	8	1
<i>Ceanothus</i> sp.	native	8	0
<i>Grindelia camporum</i>	native	8	0
<i>Penstemon heterophyllus</i>	native	1	6
<i>Baccharis pilularis</i>	native	0	6
<i>Corethrogyne filaginifolia</i>	native	5	1
<i>Dendromecon harfordii</i>	native	6	0
<i>Eriogonum fasciculatum</i>	native	3	3
<i>Penstemon spectabilis</i>	native	4	2
<i>Trichostema lanatum</i>	native	6	0
<i>Malosma laurina</i>	native	5	0
<i>Sphaeralcea ambigua</i>	native	5	0
<i>Baccharis salicifolia</i>	native	3	1
<i>Rosa californica</i>	native	3	1
<i>Encelia farinosa</i>	native	0	3
<i>Epilobium canum</i>	native	3	0
<i>Eriogonum parvifolium</i>	native	3	0
<i>Fallugia paradoxa</i>	native	3	0
<i>Helianthus annuus</i>	native	3	0
<i>Isocoma menziesii</i>	native	1	2
<i>Achillea millefolium</i>	native	0	2
<i>Astragalus trichopodus</i>	native	2	0
<i>Chilopsis linearis</i>	native	0	2
<i>Clarkia unguiculata</i>	native	2	0
<i>Frangula californica</i>	native	0	2
<i>Grindelia stricta</i>	native	2	0
<i>Phyla nodiflora</i>	native	0	2
<i>Sidalcea malviflora</i>	native	2	0
<i>Tetrandeum acaulis</i>	native	2	0
<i>Bahiopsis laciniata</i>	native	1	0
<i>Calystegia macrostegia</i>	native	1	0
<i>Dichelostemma capitatum</i>	native	1	0
<i>Eriogonum grande</i>	native	1	0
<i>Helianthus californicus</i>	native	1	0
<i>Malacothamnus clementinus</i>	native	1	0
<i>Malacothamnus fasciculatus</i>	native	1	0
<i>Monardella odoratissima</i>	native	1	0
<i>Monardella villosa</i>	native	1	0

<i>Romneya coulteri</i>	native	0	1
<i>Salvia leucophylla</i>	native	0	1
<i>Salvia pachyphylla</i>	native	0	1
<i>Sambucus mexicana</i>	native	0	1
<i>Salvia greggii</i>	non-native	0	9
<i>Gaillardia grandiflora</i>	non-native	0	8
<i>Lagerstroemia indica</i>	non-native	0	8
<i>Eremophila hygrophana</i>	non-native	0	6
<i>Salvia chamaedryoides</i>	non-native	0	6
<i>Caryopteris clandonensis</i>	non-native	0	5
<i>Melampodium leucanthum</i>	non-native	1	4
<i>Vitex agnus-castus</i>	non-native	0	5
<i>Bulbine frutescens</i>	non-native	0	4
<i>Ligustrum lucidum</i>	non-native	0	4
<i>Oenothera cf. biennis</i>	non-native	0	4
<i>Rosmarinus officinalis</i>	non-native	0	4
<i>Berlandiera lyrata</i>	non-native	3	0
<i>Cistus salviifolius</i>	non-native	0	3
<i>Lamium amplexicaule</i>	non-native	3	0
<i>Oenothera speciosa</i>	non-native	0	3
<i>Pelargonium sidoides</i>	non-native	0	3
<i>Salvia microphylla</i>	non-native	1	2
<i>Salvia uliginosa</i>	non-native	0	3
<i>Tecoma capensis</i>	non-native	0	3
<i>Verbesina encelioides</i>	non-native	0	3
<i>Calandrinia spectabilis</i>	non-native	0	2
<i>Coreopsis grandiflora</i>	non-native	0	2
<i>Duranta erecta</i>	non-native	0	2
<i>Lavandula dentata</i>	non-native	0	2
<i>Lavandula stoechas</i>	non-native	0	2
<i>Leptospermum scoparium</i>	non-native	0	2
<i>Leucophyllum frutescens</i>	non-native	0	2
<i>Perovskia atriplicifolia</i>	non-native	0	2
<i>Ageratina adenophora</i>	non-native	0	1
<i>Buddleja davidii</i>	non-native	0	1
<i>Callistemon viminalis</i>	non-native	0	1
<i>Ceratostigma plumbaginoides</i>	non-native	0	1
<i>Cercidium</i> sp.	non-native	0	1
<i>Cistus purpureus</i>	non-native	0	1
<i>Convolvulus sabatius</i>	non-native	0	1
<i>Cosmos bipinnatus</i>	non-native	0	1
<i>Felicia amelloides</i>	non-native	0	1
<i>Gaura lindheimeri</i>	non-native	0	1
<i>Gnaphalieae</i> sp.	non-native	0	1
<i>Lavandula gingsinsii</i>	non-native	0	1
<i>Monarda</i> sp.	non-native	0	1
<i>Nerium oleander</i>	non-native	0	1
<i>Pallenis maritima</i>	non-native	0	1
<i>Penstemon cf. cobaea</i>	non-native	0	1
<i>Sagina subulata</i>	non-native	0	1
<i>Salvia farinacea</i>	non-native	0	1
<i>Vitex trifolia</i>	non-native	0	1



Fig. S1. View of a sampling plot with blue vane trap deployed at a conventional plant nursery in Riverside County, California, on 21 June 2016.




Fig. S2. Diet breadth of bee individuals collected by (A) sweep netting and (B) vane trapping. Proportions represent the fraction of individual bees detected a given sampling that fall into each of three functional categories—generalist, specialist, or cleptoparasite—out of all individuals collected in that given sampling. Bars and whiskers represent mean $\pm$ SE. Samplings are arranged by season and type of nursery. For sweep netting (A), the two-way interaction between diet breadth and nursery type was statistically significant. For vane trapping (B), the three-way interaction between diet breadth, season, and nursery type was statistically significant (Table S4).




Fig. S3. Nesting location of bee individuals collected by (A) sweep netting and (B) vane trapping. Proportions represent the fraction of individual bees detected a given sampling that fall into each of three functional categories— below ground, cavity, or cleptoparasite—out of all individuals collected in that given sampling. Bars and whiskers represent mean $\pm$ SE. Samplings are arranged by season and type of nursery. For sweep netting (A), the two-way interaction between nesting location and season was statistically significant. For vane trapping (B), the three-way interaction between diet breadth, season, and nursery type was statistically significant (Table S4).




Fig. S4. Social organization of bee individuals collected by (A) sweep netting and (B) vane trapping. Proportions represent the fraction of individual bees detected a given sampling that fall into each of three functional categories—solitary, eusocial, or cleptoparasite—out of all individuals collected in that given sampling. Bars and whiskers represent mean $\pm$ SE. Samplings are arranged by season and type of nursery. For sweep netting (A), the two-way interaction between sociality and season was statistically significant. For vane trapping (B), the three-way interaction between diet breadth, season, and nursery type was statistically significant (Table S4).