

Table S2: Occurrence of species at baits after 3 hours in complex and simple natural habitats in each of the regions. Abbreviations for morphological traits are: WL: Weber's length; HL: head length; FL: femur length; PW: pronotum width; HW1: head width between the eyes; HW2: head width behind the eyes; EP: eye position (EP=HW2-HW1).

Species	Habitat		Morphological trait						
	Complex	Simple	WL	HL	FL	PW	HW1	HW2	EP
S Africa									
<i>Camponotus cinctellus</i>	1	18	2.2 1	1.5 7	2.2 9	0.9 5	0.86	1.20	0.3 4
<i>Camponotus sp. A</i>	5		2.3 3	1.5 2	1.8 9	1.4 7	0.98	1.57	0.5 8
<i>Cardiocondyla sp. A</i>	2		0.4 0	0.3 7	0.3 6	0.2 2	0.25	0.33	0.0 7
<i>Lepisiota sp. A</i>	18	1	0.7 0	0.3 7	0.6 7	0.3 7	0.33	0.42	0.0 9
<i>Lepisiota sp. B</i>	15		0.7 6	0.5 1	0.7 4	0.3 7	0.33	0.42	0.0 9
<i>Meranoplus sp. A</i>	12		0.4 1	0.8 1	0.6 0	0.8 3	0.75	0.76	0.0 1
<i>Monomorium sp. E</i>	1		0.7 0	0.6 8	0.6 2	0.6 0		0.66	
<i>Monomorium sp. A</i>		86	0.7 8	0.6 8	0.4 9	0.3 8	0.49	0.59	0.1 11<
<i>Monomorium sp. B</i>	2	5	0.3 5	0.3 4	0.1 6	0.1 8	0.21	0.26	0.0 5
<i>Monomorium sp. D</i>	2		0.4 6	0.4 4	0.3 4	0.2 2	0.30	0.35	0.0 5
<i>Myrmicaria natalensis</i>	17	48	2.1 7	1.6 0	2.7 1	1.3 0	1.38	1.85	0.4 7
<i>Odontomachus sp. A</i>	3		2.8 3	2.3 9	2.5 5	1.0 8	1.52	1.97	0.4 5
<i>Pachycondyla sp. A</i>		14	1.5 9	1.2 1	1.2 7	0.8 2	1.04	1.30	0.2 6
<i>Pachycondyla sp. B</i>		3	1.9 5	1.3 3	1.2 5	0.6 3	0.67	1.16	0.4 8
<i>Pheidole megacephala</i>	96	7	0.7 4	0.6 4	0.5 5	0.3 3	0.51	0.57	0.0 7
<i>Pheidole sp. A</i>	1		1.2 1	0.8 4	1.3 5	0.4 8	0.65	0.73	0.0 8
<i>Pheidole sp. B</i>	107	2	0.3 5	0.4 8	0.4 1	0.2 8	0.37	0.43	0.0 7
<i>Pheidole sp. C</i>	30	5	0.7 4	0.6 0	0.7 7	0.3 5	0.49	0.53	0.0 4
<i>Pheidole sp. E</i>	1	3	1.2	0.9	1.4	0.5	0.82	0.93	0.1

			1	6	5	9		1
<i>Pheidole sp. F</i>	13		0.7 7	0.6 1	0.6 8	0.3 4	0.48	0.55 0.0 7
<i>Pheidole sp. G</i>		5	0.8 9	0.6 3	0.9 0	0.4 5	0.60	0.69 0.0 8
<i>Plagiolepis sp. A</i>	4		0.3 3	0.2 9	0.2 8	0.2 1	0.18	0.26 0.0 9
<i>Plagiolepis sp. B</i>	8		0.3 7	0.3 6	0.3 2	0.2 0	0.24	0.31 0.0 7
<i>Polyrhachis sp. A</i>	8		2.8 5	2.3 3	3.0 4	1.4 3	1.23	1.82 0.5 9
<i>Technomyrmex sp. A</i>	5		0.5 2	0.5 6	0.5 7	0.3 5	0.34	0.51 0.1 7
<i>Tetramorium sp. B</i>	2		0.6 0	0.5 8	0.4 7	0.3 6	0.43	0.50 0.0 7
<i>Tetramorium sp. C</i>		3	0.9 6	0.7 9	0.9 6	0.4 8	0.61	0.71 0.1 0
<i>Tetramorium sp. D</i>	2		0.6 0	0.6 0	0.5 1	0.3 9	0.53	0.65 0.1 2
SE Australia								
<i>Crematogaster sp. A</i>	69	1	1.0 1	0.8 6	0.8 7	0.5 1	0.80	0.86 0.0 6
<i>Dolichoderus sp. A</i>	1		1.4 5	0.9 4	1.1 7	0.5 4		0.74
<i>Dolichoderus sp. B</i>	1		2.1 7	1.5 1	1.7 9	1.0 9	1.02	1.33 0.3 0
<i>Iridomyrmex sp. A</i>		15	1.4 4	1.0 4	1.7 2	0.5 8	0.47	0.78 0.3 1
<i>Iridomyrmex sp. AA</i>	7		1.6 2	1.1 1	1.6 7	0.6 3	0.66	0.86 0.2 0
<i>Iridomyrmex sp. B</i>	66	205	1.0 8	0.8 6	1.0 8	0.5 1	0.43	0.79 0.3 6
<i>Iridomyrmex sp. C</i>	1	11	1.0 1	0.8 0	0.9 8	0.4 6	0.42	0.71 0.2 9
<i>Melophorus sp. B</i>		1	1.5 0	0.9 7	1.4 6	0.7 0	0.60	0.79 0.1 9
<i>Meranoplus sp. A</i>	2	5	0.7 6	0.7 9	0.6 9	0.8 0	0.74	0.80 0.0 5
<i>Meranoplus sp. B</i>		2	0.8 5	0.9 2	0.8 4	1.0 4	0.97	1.01 0.0 4
<i>Monomorium sp. A</i>	34	56	0.5 2	0.5 0	0.4 2	0.2 5	0.33	0.42 0.0 9
<i>Monomorium sp. B</i>		1	0.4 1	0.4 0	0.2 8	0.2 1	0.27	0.32 0.0 6
<i>Myrmecia sp. A</i>	1		6.8 2	3.6 9	6.9 0	2.5 1	2.75	3.82 1.0 7
<i>Notoncus sp. A</i>	4		1.5 3	1.2 0	1.4 0	0.8 2	0.84	1.14 0.3 0

<i>Ochetellus sp. A</i>	72	25	0.7 0	0.5 9	0.5 1	0.3 5	0.32	0.54	0.2 2
<i>Paratrechina sp. A</i>	14	9	0.8 2	0.6 4	0.8 2	0.4 1	0.43	0.51	0.0 8
<i>Pheidole sp. A</i>	8		0.4 9	0.4 5	0.4 3	0.2 6	0.38	0.41	0.0 3
<i>Pheidole sp. B</i>		10	0.7 4	0.6 2	0.6 5	0.3 8	0.53	0.57	0.0 4
<i>Pheidole sp. C</i>	1	1	0.7 9	0.6 3	0.7 2	0.3 7	0.54	0.59	0.0 5
<i>Prolasius sp. A</i>	1		1.1 9	0.8 1	1.1 6	0.5 0		0.64	
<i>Prolasius sp. B</i>	2		1.2 3	0.8 2	1.1 4	0.5 6	0.49	0.72	0.2 3
<i>Rhytidoponera metallica gp sp. A</i>	9	13	1.9 1	1.4 3	1.6 7	1.0 0	1.06	1.33	0.2 8
<i>Rhytidoponera metallica gp sp. B</i>	30	4	1.3 8	1.0 9	1.1 1	0.7 4	0.79	0.95	0.1 6
<i>Tapinoma sp. A</i>	28	41	0.6 1	0.5 4	0.5 2	0.2 9	0.27	0.44	0.1 7
Sweden									
<i>Camponotus herculeanus</i>	3	3	3.0 6	2.2 4	2.5 2	1.5 2	1.51	2.20	0.6 9
<i>Formica aquilonia</i>	91	116	2.2 0	1.6 9	2.0 7	1.1 3	1.08	1.51	0.4 3
<i>Formica lemani</i>	36	71	2.0 1	1.4 2	1.8 1	0.9 1	0.83	1.20	0.3 7
<i>Formica lugubris</i>	48	61	2.6 7	2.0 3	2.5 0	1.3 1	1.38	1.87	0.4 9
<i>Leptothorax acervorum</i>		4	1.1 7	0.8 8	0.7 6	0.4 4	0.68	0.75	0.0 7
<i>Myrmica lobicornis</i>	1	2	1.4 9	1.0 9	1.1 5	0.7 0	0.89	0.98	0.0 9
<i>Myrmica ruginodis</i>	90	5	1.7 7	1.2 7	1.4 4	0.7 6	0.93	1.05	0.1 2
<i>Myrmica sulcinodis</i>		21	1.6 5	1.2 5	1.3 9	0.7 6	1.01	1.14	0.1 3