



Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Ecological Indicators



journal homepage: www.elsevier.com/locate/ecolind

Original Articles

Using functional traits of chironomids to determine habitat changes in subtropical wetlands

Yanmin Cao ^{a,b,c,*}, Peter G. Langdon ^b, Shengman Shen ^a, Hui Li ^a, Deng Pan ^a ^aCollege of Resources

and Environment, South-Central Minzu University, Wuhan 430074, China ^bSchool of Geography and Environmental Science, University of Southampton,

Southampton SO17 1BJ, UK ^cHubei Key Laboratory of Critical Zone Evolution, School of Geography and Information Engineering, China University of Geosciences, Wuhan 430078, China

ARTICLE INFO

Keywords:

Functional traits

Chironomidae

Peatlands

Water level fluctuations Organic matter

research demonstrated that functional trait groups of chironomids are more robust and sensitive than taxonomy based approaches to habitat changes, and therefore could be an alternative approach for the bioassessment of aquatic ecosystem functioning and palaeo-studies in wetlands.

1. Introduction

ABSTRACT

Ecosystem functions in wetlands are increasingly degrading under the multiple stresses of climate change and human disturbances. Traditional wetland bioassessment is usually based on taxonomic approaches but this approach has limitations. To explore the effectiveness of functional traits in response to environmental changes, we compared the traditional taxonomic composition of chironomid communities with a trait-based approach in a subtropical subalpine wetland (Central China) spanning a wide habitat gradient from dry peatland to inundated peatland pools. The results revealed that 57 % of functional trait groups but only 38 % of taxonomic groups examined were significantly different between diverse peatland habitats. *Sphagnum* moss hummocks were generally inhabited by larvae of collector-gatherers, small body-sized individuals and sprawlers, while peatland pools supported a high abundance of shredders, large body-sized larvae and burrowers. Ecotones had more niche opportunities and hence possessed high taxonomic and functional diversity. Ordination analyses indicated that three similar environmental variables (loss-on-ignition (LOI), depth to water table (DWT) and K⁺) were the most powerful explanations of the chironomid variability in both taxonomic and functional trait compositions. LOI and DWT interacted strongly and were the dominant controls on both taxonomic and trait communities. Our

Environmental bioassessments have traditionally been conducted using taxonomic structures including community composition and di

Major changes in wetlands such as shrinking and degradation due to global climate change and anthropogenic drivers have occurred in recent decades (Xi et al., 2021). Such impacts are likely to degrade the ecosystem functions and services in carbon sequestration, water purification, food supply, flood control, biodiversity conservation and recreation (Hu et al., 2017; Xi et al., 2021). Knowledge of wetland functions and development processes is of crucial importance for wetland conservation and restoration. Increasingly numbers of aquatic bio indicators, such as benthic macroinvertebrates (Morse et al., 2007; Cai et al., 2023), zooplankton (Zebral et al., 2021) and algae (Peng et al., 2023) in rivers and lakes, as well as testate amoebae in peatlands

(Mitchell et al., 2000; Qin et al., 2021), are explored and commonly used in the assessment of modern and past environment and ecological functions.

iversity (Wentzky et al., 2020; Belle et al., 2022). Taxonomic approaches are capable of predicting changes in biotic communities produced by habitat conditions and environmental change, but may not always align with ecosystem functioning which primarily relies on functional characteristics of species (Nevalainen et al., 2015). Function is undoubtedly a key component of ecosystem behavior, like resilience, which would

likely be altered with human impacts (Bellwood et al., 2019). Better ways to explore functionality are therefore crucial to understand the response of ecosystems to multiple stressors. Functional trait approaches have provided new ways to assess ecosystem functioning across spatial and temporal scales (Dawson et al., 2021; Haase et al., 2023). Several studies have reported that both taxonomic compositions and functional traits were sensitive to environmental change, but interpretations were not straightforward, highlighting the importance of their combination in

* Corresponding author at: College of Resources and Environment, South-Central Minzu University, Wuhan 430074, China. E-mail address: ymcao@scuec.edu.cn (Y. Cao).

<https://doi.org/10.1016/j.ecolind.2024.111656>

Received 20 October 2023; Received in revised form 23 January 2024; Accepted 24 January 2024

Available online 2 February 2024

1470-160X/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Ecological Indicators 159 (2024) 111656

Y. Cao et al.

exploring environment-biota relationships comprehensively (Luoto and Ojala, 2018; Piano et al., 2020; Li et al., 2019). While trait approaches are considered to have relatively stable relationships with environments, taxonomic compositions may vary considerably due to biogeographic factors and taxonomic accuracy (Dolédéc et al., 2011). As such, functional trait approaches are increasingly used to provide a deeper understanding of response mechanisms of biota to environmental change (Adeleye et al., 2023).

Chironomids are prominent aquatic insects having a wide geographic distribution and outstanding capacity to resist unfavorable conditions such as extreme temperature, hydraulic changes, trophic status and substrates (Armitage et al., 2012). The larvae are highly sensitive to environmental change and typically show clear differences in community assemblages between different habitats (Brooks et al., 2007). These advantages allow chironomids (including their subfossil remains) to serve as excellent bioindicators in biological monitoring and proxies in the environmental reconstruction of aquatic systems (Armitage et al., 2012). However, the usage of taxonomy-based measures can be problematic, as outlined above, and they can vary markedly across relatively small scales (Radkova et al., 2014; Canedo-Argüelles et al., 2020). Insufficient taxonomic resolution can obscure the true biotic structure and environmental status. This problem can be magnified in large scale comparative studies, which are essential in modeling and forecasting ecological responses to future global changes (Bonada et al., 2007). Nevertheless, chironomid larvae play a crucial role in ecosystem functional processes, serving as a key node in food-web structures and biogeochemical cycles by influencing nutrient and pollutant exchanges (Kivila et al., 2019; Gautreau et al., 2023). Given the challenges with taxonomic approaches, we focus here on functional traits and functional diversity of chironomids and their role in modern bioassessment and inference of environmental history, with a novel focus on peatlands.

We selected the Dajiuhu Wetland for this study. The wetland is situated in the Shennongjia Forest region in Central China. It shows a high heterogeneity in habitat conditions and spans a gradient of *Sphagnum* moss peatlands, natural and artificial ditches, seasonal and temporary ponds, as well as artificial peatland pools. The wetland is of importance in maintaining biodiversity and regional water supply, and has been

listed in UNESCO's World Network of Biosphere Reserves since the 1990s and designated as a wetland of international importance under the Ramsar Convention in 2014. Hence the knowledge of its changes in habitat conditions and ecosystem functions would enhance wetland conservation and sustainable resource management. Here we investigated the taxonomic and functional trait structure (composition and diversity) of chironomid communities along a habitat gradient from arid *Sphagnum* moss peatlands to permanent peatland pools and verified their consistency in temporal responses to environmental changes based on a short peat core. We aimed to (1) detect the differences of chironomid community composition (both taxonomic and functional trait) between different types of habitats; (2) assess the effectiveness of taxonomic and functional trait-based measures in identifying habitat types in complex aquatic systems; and (3) determine the environmental parameters driving faunal community composition. We hypothesized that the functional traits of chironomid communities would be robust in responding to habitat variations and could be an alternative approach for contemporary bioassessment and palaeo-studies in peatland-shallow lake complexes like the Dajiuhu Wetland.

2. Material and methods

2.1. Study area

The Dajiuhu National Wetland Park lies in the mountainous area of the western Shennongjia Forest region in central China. It belongs to a closed alpine karst basin with an area of 16 km² and an altitude of 1700–1760 m a.s.l.. The region belongs to the East Asian Monsoon climate zone and is characterized by a mid-latitude subalpine humid

climate. The current mean annual temperature is ~ 7.2°C and annual precipitation is 1560 mm with maximum rainfall of 3000 mm, whilst the annual evaporation varies from 500 to 800 mm (Li et al., 2013). Cool and high humidity conditions support peatland accumulation and have led to a large coverage of thick peat (average peat coverage of 50%) in the Dajiuhu Wetland (Zhang, 1982). The Dajiuhu Wetland has been designated as an internationally important wetland under the Ramsar Convention since 2014. It is also of vital importance in water conservation

vation for the middle route of the South to North Water Diversion Project in China (Du et al., 2008). As a hotspot of biodiversity and a key ecotone for regional ecological security, the bioassessment of ecosystem health and functions for the Dajiuhu Wetland is becoming increasingly necessary, especially with the spectre of increased future climate change.

2.2. Material and laboratory analyses

We collected a total of 44 surface sediment samples in November 2019 (12 samples), September 2020 (19 samples) and July 2021 (13 samples), respectively (Fig. 1). These sampling sites occurred along a water-level gradient across diverse habitats including *Sphagnum* moss hummocks and hollows, ditches, temporary or seasonal ponds, and peatland pools. Hummocks and hollows are two distinct growth forms of *Sphagnum* moss; the former has a higher height of moss growth above the water table than the latter. Ditches here refer to flowing or still streams, and ponds are very shallow (<1 m) and small-sized (generally less than 10 m²) depressions, and store temporary water during the wet season or after rain in the dry season. Open water areas with permanent water in the wetland are classified into peatland pools in the present study (Supplementary Fig. 1). At each site, depth to water table (DWT) was measured using a meter ruler. Portable electrodes (Hach HQ40d portable multi-meter, Hach Company, Loveland, CO, USA) were employed to measure other environmental parameters in situ, including pH (Hach HQ40d multi, PHC201), conductivity (Hach HQ40d multi, CDC401), oxidation–reduction potential (ORP; Hach HQ40d multi, MTC101), dissolved oxygen and water temperature (Hach HQ40d multi, LDO101). To collect water samples, a hole was dug at arid hummock sites, or a water sampler was used for peatland pools with relatively

deep water (maximum of ~ 1.5 m in depth). For peaty habitats (both hummocks and hollows), surface peat samples were collected manually, while for inundated sites, surface sediments were retrieved using a 1/32 m² Ekman grab. Between 5 and 12 samples for each habitat were collected (Fig. 1; Table 1).

In the laboratory, concentrations of major cations (K⁺, Na⁺, Mg²⁺ and Ca²⁺) were measured using inductively coupled plasma-atomic emission spectrometry (ICP-AES; Leeman Labs Profile, Leeman Labs, Hudson, NH, USA) and morphological phosphorus (PO₄-P) and nitrogen (NO₃-N, NH₄-N and NO₂-N) in water samples were quantified by a continuous flow autoanalyzer (Skalar San Plus, Skalar Inc., Breda, The Netherlands), with the detection limit of 10 µg L⁻¹ for NH₄-N and 1 µg L⁻¹ for the other three anions. An elemental analyzer (Vario EL cube, Elementary, Hanau, Germany) was used to determine the contents of total organic carbon (TOC) and total nitrogen (TN). Weight-loss-on ignition (LOI) was calculated after the sediments were burned for 2 h at 550°C.

Chironomid head capsules were extracted from wet sediment samples following the standard approach recommended by Brooks et al (2007) and identified to genus/species type level under a microscope of 400 magnification using the taxonomy of Brooks et al (2007), with reference to Epler (2001) and Tang (2006). The criterion of 50 minimum identifiable whole head capsules per sample was satisfied to conduct subsequent analyses. To compare the temporal responses of taxonomic and functional trait compositions of chironomid communities to peat land development, chironomids were analysed from a peat core named YLC (Yangluchang Core; 51 cm-length), which was collected from the peatland central area (Fig. 1C). The chronology and biota data

Fig. 1. Geographical locations and elevations of the Shennongjia Forestry District (A), the Dajiuhu Wetland (B) and sites of the YLC (Yangluchang Core) peat core and surface samples along diverse habitats (C).

Conductivity	(5.58)	(6.39)	(6.73)	Sprawler Climber	Sprawler Climber	Canedo-Arguelles et al. (2016); Heino (2008); Serra et al. (2016)
($\mu\text{S cm}^{-1}$) 20.34 ~	128.00 (56.44)	ORP (mV) - 19.90 ~ 9.31 ~ 85.30 (25.74)	128.80 14.19 ~ 88.50 (41.30) 96.50 ~	20.18 ~ 81.60 (52.84) 73.00 ~ 65.30 ~ 151.10 (91.06)	87.40 ~ Clinger Clinger	
272.50 (140.82)	(19.39) 13.93 ~ 19.68 (17.48) 0.00 ~ 1.33 (0.44)	3.07 ~ 21.74 (8.74)	indicating their contrasting	3.2. Taxonomic composition and diversity	abundance mean value	
DO (mg L^{-1}) 0.56 ~ 7.12 (2.53)	1.08 ~ 10.71 (2.76)	10.30 ~ 16.96 (13.93) 1.03 ~ 1.88 (1.22)	environmental con ditions. The first two	A total of 4792 head capsules (HCs)	dominant genera were	
TOC (g kg^{-1}) 146.21 ~ 390.71 (310.83)	0.51 ~ 3.02 (1.32)	0.99 ~ 8.30 (3.98)	axes of the PCA	belonging to 52 genera	<i>Limnophyes</i> (16.3 %),	
TN (g kg^{-1}) 11.66 ~ 24.20 (19.24)	0.20 ~ 1.14 (0.43)	0.47 ~ 0.75 (0.61)	explained 51 % of total variation, with the first	of four subfamilies	<i>Psilome triocnemus</i> (10.1 %), <i>Chironomus</i> (6.9 %),	
C/N 14.64 ~ 29.47 (19.02)	0.01 ~ 231.50 (174.29) 0.55 ~ 9.51	0.21 ~ 0.92 (0.51)	axis explaining 35.7 % and the second one	(Tanypodinae, Chironominae,	<i>Neozavrelia</i> (5.7 %), <i>Tribelos</i> (5.3 %) and <i>Pseudosmittia</i> (5.1 %).	
K ⁺ (mg L^{-1}) 0.27 ~ 1.90 (1.06)	(5.50)	213.80 (156.24) 4.65 ~ 11.73	explaining 15.3 %. The first axis was highly	Orthocladiinae and Dia mesinae) were picked	The community composition in	
Ca ²⁺ (mg L^{-1}) 1.48 ~ 13.35 (3.85)	3.22 ~ 27.50 (10.48) 11.26 ~ 20.24 (15.22) 0.00 ~ 1.20	(7.89)	positively correlated with water pH and	from 44 surface samples (Fig. 3). Chironominae	<i>Sphagnum</i> hummocks	
Na ⁺ (mg L^{-1}) 0.22 ~ 1.90 (0.59)	(0.57)	1.33 ~ 8.34 (4.33)	cations (Mg^{2+} and Ca^{2+}), and negatively	were the most abundant subfamily occupying	was clearly different from other habitats. The	
Mg ²⁺ (mg L^{-1}) 0.17 ~ 1.50 (0.50)	- 0.09 ~ 14.15 (4.46)	9.35 ~ 20.02 (11.92) 0.28 ~ 1.34 (0.85)	correlated with LOI and DWT. The second axis	57.3 % of all HCs on average, followed by	dominant taxa were	
0.01 ~	0.40 ~ 2.59 (1.24)	8.44 ~ 16.28 (10.34) 0.41 ~ 4.46 (1.29)	showed a gradient	Orthocladiinae (40.1 %). At the genus-level,	terrestrial/semi terrestrial chironomids, such as	
NO ₃ -N (mg L ~ 280.00 (207.51) 3.18 ~ 10.84 (6.35)	0.03 ~ 1.32 (0.54)	1.01 ~ 2.13 (1.52)	associated with high concentrations of Na ⁺	Pol ypedilum showed the largest percentage	<i>Psilometriocnemus</i> (26.6 %), <i>Limnophyes</i>	
62.98 ~ 383.13 (283.68) 3.91 ~ 26.35	9.40 (7.27)	0.00 ~ hummocks and pools,	and low concentrations of K ⁺ .			
¹)	27.12 ~ 316.10 (114.98)					
NH ₄ ⁺ -N (mg L^{-1})	1.26 (0.37)	0.09 (0.04) 0.04 ~ 0.15 (0.11) 0.00 ~ 0.01	and <i>Pseudosmittia</i> (16.5 %) (Fig. 3).	absent in these relatively arid	(11.9 %) and <i>Tribelos</i> (11.1 %),	terrestrial/semi terrestrial taxa such
NO ₂ -N (mg L^{-1})	0.00 ~ 0.08 (0.01)	(0.00) 0.01 ~ 0.02 (0.01)	<i>Pseudorthocladius</i> also displayed an	hummock samples. <i>Sphagnum</i> hollow	and aquatic taxa such as <i>P.</i>	as <i>Psilometriocnemus</i> (4.9 %), <i>Neozavrelia</i>
PO ₄ -P (mg L^{-1})	0.50 (0.11) 0.00 ~ 0.46 (0.13) 0.00 ~ 0.19	0.33 (0.08) 0.00 ~ 0.78 (0.18) 0.00 ~ 0.21	increase in its frequency and abun	samples were characterized by the	<i>nubeculosum</i> (18.1 %). Although	(2.1 %), <i>Tribelos</i> (4.8 %),
0.15 (0.05)	(0.04) 0.00 ~ 0.01 (0.00)	(0.04) 0.00 ~ 0.01 (0.00)	dance compared with other habitats.	coexistence of abundant semi	<i>Limnophyes</i> (17.6 %) was dominant in and	<i>Pseudosmittia</i> (1.9 %)
0.00 ~ 1.73 (0.20)	0.55 (0.12) 0.00 ~ 0.25 (0.13) 0.00 ~ 0.09	(21.9 %), <i>Neozavrelia</i> (16.9 %)	Aquatic chironomid genera were almost	terrestrial taxa, such ditches and ponds,	<i>Psilometriocnemus</i> other	
0.00 ~	(0.01) 0.00 ~ 0.01					(0.8 %)

ranged from 4.6 % to 88.7 % with the mean values ranging from 12.2 % to 67.8 % from pools to arid *Sphagnum* hummocks respectively. Among peatland samples, three sites showed unexpectedly low LOI values: one sample from a *Sphagnum* moss hummock (29.6 % of LOI) and two from hollows (20 % and 8 % of LOI, respectively). TOC and TN, which also represent primary production, changed in a similar trend to LOI across different habitats. *Sphagnum* hollows in the peatland showed the highest acidity (mean pH of 5.58), but lowest ion concentrations (mean con

ductivity of 25.74 $\mu\text{S cm}^{-1}$). The details of ranges and averages for major environmental parameters in different habitats are given in Table 1. The PCA of environmental parameters showed that multiple surface samples overlap (Fig. 2), which was expected given that a large proportion of sites were sampled from transitional zones adjoining different habitats. Clear divergence existed between samples from peatland

were reduced in their frequency of abundance. Typical aquatic taxa, such as *Chironomus* and *Polypedilum*, occupied pool habitats with relatively high water-levels (22–147 cm in depth). *Limnophyes* abundances were reduced, and other semi-terrestrial genera were almost absent. The Kruskal-Wallis rank sum test results suggested that significant differences were shown by three subfamilies/tribes (60 % of all sub families/tribes) and 20 genera (38 %) between diverse habitats (Figs. 4 and 5). The relative abundance of Chironomini rose with the water-level in different habitats (Fig. 4), while Orthocladiinae showed an inverse relationship (Fig. 5). *Polypedilum* contributed most to the variation of Chironomini, while *Psilometriocnemus* explained the most variation of Orthocladiinae.

HC density ranged from 4 to 114 individuals g^{-1} and showed insignificant differences between the five habitats (Fig. 6A). Taxonomic richness (Taxa S) and diversity (Shannon H) in arid peatland samples

Fig. 2. Biplot of principal component analysis (PCA) showing the environmental differences between five different habitats from all the surface samples.

Fig. 3. Percentage abundance of chironomid taxa in different habitats (samples in each habitat were ordered based on DWT, so the driest samples are at the top, and the wettest at the base). The five habitats were abbreviated as follows: Hu-*Sphagnum* hummock; Ho-*Sphagnum* hollow; Di-ditch; Po-pond; Pl-pool.

were significantly lower than in other inundated habitats (Fig. 6B1 and B2). Ditches supported the most diverse assemblages, with 25 different genera on average, varying from 5 to 47 in each sample. Pools provided habitats for 21 genera while *Sphagnum* hummocks supported 8 genera on average.

3.3. Functional trait structure

For functional traits, M15 was the most common morphological type and the abundance was 34.6 % on average. Collector-gatherers (CG)

was the most common functional feeding group, and sprawlers occupied nearly 40 % across four habit types. The body size of chironomid larvae

in this study was generally medium to large. 41 % of individuals exceeded 10 mm, but no individuals were smaller than 2.5 mm. Significant differences between the five habitats were shown in 16 functional traits (57 %) verified by Kruskal-Wallis rank sum tests (Figs. 7 and 8). All five functional feeding groups displayed distinct differences between *Sphagnum* hummocks and pools (Fig. 7A1-A5). CG occupied nearly 97 % of larvae in *Sphagnum* hummocks, while

shredders (Shr) (43.6 %) was the most dominant feeding group within pools. M15 showed a generally declining tendency with water-level increases but M3 showed the reverse pattern (Fig. 7B7 and B1). Body size increased, but sprawlers decreased gradually with water-level increases (Fig. 8). Similar to taxonomic richness, ditches supported more diverse functional groups

Fig. 4. Box-Whisker plots of relative abundance of the tribe Chironomini (A) and the dominant genera (B-F) belonging to the subfamily Chironominae. The upmost, lowest and median line on the box represented the maximum, minimum and median abundance of each genus, respectively. Letters (i.e. a and b) on the bar indicated the significance between habitats tested by Kruskal-Wallis multiple comparisons.

Fig. 5. Box-Whisker plots of relative abundance of the subfamily Orthocladinae (A) and the dominant genera (B-D) having significant differences between habitats. Lines and lower-case letters in the plots are similar to [Fig. 4](#).

Fig. 6. Box-Whisker plots of HC density (A), taxonomic (B1-B2) and functional (C1-C3) diversity. Lines and lower-case letters in the plots are similar to [Fig. 4](#).

Fig. 7. Box-Whisker plots of percentages of functional traits showing significant differences between habitats. A1-A5 represents the traits of functional feeding groups and B1-B7 are trait groups of morphological type. Lines and lower-case letters in the plots are similar to Fig. 4.

Fig. 8. Box-Whisker plots of percentages of trait groups for body size (A-B) and locomotion type (C-D) showing significant differences between habitats. Lines and lower-case letters in the plots are similar to Fig. 4.

indicated by FRic (Fig. 6C2). Dissimilar to FRic, FEve and FDiv failed to show significant differences between the five habitats (Fig. 6C1 and C3).

3.4. Environmental factors controlling taxonomic and functional trait structures

The dbRDA results revealed three significant environmental variables (i.e. LOI, DWT and K^+) for both community characterizations (taxonomic and trait compositions), capturing around 39 % of the total variation in chironomid communities (Fig. 9). For the taxonomic com

munity, VPA suggested the unique effect of LOI and DWT was 10.8 % and 4.0 %, respectively, and their combined effect was 14.8 % (Fig. 10A). Similar results were shown for functional trait composition. Strong interactions also existed between LOI and DWT and occupied 18.5 % of functional trait variation. However, the unique effect was only 9.5 % for LOI and 5.5 % for DWT (Fig. 10B).

3.5. Taxonomy and traits in the YLC peat core

For taxonomic compositions, chironomids throughout the whole core were dominated by *Psilometriocnemus*, *Pseudosmittia*, *Limnophyes* and *Polypedilum* (details in Hou et al. (2022)). Major trait categories of

Fig. 10. Variation partitioning analyses revealed the unique and shared variations (%) of three significant environmental factors on taxonomic (A) and functional trait (B) compositions.

Fig. 11. Stratigraphy of major taxonomic and trait components of chironomid communities in the YLC peat core, and scores of the PC1 axis and the relations between traits and taxonomy-based PCs. Statistically significant relationships with a p value less than 0.05 or 0.01 were represented by symbol * and **, respectively. The period covered with a grey bar has no data.

chironomids are shown in Fig. 11. PCs between taxonomic and different trait compositions displayed significantly negative or positive relationships, indicating their synchronous responses to past environmental changes.

4. Discussion

4.1. Environmental variables influencing the chironomid community

Ordination analyses produced similar results for taxonomic composition and functional traits, showing that LOI, DWT and K^+ were the most important environmental factors controlling chironomid communities. LOI showed large discrepancies between different habitats. Peatlands store sequestered carbon and generally display a relatively high organic content (Leifeld and Menichetti, 2018) and hence it was

unexpected that there were three sampling sites having very low LOI values. Two of them were located near ditches, which flow from sandy areas surrounding the basin and carry high mineral content, explaining the relatively low organic matter of the nearby sites. It has been widely reported that organic matter (measured as DOC, TOC or LOI) and/or substrate type is an important control of chironomid communities in

lakes (Larocque et al., 2006; Nyman et al., 2005; Luoto et al., 2016), streams (Entrekin et al., 2007), rivers (Syrovatka and Brabec, 2006; Leszczynska et al., 2019) as well as peatlands (Van Duinen et al., 2006). The quantity of organic matter is crucial to zoobenthos biomass and community composition as it provides both food supply and habitats for organisms (Luoto et al., 2016). Organic matter exclusion in freshwater ecosystems would lead to decreased chironomid abundances and an altered chironomid community assemblage (Entrekin et al.,

2007). Nevertheless, a high organic matter content is often accompanied with low pH, poor light and depleted hypolimnetic oxygen (Luoto et al., 2016; Premke et al., 2010), which might exert an inhibitory effect to the survival of zoobenthos. Substrate quality determines the feeding behaviours of invertebrates (Henriques-Oliveira et al., 2003) and collector gatherers are often found in high humic habitats, while collector

filterers prefer relatively low organic content (Kivilä et al., 2019). Shredders generally feed on coarse particulate organic matter (CPOM > 1 mm), such as leaf litter and woody debris. However, in our study, this feeding group mainly consisted of the aquatic genus *Polypedilum*, for which algae are an important part of the diet (Henriques-Oliveira et al., 2003), and therefore they were abundant in pool habitats in the Dajiuhu Wetland. As mouthpart evolution is strongly impacted by feeding

Y. Cao et al.

ecology, morphological traits are thought to be dependent on feeding traits and serve as a good surrogate for habitat changes (Antczak Orlewska et al., 2021). The M14 morphological type are mainly composed of taxa preferring algal-rich conditions (e.g. *P. nubeculosum* type), and are thus abundant in inundated habitats. M15 is a morphological type having narrow or foot-shaped ventromental plates and mainly consisted of collector-gatherers adapted to semi-terrestrial environments such as *Psilometriocnemus*, *Pseudosmittia*, *Limnophyes* and *Pseudorthocladius*, and therefore showed an opposite distribution to the M14 morphological type. For body size, habitats with high-quality food supply are undoubtedly prone to support large individuals to improve their competitiveness and survive in relatively stable conditions like peatland pools here (Iglesias et al., 2012).

Water level is an important environmental filter for biotic communities in both taxonomic and functional trait compositions (Canedo-Argüelles et al., 2016). Chironomid taxonomic compositions showed distinctive variations along the water depth gradient in the current study, in agreement with the relatively limited chironomid work on peatlands (Cao et al., 2019). Semi-terrestrial species (i.e. *Psilometriocnemus*, *Pseudosmittia*, *Neozavrelia*, *Pseudorthocladius* and *Tribelos*) are found in dry moss peats whereas lacustrine taxa (i.e. *Polypedilum*, *Glyptotendipes* and *Chironomus*) live predominantly in inundated habitats like pools. This finding might encourage the usage of a taxonomy based approach of chironomids in tracking the long time-scale history of peatland ontogeny, as peatlands might experience several complex wet-dry transition phases during peat accumulation (Morris et al., 2015). For traits, our results were consistent with other reports in that collectors (especially gatherers) and sprawlers were abundant in intermittent or perennial drought habitats (Canedo-Argüelles et al., 2016; Jovem-Azevedo et al., 2019). Conversely, shredders generally show a lack of drought resistance (Bazzanti et al., 2009), as *Sphagnum* hummocks with continuous drought could be less suitable for shredders to colonize. Drought undoubtedly acted as an ecological filter of body size, as small individuals are commonly found in extreme environments suffering relatively high levels of disturbance (Gomes et al., 2018). Reduced body size allows a high rate of reproduction and growth, as well as a relatively high capacity to surviving disturbances (Bonada et al., 2007).

An orthogonal relationship was identified between K^+ and variables of LOI and DWT (Fig. 12), suggesting that K^+ is independent of the other two parameters, and hence it may be related to the mineral type that is influencing the chironomid communities. Potassium mostly exists in the lattice of aluminosilicate minerals (mainly feldspars and micas) in the

form of ions, but it is generally difficult to be released naturally. The catchment area of the Dajiuhu Wetland is mostly composed of

limestones and dolomites, with a few sandstones and shales in the west (Du et al., 2008). Significant positive relationships were detected between K^+ and Ca^{2+} , as well as Mg^{2+} (Fig. 12), revealing their possible similar sources, indicating that potassium might be released during bedrock decomposition. Potassic fertilizer is increasingly needed in agricultural cultivation in the form of water-soluble salts such as KCl (Skorina and Allanore, 2015). Massive ditches were dug, and natural peatlands were reclaimed in the Dajiuhu basin since the 1980s to pro-

mote local agricultural economic development (Zhou et al., 2017). Agricultural activities within and around the Dajiuhu Wetland might contribute to the potassium supply via running water and/or aeolian transport. The significance of K^+ for chironomid communities could be through a range of related processes. Ionic balance is important for chironomid communities, of which K^+ has a role (Jonusaite et al., 2011) and indirect processes will also be important, such as via substrate shaping through moss growth (Clymo and Hayward, 1982; Rydin and Jeglum, 2013; Cao et al., 2019). Potassium deficiency may trigger hypokalemia for specific taxa and threaten their survival (Belowitz et al., 2014), and it also shapes functional trait communities (Antczak Orlewska et al., 2021). Shredders in our study are positively associated with K^+ , and they were mainly dominated by mining taxa, such as *Endochironomus*, *Glyptochironomus* and *Polypedilum*. These larvae commonly live in lacustrine habitats with coarse organic detritus (Bijl makers, 1983) containing a mixture of rotten plant and animal material which is therefore likely enriched in potassium (Belowitz and O'Donnell, 2013).

The transverse zones between pools and *Sphagnum* moss hummocks in our study supported diverse taxonomic and trait compositions. Edge effects were especially pronounced in ditch habitats since they exhibited the highest abundance, as well as taxonomic and functional diversity of chironomid communities. Natural ecotones possess various habitats and provide more niche opportunities for diverse species assemblages with different functional traits (Kark, 2013), highlighting the vital importance of ecotones in biodiversity maintenance.

4.2. Functional traits are robust in relation to habitat change

Increasing numbers of studies have shown that biotic trait composition is more sensitive to environmental variations than traditional taxonomic structures (Jiang et al., 2019; Paiva et al., 2023), and as such could be a useful bioindicator (Paiva et al., 2023). This aligns with our

study as it showed that in total half of all functional trait categories were sensitive to habitat changes, while only 38 % of traditional taxonomic compositions were. At the temporal scale, five categories of traits showed consistent and synchronous trends. This reveals that each trait could provide an independent functional response to environmental change (Fig. 9), and hence these traits could be used to infer changes in environment and ecosystem functions. It would be more beneficial to comparative analysis across studies (both modern bioassessment and paleoenvironmental studies) than the taxonomy-based approach, in which the influence of biogeographic factors and biological interactions within communities on taxonomic structures (Heino and Tolonen, 2017) should be considered. Moreover, taxonomic compositional changes

could reveal the replacement and gain or loss of species (Mayfield et al., 2022), but traits can reveal changes in ecosystem functions under disturbance, which is conducive to ecosystem management (Haase et al., 2023). Our results indicate that the trait composition of chironomid communities has a robust response to environmental change at spatio-temporal scales.

4.3. Conclusions and implications for bioassessment and paleoenvironmental reconstruction

Our study showed that more chironomid functional trait categories compared to taxonomic assemblages could be used to recognise habitat

Fig. 12. Spearman correlations between different environmental variables. changes at spatial scales. This result was also verified by sedimentary

10

Ecological Indicators 159 (2024) 111656

Y. Cao et al.

data, which record the response of biotic communities to environmental change across temporal scales. Both taxonomic and functional trait communities were primarily controlled by sedimentary organic matter content, depth to water table and K^+ . We therefore conclude that a functional trait-based approach could be a good alternative to a traditional taxonomy-based approach in identifying the response of chironomid assemblages to wetland development. Considering the limitations of taxonomic identification accuracy, especially in peat samples, chironomid functional trait composition is strongly recommended to serve as a surrogate in assessments of environments and ecosystem functions, as well as in palaeo-studies in a range of wetlands.

CRedit authorship contribution statement

Yanmin Cao: Writing – review & editing, Writing – original draft, Investigation, Funding acquisition, Conceptualization. **Peter G. Langdon:** Writing – review & editing. **Shengman Shen:** Writing – review & editing, Methodology. **Hui Li:** Resources, Investigation. **Deng Pan:** Writing – review & editing, Methodology.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We acknowledge Chen Xu, Hou Guilin, Ge Wenlong and Peng Jia for their kind help with their field and laboratory assistance. This work was supported by the National Natural Science Foundation of China (42271165) and the Fundamental Research Funds for the Central Universities, South-Central Minzu University (CZQ23009). Y. Cao was supported by the China Scholarship Council (grant number 202008420139). We thank the very helpful reviewer and editorial

comments in shaping this manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecolind.2024.111656>.

References

- Adeleye, M.A., Haberle, S.G., Gallagher, R., Andrew, S.C., Herbert, A., 2023. Changing plant functional diversity over the last 12,000 years provides perspectives for tracking future changes in vegetation communities. *Nat. Ecol. Evol.* 7, 224–235.
- Antczak-Orlewska, O., Płociennik, M., Sobczyk, R., Okupny, D., Stachowicz-Rybka, R., Rzdokiewicz, M., Siciński, J., Mroczkowska, A., Krapić, M., Słowiński, M., Kittel, P., 2021. Chironomidae Morphological Types and Functional Feeding Groups as a Habitat Complexity Vestige. *8*, 583831.
- Armitage, P.D., Pinder, L., Cranston, P., 2012. *The Chironomidae: biology and ecology of non-biting midges*. Springer Science & Business Media, Dordrecht.
- Bazzanti, M., Della Bella, V., Grezzi, F., 2009. Functional characteristics of macroinvertebrate communities in Mediterranean ponds (Central Italy): Influence of water permanence and mesohabitat type. *Ann. Limnol. - Int. J. Lim.* 45, 29–39.
- Belle, S., Klaus, F., González Sagrario, M.d.l.A., Vrede, T., Goedkoop, W., 2022. Unravelling chironomid biodiversity response to climate change in subarctic lakes across temporal and spatial scales. *Hydrobiologia* 849, 2621–2633.
- Bellwood, D.R., Streitz, R.P., Brandt, S.J., Tebbett, S.B., 2019. The meaning of the term ‘function’ in ecology: A coral reef perspective. *Funct. Ecol.* 33, 948–961.
- Belowitz, R., Leonard, E.M., O'Donnell, M.J., 2014. Effects of exposure to high concentrations of waterborne TI on K and TI concentrations in *Chironomus riparius* larvae. *Comp. Biochem. Physiol. C. Toxicol. Pharmacol.* 166, 59–64.
- Belowitz, R., O'Donnell, M.J., 2013. Ion-selective microelectrode measurements of TI⁺ and K⁺ transport by the gut and associated epithelia in *Chironomus riparius*. *Aquat. Toxicol.* 138–139, 70–80.
- Bijlmakers, L., 1983. *De verspreiding en oecologie van chironomide larven (Chironomidae: Diptera)*. Twee Vennen in De Omgeving Van Oisterwijk (n.br.). K.U. Nijmegen, Versl.
- Bonada, N., Dolédec, S., Statzner, B., 2007. Taxonomic and biological trait differences of stream macroinvertebrate communities between mediterranean and temperate regions: implications for future climatic scenarios. *Glob. Chang. Biol.* 13, 1658–1671.
- Borcard, D., 1992. Partialling out the spatial component of ecological variation. *Ecology* 71, 1045–1055.
- Brooks, S.J., Langdon, P., Heiri, O., 2007. *The identification and use of Palaeartic Chironomidae larvae in palaeoecology*, QRA Technical Guide No. 10. Quaternary Research Association, London.
- Cai, Y., Dong, R., Kattel, G., Zhang, Y., Peng, K., Gong, Z., 2023. Macroinvertebrate diversity and ecosystem functioning across the eutrophication gradients of the middle and lower reaches of Yangtze River lakes (China). *Ecol. Evol.* 13, e9751.
- Canedo-Argüelles, M., Bogan, M.T., Lytle, D.A., Prat, N., 2016. Are Chironomidae (Diptera) good indicators of water scarcity? Dryland streams as a case study. *Ecol.*

- Ind. 71, 155–162.
- Canedo-Argüelles, M., Gutiérrez-Cano, C., Acosta, R., Castro-Lopez, D., Cid, N., Fortuno, P., Munné, A., Múrria, C., Pimentão, A.R., Sarremejane, R., Soria, M., Tarrats, P., Verkaik, I., Prat, N., Bonada, N., 2020. As time goes by: 20 years of changes in the aquatic macroinvertebrate metacommunity of Mediterranean river networks. *J. Biogeogr.* 47, 1861–1874.
- Cao, Y., Langdon, P.G., Yan, Y., Wang, S., Zheng, Z., Zhang, Z., 2019. Chironomid communities from subalpine peatlands in subtropical China as indicators of environmental change. *J. Paleolimnol.* 62, 165–179.
- Clymo, R., Hayward, P., 1982. *The ecology of Sphagnum, Bryophyte ecology*. Springer, Dordrecht.
- Dawson, S.K., Carmona, C.P., González-Suárez, M., Jonsson, M., Chichorro, F., Mallen Cooper, M., Melero, Y., Moor, H., Simaika, J.P., Duthie, A.B., 2021. The traits of “trait ecologists”: An analysis of the use of trait and functional trait terminology. *Ecol. Evol.* 11, 16434–16445.
- Dolédéc, S., Phillips, N., Townsend, C., 2011. Invertebrate community responses to land use at a broad spatial scale: trait and taxonomic measures compared in New Zealand rivers. *Freshwater Biol.* 56, 1670–1688.
- Du, Y., Cai, S.M., Wang, X.L., He, B.Y., Xu, G.L., Jiang, M.X., Xue, H.P., Xiao, F., 2008. Environmental background and ecological restoration of the Dajihu sub-alpine wetland in Mt. Shennongjia. *Resour. Environ. Yangtze Basin* 17 (6), 915–919 in Chinese.
- Dunn, O.J., 1964. Multiple Comparisons Using Rank Sums. *Technometrics* 6, 241–252.
- Entrekín, S.A., Wallace, J.B., Eggert, S.L., 2007. The response of Chironomidae (Diptera) to a long-term exclusion of terrestrial organic matter. *Hydrobiologia* 575, 401–413.
- Epler, J.H. 2001. *Identification manual for the larval Chironomidae (Diptera) of North and South Carolina*. North Carolina Department of Environment and Natural Resources, Division of Water Quality.
- Gautreau, E., Volatier, L., Nogaro, G., Gouze, E., Marmonier, P., Mermillod-Blondin, F., 2023. Interactions between microbial activity and bioturbation modes of benthic invertebrates determine nutrient releases from reservoir sediments. *Freshwater Biol.* 68, 245–259.
- Gomes, W.I.A., Jovem-Azevêdo, D.d.S., Paiva, F.F., Milesi, S.V., Molozzi, J., 2018. Functional attributes of Chironomidae for detecting anthropogenic impacts on reservoirs: A biomonitoring approach. *Ecol. Ind.* 93, 404–410.
- Haase, P., Bowler, D.E., Baker, N.J., et al., 2023. The recovery of European freshwater biodiversity has come to a halt. *Nature* 620, 582–588.
- Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: paleontological statistics software package for education and data analysis. *Palaeontol. Electron.* 4, 9.
- Heino, J., 2008. Patterns of functional biodiversity and function-environment relationships in lake littoral macroinvertebrates. *Limnol. Oceanogr.* 53, 1446–1455.
- Tolonen, K.T., 2017. Ecological drivers of multiple facets of beta diversity in a lentic macroinvertebrate metacommunity. *Limnol. Oceanogr.* 62, 2431–2444.
- Henriques-Oliveira, A.L., Nessimian, J.L., Dorville, L.F.M., 2003. Feeding habits of Chironomid larvae (Insecta: Diptera) from a stream in the Floresta da Tijuca, Rio de Janeiro, Brazil. *Braz. J. Biol.* 63, 269–281.
- Hou, G.L., Cao, Y.M., Tian, W.L., Deng, P., Shen, S.M., Chen, X., 2022. Response of chironomid assemblages to the changes of wetness in Dajihu Peatland of Hubei Province during the past 400 years. *J. Lake Sci.* 34, 1030–1042 in Chinese.
- Hu, S., Niu, Z., Chen, Y., Li, L., Zhang, H., 2017. Global wetlands: Potential distribution, wetland loss, and status. *Sci. Total Environ.* 586, 319–327.
- Iglesias, S., Tracy, C., Bedford, G., Christian, K., 2012. Habitat differences in body size and shape of the Australian agamid lizard, *Lophognathus temporalis*. *J. Herpetol.* 46, 297–303.
- Jiang, X., Pan, B., Song, Z., Xie, Z., 2019. Do functional traits of chironomid assemblages respond more readily to eutrophication than taxonomic composition in Chinese floodplain lakes? *Ecol. Ind.* 103, 355–362.
- Jonusaite, S., Kelly, S.P., Donini, A., 2011. The physiological response of larval *Chironomus riparius* (Meigen) to abrupt brackish water exposure. *J. Comp. Physiol. B* 181, 343–352.
- Jovem-Azevêdo, D., Bezerra-Neto, J.F., Azevêdo, E.L., Gomes, W.I.A., Molozzi, J., Feio, M.J., 2019. Dipteran assemblages as functional indicators of extreme droughts. *J. Arid Environ.* 164, 12–22.
- Kark, S., 2013. Effects of ecotones on biodiversity. *Encyclopedia of Biodiversity* 142, 1.
- Kivilä, E.H., Luoto, T.P., Rantala, M.V., Kiljunen, M., Rautio, M., Nevalainen, L., 2019. Environmental controls on benthic food web functions and carbon resource use in subarctic lakes. *Freshwater Biol.* 64, 643–658.
- Labrière, E., Legendre, P., Shipley, B., Labrière, M.E., 2014. Measuring functional diversity from multiple traits, and other tools for functional ecology. *R Package FD*.
- 11
- Ecological Indicators 159 (2024) 111656*
- Martins, I., Castro, D.M.P., Macedo, D.R., Hughes, R.M., Callisto, M., 2021. Anthropogenic impacts influence the functional traits of Chironomidae (Diptera) assemblages in a neotropical savanna river basin. *Aquat. Ecol.* 55, 1081–1095.
- Mason, N.W.H., Mouillot, D., Lee, W.G., Wilson, J.B., 2005. Functional richness, functional evenness and functional divergence: the primary components of functional diversity. *Oikos* 111, 112–118.
- Mayfield, R.J., Dearing, J.A., Doncaster, C.P., Langdon, P.G., 2022. Stability of chironomid community structure during historic climatic and environmental change in subarctic Alaska. *Limnol. Oceanogr.* 67, S444–S460.
- Mitchell, E.A.D., Buttler, A., Grosvernier, P., Rydin, H., Albinsson, C., Greenup, A.L., Heijmans, M.M.P.D., Hoosbeek, M.R., Saarinen, T., 2000. Relationships among testate amoebae (Protozoa), vegetation and water chemistry in five Sphagnum dominated peatlands in Europe. *New Phytol.* 145, 95–106.
- Morris, P.J., Baird, A.J., Young, D.M., Swindles, G.T., 2015. Untangling climate signals from autogenic changes in long-term peatland development. *Geophys. Res. Lett.* 42, 10,788–710,797.
- Morse, J.C., Bae, Y.J., Munkhjargal, G., Sangpradub, N., Tanida, K., Vshivkova, T.S., Wang, B., Yang, L., Yule, C.M., 2007. Freshwater biomonitoring with macroinvertebrates in East Asia. *Front. Ecol. Environ.* 5, 33–42.
- Nevalainen, L., Luoto, T.P., Manca, M., Weisse, T., 2015. A paleolimnological perspective on aquatic biodiversity in Austrian mountain lakes. *Aquat. Sci.* 77, 59–69.
- Nyman, M., Korhola, A., Brooks, S.J., 2005. The distribution and diversity of Chironomidae (Insecta: Diptera) in western Finnish Lapland, with special emphasis on shallow lakes. *Glob. Ecol. Biogeogr.* 14, 137–153.
- Ogle, D., Wheeler, P., Dinno, A., 2020. FSA: fisheries stock analysis. R package version. Chapman & Hall/CRC, Boca Raton, p. 30.
- Paiva, F.F., Melo, D.B.D., Dolbeth, M., Molozzi, J., 2023. Functional threshold responses of benthic macroinvertebrates to environmental stressors in reservoirs. *J. Environ. Manage.* 329, 116970.
- Peng, J., Zeng, L., Huang, X., Chen, X., 2023. Climate, atmospheric deposition and catchment process interact to trigger recent diatom community reorganization in
- Y. Cao et al.
- Larentis, C., Pavanelli, C.S., Delariva, R.L., 2022. Do environmental conditions modulated by land use drive fish functional diversity in streams? *Hydrobiologia* 849, 4465–4483.
- Larocque, I., Pienitz, R., Rolland, N., 2006. Factors influencing the distribution of chironomids in lakes distributed along a latitudinal gradient in northwestern Quebec, Canada. *Can. J. Fish. Aquat. Sci.* 63, 1286–1297.
- Legendre, P., Anderson, M.J., 1999. Distance-based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecol. Monogr.* 69, 1–24.
- Legendre, P., Legendre, L., 1998. *Numerical ecology*, 2nd ed. Elsevier, Amsterdam.
- Leifeld, J., Menichetti, L., 2018. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nat. Commun.* 9, 1071.
- Leszczynska, J., Grzybkowska, M., Głowacki, Ł., Dukowska, M., 2019. Environmental Variables Influencing Chironomid Assemblages (Diptera: Chironomidae) in Lowland Rivers of Central Poland. *Environ. Entomol.* 48, 988–997.
- Li, Z., Wang, J., Liu, Z., Meng, X., Heino, J., Jiang, X., Xiong, X., Jiang, X., Xie, Z., 2019. Different responses of taxonomic and functional structures of stream macroinvertebrate communities to local stressors and regional factors in a subtropical biodiversity hotspot. *Sci. Total Environ.* 655, 1288–1300.
- Li, J., Zheng, Z., Huang, K., Yang, S., Chase, B., Valsecchi, V., Carré, M., Cheddadi, R., 2013. Vegetation changes during the past 40,000 years in Central China from a long fossil record. *Quat. Int.* 310, 221–226.
- Luoto, T.P., Ojala, A.E.K., 2018. Controls of climate, catchment erosion and biological production on long-term community and functional changes of chironomids in High Arctic lakes (Svalbard). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 505, 63–72.
- Luoto, T.P., Rantala, M.V., Galkin, A., Rautio, M., Nevalainen, L., 2016. Environmental determinants of chironomid communities in remote northern lakes across the treeline – Implications for climate change assessments. *Ecol. Ind.* 61, 991–999.
- Mandaville, S., 2002. Benthic macroinvertebrates in freshwaters: Taxa tolerance values, metrics, and protocols. Soil & Water Conservation Society of Metro Halifax, Nova Scotia.

- alpine lakes of the eastern monsoonal region of China. *Quaternary Sci Rev* 310, 108131.
- Piano, E., Doretto, A., Mammola, S., Falasco, E., Fenoglio, S., Bona, F., 2020. Taxonomic and functional homogenisation of macroinvertebrate communities in recently intermittent Alpine watercourses. *Freshwater Biol* 65, 2096–2107.
- Premke, K., Karlsson, J., Steger, K., Gudas, C., von Wachenfeldt, E., Tranvik, L.J., 2010. Stable isotope analysis of benthic fauna and their food sources in boreal lakes. *J. N. Am. Benthol. Soc.* 29, 1339–1348.
- Qin, Y., Li, H., Mazei, Y., Kurina, I., Swindles, G.T., Bobrov, A., Tsyganov, A.N., Gu, Y., Huang, X., Xue, J., Lamentowicz, M., Marcisz, K., Roland, T., Payne, R.J., Mitchell, E. A.D., Xie, S., 2021. Developing a continental-scale testate amoeba hydrological transfer function for Asian peatlands. *Quaternary Sci Rev* 258, 106868.
- Radkova, V., Syrovatka, V., Bojkova, J., Schenkova, J., Kroupalova, V., Horsak, M., 2014. The importance of species replacement and richness differences in small-scale diversity patterns of aquatic macroinvertebrates in spring fens. *Limnologia* 47, 52–61.
- Rydin, H., Jeglum, J.K., 2013. *The biology of peatlands*, 2e. Oxford University Press, Oxford.
- Saulino, H.H., Leite-Rossi, L.A., Trivinho-Strixino, S., 2017. The effect of small reservoirs on chironomid diversity and trait composition in Savanna streams: evidence for Serial Discontinuity Concept. *Hydrobiologia* 793, 109–119.
- Schleuter, D., Daufresne, M., Massol, F., Argillier, C., 2010. A user's guide to functional diversity indices. *Ecol. Monogr.* 80, 469–484.
- Serra, S.R.Q., Cobo, F., Graça, M.A.S., Dolédec, S., Feio, M.J., 2016. Synthesising the trait information of European Chironomidae (Insecta: Diptera): Towards a new database. *Ecol. Ind.* 61, 282–292.
- Skorina, T., Allanore, A., 2015. Aqueous alteration of potassium-bearing aluminosilicate minerals: from mechanism to processing. *Green Chem.* 17, 2123–2136.
- Smilauer, P., Lepš, J., 2014. *Multivariate Analysis of Ecological Data Using CANOCO 5*. Cambridge University Press, Cambridge.
- Syrovatka, V., Brabec, K., 2006. Effects of physical factors on chironomid larvae distribution and community structure at a mesohabitat scale. *Internationale Vereinigung Für Theoretische Und Angewandte Limnologie: Verhandlungen* 29 (4), 1845–1848.
- Tang, H.Q., 2006. *Biosystematic study on the chironomid larvae in China (Diptera: Chironomidae)*. Nankai University, Tianjin. Ph. D thesis (in Chinese).
- Van Duinen, G., Vermonden, K., Brock, A., Leuven, R., Smolders, A., Van Der Velde, G., Verberk, W., Esselink, H., 2006. Basal food sources for the invertebrate food web in nutrient poor and nutrient enriched raised bog pools. *Proc. Section Exp. Appl. Entomol.-Netherlands Entomological Society* 17, 37.
- Villéger, S., Mason, N.W.H., Mouillot, D., 2008. New multidimensional functional diversity indices for a multifaceted framework in functional ecology. *Ecology* 89, 2290–2301.
- Wang, J.C., Wang, X.H., 2011. *Chironomidae larval in North China*. China Yanshi Press, Beijing.
- Wentzky, V.C., Tittel, J., Jäger, C.G., Bruggeman, J., Rinke, K., 2020. Seasonal succession of functional traits in phytoplankton communities and their interaction with trophic state. *J. Ecol.* 108, 1649–1663.
- Xi, Y., Peng, S., Ciais, P., Chen, Y., 2021. Future impacts of climate change on inland Ramsar wetlands. *Nat. Clim. Chang.* 11, 45–51.
- Zebal, Y.D., Righi, B.D.P., Anni, I.S.A., Escarrone, A.L.V., Roza, M., Vieira, C.E.D., Costa, P.G., Bianchini, A., 2021. Pollution levels and biomarker responses in zooplankton from three hydrographic regions of southern Brazil: An integrated approach for water quality monitoring. *J. Environ. Chem. Eng.* 9, 106180.
- Zhang, Z.Z., 1982. The formation and distribution of bog in Shennongjia. *J. Northeast Normal University (nat. Sci.)* 1, 95e103 in Chinese.
- Zhou, W.C., Shen, Y.H., Cui, H.X., Zhang, Z.Q., Yang, J.Y., 2017. The countermeasures for protection and management of Dajiu Lake Wetland in Shennongjia. *Wetland Sci. Manage.* 13 (2), 34–37 in Chinese.