

Ecosystem functions and functional traits for the study of phytophagous scarab beetles (Coleoptera: Scarabaeidae)

Supporting Information

Table S1. Traits of phytophagous Scarabaeidae species selected from the entomological collection of the University of the Atlantic, UARC. Abbreviations: TL: Total length, PL: pronotum length, PW: pronotum width, ID: interocular distance, ED: eye diameter, EL: elytral length, EW: elytral width, TT: total thickness, ET: elytral thickness, LeBoRat: leg length to body length ratio, MCL: Maximum Metacoxa Length, BIO: biomass, WA: wing area, WW: wing width, WH: wing height, WL: wing loading, TrGr: trophic group.

Species	TL	PL	PW	ID	ED	EL	EW	TT	ET	LeBoRat	MCL	BIO	WA	WW	WH	WL	TrGr
<i>Amithao lafertei</i>	24.5	7.7	10.2	3	2.1	16.8	11.7	6.8	3.2	0.8	3.6	0.1	170.5	27.1	10	2.7	4
<i>Aspidolea singularis</i>	14	3.8	6.1	2.3	1.6	10.2	3.9	6	2.5	0.8	1.7	0.1	112.8	21.1	7	3	1
<i>Astaena</i> sp	9.9	2.2	3.9	1.5	1.1	7.7	3.9	4.7	2.1	1	1.7	0	70.2	15.2	5.5	2.7	1
<i>Barybas</i> sp	6.2	1.8	3.1	1.1	0.8	4.4	1.9	2.8	1.2	0.9	0.7	0	27.4	10.5	3.6	2.9	1
<i>Ceraspis innotata</i>	9.9	3.8	4.9	1.8	1	8.3	6	4.1	2	1	1.3	0	72.2	17.1	5.7	3	1
<i>Ceraspis quadripustulata</i>	7.5	2.9	3.9	1.6	0.8	6.8	4.6	4.1	1.8	1.3	0.9	0	44.8	13.6	4.3	3.1	1
<i>Cyclocephala amazona</i>	11.9	3.6	5.4	2	1.5	8.3	3.4	4.7	2.1	0.8	1.5	0	68.4	16.5	5.2	3.2	2
<i>Cyclocephala fulgurata</i>	13.3	3.7	6	2	1.4	9.7	4.8	5	2.7	0.8	1.6	0.1	99.5	19.3	6.6	2.9	2
<i>Cyclocephala lunulata</i>	13	3.8	5.8	1.9	1.5	9.1	4.8	4.9	2.2	0.7	1.5	0.1	83.6	17.7	6	2.9	2
<i>Cyclocephala maffufa</i>	23.4	7	10.3	3.9	2.5	16.3	8.1	8.6	3.9	0.7	2.3	0.3	245.2	29.5	10.6	2.8	2
<i>Cyclocephala ovulum</i>	10.7	3.1	5	1.6	1.2	7.6	3.7	4.1	2	0.8	1.2	0	70.8	15.9	5.7	2.8	2
<i>Cyclocephala prelli</i>	17	4.9	7.6	2.9	2	12.2	6	5.8	2.9	0.8	2.1	0.1	123.7	21.5	7.3	3	2
<i>Golofa</i> sp	30.8	9.8	12.8	2.9	2.7	21	11.1	12.7	7.1	0.7	3.1	0.9	425.1	39	13.6	2.9	3
<i>Hoplopyga liturata</i>	15.6	5.2	7.5	1.7	1.3	10.5	8.7	5	3.7	0.8	2.4	0	63	15.6	6	2.6	4
<i>Isonychus</i> sp	8.9	2.6	3.6	1.4	0.8	6.3	2.6	4.7	1.6	1.1	1.6	0	47.7	13.9	4.8	2.9	1
<i>Leucothyreus</i> sp1	9.2	2.3	4.5	2.2	1.2	6.8	2.9	4.2	1.7	0.9	1.1	0	74.2	16.6	5.8	2.8	1
<i>Leucothyreus</i> sp2	12	2.8	5.1	2.7	1.5	9.2	4.4	5	2.1	0.8	1	0	94.1	18.9	6.3	3	1
<i>Paranomala inconstans</i>	14.5	3.5	6.2	2	1.4	11	4.5	7.6	3.2	0.9	2.4	0.1	132.5	21.7	7.6	2.8	1
<i>Paranomala</i> sp3	9.5	2.5	4	1.4	0.9	7	2.8	4.3	1.7	0.9	1.7	0	55.7	13.9	5.1	2.8	1
<i>Pelidnota glaberrima</i>	18.5	5	7.8	2.9	1.9	13.5	6.9	7.9	3.1	0.6	2.1	0.2	190.3	27.6	8.8	3.1	1
<i>Pelidnota polita</i>	23.2	5.4	11	4.3	2.5	17.8	9.3	11.1	5.1	0.7	2.2	0.5	328.2	35.5	11.8	3	1
<i>Phileurus dydimus</i>	34.9	11	13.7	5	3	23.9	24.7	13.5	5.2	0.9	4	0.5	550.5	41	11	3.7	5
<i>Pholeurus valgus</i>	20.6	7	9.4	3.3	2	13.6	10	6	4	1	2	0.1	130.5	23.9	10	2.4	5
<i>Phyllophaga aguadita</i>	19.1	4.1	7.7	2.9	2.2	15	10.5	9	5.2	1	2.1	0.1	196.6	27.5	9.6	2.9	1

Species	TL	PL	PW	ID	ED	EL	EW	TT	ET	LeBoRat	MCL	BIO	WA	WW	WH	WL	TrGr
<i>Phyllophaga brevisetosa</i>	16.2	3.8	6.5	2.4	1.5	12.4	9.2	8.3	4.6	1.1	1.8	0.1	147	23	7.9	2.9	1
<i>Phyllophaga logicornis</i>	22	4.5	8.4	2.9	2.1	17.6	8.1	10.1	4.8	0.8	2.5	0.2	259.1	30.7	10.9	2.8	1
<i>Phyllophaga obsoleta</i>	16.3	3.6	6.6	2.4	1.8	12.7	8.5	7.7	3.8	1.1	1.9	0.1	142.5	23.2	8.2	2.8	1
<i>Phyllophaga pruinipennis</i>	19.5	4.2	7.9	3	2.2	15.3	10.9	8.8	4.8	1.1	2.2	0.1	203.5	27.6	9.9	2.8	1
<i>Phyllophaga</i> sp	15.4	3.6	5.8	2.1	1.4	11.8	7.3	6.7	3.9	0.9	1.9	0.1	113.2	21.2	7	3	1
<i>Plectris</i> sp	13.4	3.5	5.1	2	1.3	9.9	3.7	6.5	2.9	1	2	0.1	120.2	21.2	7.2	2.9	1
<i>Stenocrates laevicollis</i>	17.1	4.6	8.1	2.7	1.8	12.6	6.5	7.4	3.1	0.8	2	0.3	146.1	23.4	8.3	2.8	1
<i>Tomarus fossor</i>	13.1	4.4	6.6	2	1.3	8.6	4.1	5.7	3.5	0.7	2	0.1	82.7	18.4	6	3	3