

Supplementary Materials: Use of Carabids for the Post-Market Environmental Monitoring of Genetically Modified Crops

Oxana Skoková Habušťová, Zdeňka Svobodová, Ľudovít Cagán and František Sehnal

Table S1. The sample dates of deployment of pitfall traps at the locality western Slovakia in 2014 and 2015.

2014	2015	Stage of maize
May 15	May 21	BBCH 09
June 10	June 04	BBCH 11
June 24	June 18	BBCH 13
July 8	July 01	BBCH 17
July 22	July 16	BBCH 34
August 5	July 30	BBCH 53
August 19	August 12	BBCH 63
September 3	August 27	BBCH 69
September 16	September 10	BBCH 79
October 1	September 24	BBCH 89

Table S2. The abundance of carabids in the examined localities in 2002–2015.

Tribe	Species	Localities					Total
		South Bohemia 1	South Bohemia 2	South Bohemia 3	Central Bohemia	western Slovakia	
Pterostichini	<i>Abax parallelus</i> (Duftschmid, 1812)	0	0	0	1	0	1
Harpalini	<i>Acupalpus meridianus</i> (Linnaeus, 1761)	0	0	0	1	1	2
Platynini	<i>Agonum gracilipes</i> (Duftschmid, 1812)	0	11	3	0	0	14
Platynini	<i>Agonum muelleri</i> (Herbst, 1784)	42	211	3	0	0	256
Platinini	<i>Agonum sexpunctatum</i> (Linnaeus, 1758)	1	5	3	0	0	9
Platynini	<i>Agonum versutum</i> (Sturm, 1824)	0	0	1	0	0	1
Zabrini	<i>Amara aenea</i> (De Geer, 1774)	0	75	6	1	1	83
Zabrini	<i>Amara apricaria</i> (Paykull, 1790)	0	6	0	0	1	7
Zabrini	<i>Amara aulica</i> (Panzer, 1796)	2	2	3	0	0	7
Zabrini	<i>Amara consularis</i> (Duftschmid 1812)	3	1	0	139	0	143
Zabrini	<i>Amara cursitans</i> (Zimmermann, 1831)	1	5	0	0	0	6
Zabrini	<i>Amara eurynota</i> (Panzer, 1797)	1	55	0	0	0	56
Zabrini	<i>Amara familiaris</i> (Duftschmid 1812)	0	8	0	0	0	8
Zabrini	<i>Amara littorea</i> (C.G. Thomson, 1857)	1	6	0	0	0	7
Zabrini	<i>Amara nitida</i> (Sturm, 1825)	0	1	0	0	0	1
Zabrini	<i>Amara ovata</i> (Fabricius, 1792)	3	8	0	6	0	17
Zabrini	<i>Amara similata</i> (Gyllenhal, 1810)	0	59	0	0	0	59

Zabrini	<i>Amara tibialis</i> (Paykull, 1798)	0	2	0	0	0	2
Platynini	<i>Anchomenus dorsalis</i> (Pontoppidan, 1763)	3	3	10	12	1071	1099
Harpalini	<i>Anisodactylus binotatus</i> (Fabricius, 1787)	0	0	2	0	3	5
Harpalini	<i>Anisodactylus signatus</i> (Panzer, 1797)	0	3	1	2	1	7
Licinini	<i>Badister bullatus</i> (Schränk, 1798)	0	0	1	3	0	4
Licinini	<i>Badister lacertosus</i> (Sturm, 1815)	0	2	0	0	0	2
Licinini	<i>Badister unipustulatus</i> (Bonelli, 1813)	0	1	0	0	0	1
Bembidiini	<i>Bembidion biguttatum</i> (Fabricius, 1779)	0	3	0	0	0	3
Bembidiini	<i>Bembidion femoratum</i> (Sturm, 1825)	20	1	2	0	0	23
Bembidiini	<i>Bembidion guttula</i> (Fabricius, 1792)	0	1	0	0	0	1
Bembidiini	<i>Bembidion illigeri</i> (Netolitzky, 1914)	0	2	0	0	0	2
Bembidiini	<i>Bembidion lampros</i> (Herbst, 1784)	218	163	18	63	0	462
Bembidiini	<i>Bembidion obtusum</i> (Audinet-Serville, 1821)	0	0	0	45	1	46
Bembidiini	<i>Bembidion properans</i> (Stephens, 1828)	34	34	2	41	3	114

Table S2. Cont.

Tribe	Species	Localities					Total
		South Bohemia 1	South Bohemia 2	South Bohemia 3	Central Bohemia	western Slovakia	
Bembidiini	<i>Bembidion quadrimaculatum</i> (Linnaeus, 1761)	813	728	80	59	0	1680
Brachinini	<i>Brachinus crepitans</i> (Linnaeus, 1758)	0	0	0	13	335	348
Brachinini	<i>Brachinus ganglbaueri</i> (Schauberger, 1921)	0	0	0	0	294	294

Brachinini	<i>Brachinus explodens</i> (Duftschmid, 1812)	0	0	0	0	71	71
Brachinini	<i>Brachinus psophia</i> (Audinet-Serville 1821)	0	0	0	0	31	31
Platynini	<i>Calathus ambiguus</i> (Paykull, 1790)	13	3	0	1	14	31
Platynini	<i>Calathus erratus</i> (C. R. Sahlberg, 1827)	0	6	0	2	0	8
Platynini	<i>Calathus fuscipes</i> (Goeze, 1777)	814	566	473	804	154	2811
Platynini	<i>Calathus melanocephalus</i> (Linnaeus, 1758)	71	7	13	0	0	91
Carabini	<i>Calosoma auropunctatum</i> (Herbst, 1784)	0	0	0	0	2	2
Carabini	<i>Carabus granulatus</i> (Linnaeus, 1758)	2	493	101	0	0	596
Carabini	<i>Carabus hortensis</i> (Linnaeus, 1758)	0	2	0	0	0	2
Carabini	<i>Carabus scheidleri</i> (Panzer, 1799)	36	1	0	0	0	37
Carabini	<i>Carabus violaceus</i> (Linnaeus, 1758)	0	11	2	0	1	14
Callistini	<i>Chlaenius nitidulus</i> (Schrank, 1781)	0	2	0	0	0	2
Clivinini	<i>Clivina fossor</i> (Linnaeus, 1758)	161	135	29	0	0	325
Platynini	<i>Dolichus halensis</i> (Schaller, 1783)	0	0	0	42	52	94
Harpalini	<i>Harpalus affinis</i> (Schrank, 1781)	97	484	117	60	162	920
Harpalini	<i>Harpalus caspius roubali</i> (Schauberger, 1928)	0	0	0	0	1	1
Harpalini	<i>Harpalus distinguendus</i> (Duftschmid, 1812)	0	20	0	124	60	204
Harpalini	<i>Harpalus honestus</i> (Duftschmid, 1812)	0	0	0	3	0	3
Harpalini	<i>Harpalus rubripes</i> (Dufschmid, 1812)	0	8	2	2724	0	2734
Harpalini	<i>Harpalus signaticornis</i> (Duftschmid in 1812)	0	1	0	1	0	2
Harpalini	<i>Harpalus smaragdinus</i> (Duftschmid in 1812)	1	1	0	0	0	2

Harpalini	<i>Harpalus tardus</i> (Panzer, 1797)	0	0	0	0	1	1
Platynini	<i>Laemostenus terricola</i> (Herbst, 1784)	0	0	0	0	38	38
Trechini	<i>Lasiotrechus discus</i> (Fabricius, 1792)	1	0	0	0	0	1
Nebrini	<i>Leistus ferrugineus</i> (Linnaeus, 1758)	0	0	0	0	15	15
Loricerini	<i>Loricera pilicornis</i> (Fabricius, 1775)	37	2	11	1	0	51
Lebiini	<i>Microlestes maurus</i> (Sturm, 1827)	4	16	1	12	4	37
Lebiini	<i>Microlestes minutulus</i> (Goeze 1777)	3	8	0	0	0	11

Table S2. Cont.

Tribe	Species	Localities					Total
		South Bohemia 1	South Bohemia 2	South Bohemia 3	Central Bohemia	western Slovakia	
Nebtriini	<i>Nebria brevicollis</i> (Fabricius, 1792)	19	0	1	0	0	20
Notiophilini	<i>Notiophilus aquaticus</i> (Linnaeus, 1758)	0	1	1	0	0	2
Notiophilini	<i>Notiophilus palustris</i> (Duftschmid, 1812)	3	0	0	59	1	63
Notiophilini	<i>Notiophilus pusillus</i> (G.R.Waterhouse, 1833)	0	0	0	13	0	13
Harpalini	<i>Ophonus azureus</i> (Fabricius, 1775)	0	2	1	0	0	3
Harpalini	<i>Ophonus nitidulus</i> (Stephens, 1828)	0	0	0	1	0	1
Platynini	<i>Platynus assimilis</i> (Paykull, 1790)	0	2	1	0	0	3
Pterostichini	<i>Poecilus cupreus</i> (Linnaeus, 1758)	874	13 234	1467	250	150	15 975
Pterostichini	<i>Poecilus lepidus</i> (Leske 1785)	0	0	0	0	63	63
Pterostichini	<i>Poecilus sericeus</i> (Fischer von Waldheim, 1824)	0	0	0	0	77	77

Pterostichini	<i>Poecilus versicolor</i> (Sturm, 1824)	1	1499	207	3	0	1710
Pterostichini	<i>Pseudoophonus griseus</i> (Panzer, 1796)	0	17	0	5	0	22
Pterostichini	<i>Pseudoophonus rufipes</i> (De Geer)	239	932	144	0	6556	7871
Pterostichini	<i>Pterostichus melanarius</i> (Illiger, 1798)	1740	2978	12 506	995	78	18 297
Pterostichini	<i>Pterostichus melas</i> (Creutzer, 1799)	0	0	0	0	1	1
Pterostichini	<i>Pterostichus niger</i> (Schaller, 1783)	2	87	21	0	1	111
Pterostichini	<i>Pterostichus nigrata</i> (Paykull, 1790)	0	52	11	0	0	63
Pterostichini	<i>Pterostichus ovoideus</i> (Sturm, 1824)	0	2	0	0	0	2
Pterostichini	<i>Pterostichus strenuus</i> (Panzer, 1796)	13	10	0	0	0	23
Pterostichini	<i>Pterostichus vernalis</i> (Panzer, 1795)	7	3	0	1	0	11
Pterostichini	<i>Stomis pumicatus</i> (Panzer, 1795)	0	0	0	4	0	4
Trechini	<i>Trechus quadristriatus</i> (Schränk, 1781)	204	33	329	160	115	841
Harpalini	<i>Trichotichnus laevicollis</i> (Duftschmid, 1812)	0	1	0	0	0	1
Zabrinini	<i>Zabrus tenebrioides</i> (Goeze, 1777)	0	0	0	180	42	222

Table S3. Incidence and functional traits of carabids¹ captured in localities South Bohemia 1, 2 and 3, Central Bohemia and western Slovakia in 2002–2015.

Tribe	Species	Incidence	Body size	Habitat affinity	Humidity affinity	Breeding period	Food specialization
Pterostichini	<i>Abax parallelus</i> (Duftschmid, 1812)	Least concern	B	Silvicolous	Hygrophilous	Spring	Carnivorous
Harpalini	<i>Acupalpus meridianus</i> (Linnaeus, 1761)	Least concern	D	Open biotopes	Mesophilous	Spring	Omnivorous
Platynini	<i>Agonum gracilipes</i> (Duftschmid, 1812)	Least concern	C	Silvicolous	Eurytopic	Spring	Carnivorous
Platynini	<i>Agonum muelleri</i> (Herbst, 1784)	Least concern	C	Eurytopic	Hygrophilous	Spring	Carnivorous
Platinini	<i>Agonum sexpunctatum</i> (Linnaeus, 1758)	Least concern	C	Eurytopic	Hygrophilous	Spring	Carnivorous
Platynini	<i>Agonum versutum</i> (Sturm, 1824)	Least concern	C	Silvicolous	Hygrophilous	Spring	Carnivorous
Zabrini	<i>Amara aenea</i> (De Geer, 1774)	Least concern	C	Open biotopes	Xerophilous	Spring	Omnivorous
Zabrini	<i>Amara apricaria</i> (Paykull, 1790)	Least concern	C	Open biotopes	Xerophilous	Autumn	Omnivorous
Zabrini	<i>Amara aulica</i> (Panzer, 1796)	Least concern	B	Open biotopes	Mesophilous	Autumn	Omnivorous
Zabrini	<i>Amara consularis</i> (Duftschmid 1812)	Least concern	C	Eurytopic	Eurytopic	Autumn	Omnivorous
Zabrini	<i>Amara cursitans</i> (Zimmermann, 1831)	Least concern	C	Eurytopic	Eurytopic	Autumn	Omnivorous

Zabrini	<i>Amara eurynota</i> (Panzer, 1797)	Least concern	C	Open biotopes	Mesophilous	Spring	Omnivorous
Zabrini	<i>Amara familiaris</i> (Duftschmid 1812)	Least concern	C	Open biotopes	Mesophilous	Spring	Omnivorous
Zabrini	<i>Amara littorea</i> (C.G. Thomson, 1857)	Least concern	C	Open biotopes	Xerophilous	Spring	Omnivorous
Zabrini	<i>Amara nitida</i> (Sturm, 1825)	Least concern	C	Open biotopes	Xerophilous	Spring	Omnivorous
Zabrini	<i>Amara ovata</i> (Fabricius, 1792)	Least concern	C	Silvicolous	Hygrophilous	Spring	Omnivorous

Table S3. Cont.

Tribe	Species	Incidence	Body size	Habitat affinity	Humidity affinity	Breeding period	Food specialization
Zabrini	<i>Amara similata</i> (Gyllenhal, 1810)	Least concern	C	Open biotopes	Mesophilous	Spring	Omnivorous
Zabrini	<i>Amara tibialis</i> (Paykull, 1798)	Least concern	D	Open biotopes	Mesophilous	Spring/summer	Omnivorous
Platynini	<i>Anchomenus dorsalis</i> (Pontoppidan, 1763)	Least concern	C	Open biotopes	Hygrophilous	Spring	Carnivorous
Harpalini	<i>Anisodactylus binotatus</i> (Fabricius, 1787)	Least concern	C	Open biotopes	Mesophilous	Spring	Omnivorous
Harpalini	<i>Anisodactylus signatus</i> (Panzer, 1797)	Least concern	B	Open biotopes	Hygrophilous	Spring	Omnivorous
Licinini	<i>Badister bullatus</i>	Least concern	D	Eurytopic	Hygrophilous	Spring	Carnivorous

(Schränk, 1798)							
Licinini	<i>Badister lacertosus</i> (Sturm, 1815)	Least concern	C	Silvicolous	Hygrophilous	Spring	Carnivorous
Licinini	<i>Badister unipustulatus</i> (Bonelli, 1813)	Least concern	C	Silvicolous	Hygrophilous	Spring	Carnivorous
Bembidiini	<i>Bembidion biguttatum</i> (Fabricius, 1779)	Least concern	D	Silvicolous	Hygrophilous	Spring	Carnivorous
Bembidiini	<i>Bembidion femoratum</i> (Sturm, 1825)	Least concern	D	Eurytopic	Mesophilous	Spring	Carnivorous
Bembidiini	<i>Bembidion guttula</i> (Fabricius, 1792)	Least concern	D	Eurytopic	Hygrophilous	Spring	Carnivorous
Bembidiini	<i>Bembidion illigeri</i> (Netolitzky, 1914)	Least concern	D	Eurytopic	Hygrophilous	Spring	Carnivorous
Bembidiini	<i>Bembidion lampros</i> (Herbst, 1784)	Least concern	D	Open biotopes	Eurytopic	Spring	Carnivorous
Bembidiini	<i>Bembidion obtusum</i> (Audinet-Serville, 1821)	Least concern	D	Eurytopic	Eurytopic	Autumn/early spring	Carnivorous
Bembidiini	<i>Bembidion properans</i> (Stephens, 1828)	Least concern	D	Open biotopes	Hygrophilous	Spring	Carnivorous
Bembidiini	<i>Bembidion quadrimaculatum</i> (Linnaeus, 1761)	Least concern	D	Open biotopes	Eurytopic	Spring	Carnivorous

Table S3. *Cont.*

Tribe	Species	Incidence	Body size	Habitat affinity	Humidity affinity	Breeding period	Food specialization
-------	---------	-----------	-----------	------------------	-------------------	-----------------	------------------------

Brachinini	<i>Brachinus crepitans</i> (Linnaeus, 1758)	Least concern	C	Open biotopes	Xerophilous	Summer	Carnivorous
Brachinini	<i>Brachinus ganglbaueri</i> (Schauberger, 1921)	Least concern	C	Open biotopes	Hygrophilous	Spring	Carnivorous
Brachinini	<i>Brachinus explodens</i> (Duftschmid, 1812)	Least concern	D	Open biotopes	Xerophilous	Spring	Carnivorous
Brachinini	<i>Brachinus psophia</i> (Audinet-Serville 1821)	CZ: endangered SK: least concern	C	Open biotopes	Hygrophilous	Spring	Carnivorous
Platynini	<i>Calathus ambiguus</i> (Paykull, 1790)	Least concern	C	Open biotopes	Xerophilous	Autumn	Carnivorous
Platynini	<i>Calathus erratus</i> (C. R. Sahlberg, 1827)	Least concern	C	Open biotopes	Xerophilous	Autumn	Carnivorous
Platynini	<i>Calathus fuscipes</i> (Goeze, 1777)	Least concern	B	Open biotopes	Xerophilous	Autumn	Carnivorous
Platynini	<i>Calathus melanocephalus</i> (Linnaeus, 1758)	Least concern	C	Open biotopes	Mesophilous	Autumn	Carnivorous
Carabini	<i>Calosoma auropunctatum</i> (Herbst, 1784)	CZ: vulnerable SK: endangered	A	Open biotopes	Xerophilous	Spring	Carnivorous
Carabini	<i>Carabus granulatus</i> (Linnaeus, 1758)	Least concern	B	Silvicolous	Hygrophilous	Spring	Carnivorous
Carabini	<i>Carabus hortensis</i> (Linnaeus, 1758)	Least concern	A	Silvicolous	Hygrophilous	Autumn	Carnivorous
Carabini	<i>Carabus scheidleri</i> (Panzer, 1799)	Least concern	A	Silvicolous	Eurytopic	Autumn	Carnivorous
Carabini	<i>Carabus violaceus</i> (Linnaeus, 1758)	Least concern	A	Silvicolous	Eurytopic	Autumn	Carnivorous

Callistini	<i>Chlaenius nitidulus</i> (Schränk, 1781)	Least concern	B	Open biotopes	Hygrophilous	Spring	Carnivorous
Clivinini	<i>Clivina fossor</i> (Linnaeus, 1758)	Least concern	C	Open biotopes	Hygrophilous	Spring	Carnivorous
Platynini	<i>Dolichus halensis</i> (Schaller, 1783)	Least concern	B	Open biotopes	Xerophilous	Late summer/ autumn	Carnivorous

Table S3. *Cont.*

Tribe	Species	Incidence	Body size	Habitat affinity	Humidity affinity	Breeding period	Food specialization
Harpalini	<i>Harpalus affinis</i> (Schränk, 1781)	Least concern	C	Open biotopes	Eurytopic	Spring/summer/ autumn	Omnivorous
Harpalini	<i>Harpalus caspius roubali</i> (Schauberger, 1928)	Least concern	B	Open biotopes	Xerophilous	Autumn	Omnivorous
Harpalini	<i>Harpalus distinguendus</i> (Duftschmid, 1812)	Least concern	C	Open biotopes	Eurytopic	Spring/summer	Omnivorous
Harpalini	<i>Harpalus honestus</i> (Duftschmid, 1812)	Least concern	C	Open biotopes	Xerophilous	Spring	Omnivorous
Harpalini	<i>Harpalus rubripes</i> (Duftschmid, 1812)	Least concern	C	Open biotopes	Eurytopic	Spring	Omnivorous
Harpalini	<i>Harpalus signaticornis</i> (Duftschmid in 1812)	Least concern	C	Open biotopes	Xerophilous	Spring/autumn	Omnivorous
Harpalini	<i>Harpalus smaragdinus</i> (Duftschmid in 1812)	Least concern	C	Open biotopes	Xerophilous	Spring/autumn	Omnivorous
Harpalini	<i>Harpalus tardus</i>	Least concern	C	Open biotopes	Eurytopic	Spring	Omnivorous

(Panzer, 1797)							
Platynini	<i>Laemostenus terricola</i> (Herbst, 1784)	CZ: near threatened SK: least concern	B	Open biotopes	Xerophilous	Autumn	Carnivorous
Trechini	<i>Lasiotrechus discus</i> (Fabricius, 1792)	Least concern	D	Eurytopic	Hygrophilous	Spring	Carnivorous
Nebrini	<i>Leistus ferrugineus</i> (Linnaeus, 1758)	Least concern	C	Eurytopic	Hygrophilous	Autumn	Carnivorous
Loricerini	<i>Loricera pilicornis</i> (Fabricius, 1775)	Least concern	C	Eurytopic	Hygrophilous	Spring/summer	Carnivorous
Lebiini	<i>Microlestes maurus</i> (Sturm, 1827)	Least concern	D	Eurytopic	Eurytopic	Spring	Omnivorous
Lebiini	<i>Microlestes minutulus</i> (Goeze 1777)	Least concern	D	Eurytopic	Eurytopic	Spring	Carnivorous
Nebtriini	<i>Nebria brevicollis</i> (Fabricius, 1792)	Least concern	B	Eurytopic	Hygrophilous	Autumn	Carnivorous
Notiophilini	<i>Notiophilus aquaticus</i> (Linnaeus, 1758)	Least concern	D	Silvicolous	Hygrophilous	Spring/summer	Carnivorous

Table S3. Cont.

Tribe	Species	Incidence	Body size	Habitat affinity	Humidity affinity	Breeding period	Food specialization
Notiophilini	<i>Notiophilus palustris</i> (Duftschmid, 1812)	Least concern	D	Silvicolous	Hygrophilous	Spring	Carnivorous
Notiophilini	<i>Notiophilus pusillus</i> (G.R.Waterhouse, 1833)	Least concern	D	Eurytopic	Eurytopic	Spring	Carnivorous
Harpalini	<i>Ophonus azureus</i>	Least concern	C	Open biotopes	Xerophilous	Autumn	Granivorous

	(Fabricius, 1775)						
Harpalini	<i>Ophonus nitidulus</i> (Stephens, 1828)	Least concern	C	Open biotopes	Xerophilous	Spring	Granivorous
Platynini	<i>Platynus assimilis</i> (Paykull, 1790)	Least concern	C	Silvicolous	Hygrophilous	Spring	Carnivorous
Pterostichini	<i>Poecilus cupreus</i> (Linnaeus, 1758)	Least concern	B	Eurytopic	Eurytopic	Spring	Omnivorous
Pterostichini	<i>Poecilus lepidus</i> (Leske 1785)	Least concern	B	Eurytopic	Xerophilous	Spring	Carnivorous
Pterostichini	<i>Poecilus sericeus</i> (Fischer von Waldheim, 1824)	CZ: vulnerable SK: least concern	B	Eurytopic	Mesophilous	Spring	Carnivorous
Pterostichini	<i>Poecilus versicolor</i> (Sturm, 1824)	Least concern	C	Open biotopes	Hygrophilous	Spring	Carnivorous
Pterostichini	<i>Pseudoophonus griseus</i> (Panzer, 1796)	Least concern	C	Open biotopes	Mesophilous	Autumn	Omnivorous
Pterostichini	<i>Pseudoophonus rufipes</i> (De Geer)	Least concern	B	Open biotopes	Eurytopic	Autumn	Omnivorous
Pterostichini	<i>Pterostichus melanarius</i> (Illiger, 1798)	Least concern	B	Eurytopic	Mesophilous	Autumn	Carnivorous
Pterostichini	<i>Pterostichus melas</i> (Creutzer, 1799)	Least concern	B	Silvicolous	Mesophilous	Autumn	Carnivorous
Pterostichini	<i>Pterostichus niger</i> (Schaller, 1783)	Least concern	B	Silvicolous	Hygrophilous	Autumn	Carnivorous
Pterostichini	<i>Pterostichus nigrita</i> (Paykull, 1790)	Least concern	B	Eurytopic	Hygrophilous	Spring	Carnivorous
Pterostichini	<i>Pterostichus ovoideus</i>	Least concern	C	Silvicolous	Hygrophilous	Spring	Carnivorous

(Sturm, 1824)

Table S3. Cont.

Tribe	Species	Incidence	Body size	Habitat affinity	Humidity affinity	Breeding period	Food specialization
Pterostichini	<i>Pterostichus strenuus</i> (Panzer, 1796)	Least concern	C	Silvicolous	Hygrophilous	Spring	Carnivorous
Pterostichini	<i>Pterostichus vernalis</i> (Panzer, 1795)	Least concern	C	Eurytopic	Hygrophilous	Spring/summer	Carnivorous
Pterostichini	<i>Stomis pumicatus</i> (Panzer, 1795)	Least concern	C	Eurytopic	Hygrophilous	Spring	Carnivorous
Trechini	<i>Trechus quadristriatus</i> (Schränk, 1781)	Least concern	D	Open biotopes	Mesophilous	Autumn	Carnivorous
Harpalini	<i>Trichotichnus laevicollis</i> (Duftschmid, 1812)	Least concern	C	Silvicolous	Mesophilous	Spring/autumn	Omnivorous
Zabrinini	<i>Zabrus tenebrioides</i> (Goeze, 1777)	Least concern	B	Open biotopes	Xerophilous	Autumn	Omnivorous

¹ The tribe, species, body size, habitat and humidity affinity, breeding period [1], incidence [2,3] and food specializations [4] were determined. Body size (mid-range): A: >22 mm, B: 11–21.9 mm, C: 6–10.9 mm, D: <5.9 mm; Silvicolous: preferring woodlands; Open biotopes: preferring open areas; Eurytopic: adaptable to various environmental conditions; Hygrophilous: preferring moist places; Mesophilous: preferring intermediate or moderate environmental conditions, avoid extremes of moisture or dryness; Xerophilous: preferring dry environmental conditions.

References

1. Hürka, K. *Carabidae of the Czech and Slovak Republics*; Kabourek: Zlín, Czech Republic, 1996; p. 565.
2. Veselý, P.; Moravec, P.; Stanovský, J. Carabidae (střevlíkovití). In *Červený Seznam Ohrožených Druhů České Republiky*; Farkáč, J., Král, D., Škorpík, M., Eds.; AOPK: Praha, Czech Republic, 2005; pp. 406–411.
3. Holecová, M.; Franc, V. Red (Ecosozological) list of beetles (Coleoptera) of Slovakia. In *Red List of Plants and Animals of Slovakia. Ochr. Prir.* **2001**, 20 (Supp.), 111–128.

4. Larochelle, A. The food of carabid beetles (Coleoptera: Carabidae, including Cicindelinae). *Faberies* **1990**, *5*, 1–132.