

Sensitivity of core results to Rousseau et al. KS-window catalogue

Green p-value cells are significant at 0.05; red cells are not significant. The final column flags whether the significance decision is unchanged.

Event	Test	0.4-4 kyr	0.6-4 kyr	Same decision
Strong	Randomness: Poisson null	0.083 (n.s.)	0.460 (n.s.)	yes
Strong	Randomness: fixed-N null	0.049 (sig)	0.414 (n.s.)	no
Strong	Rayleigh: precession phase	0.0019 (sig)	0.0010 (sig)	yes
Strong	Poisson: LR04+CO2 vs stationary	0.0092 (sig)	0.012 (sig)	yes
Strong	Poisson: phase after LR04+CO2	0.0005 (sig)	0.0003 (sig)	yes
Strong	Poisson: LR04 after CO2+phase	0.045 (sig)	0.034 (sig)	yes
Strong	Poisson: CO2 after LR04+phase	0.905 (n.s.)	0.936 (n.s.)	yes
Strong	Poisson: full model vs stationary	6.4e-05 (sig)	5.0e-05 (sig)	yes
Weak	Randomness: Poisson null	0.028 (sig)	0.217 (n.s.)	no
Weak	Randomness: fixed-N null	0.0082 (sig)	0.167 (n.s.)	no
Weak	Rayleigh: precession phase	0.0027 (sig)	0.0012 (sig)	yes
Weak	Poisson: LR04+CO2 vs stationary	4.3e-05 (sig)	0.0012 (sig)	yes
Weak	Poisson: phase after LR04+CO2	0.0024 (sig)	0.0011 (sig)	yes
Weak	Poisson: LR04 after CO2+phase	0.505 (n.s.)	0.631 (n.s.)	yes
Weak	Poisson: CO2 after LR04+phase	0.057 (n.s.)	0.110 (n.s.)	yes
Weak	Poisson: full model vs stationary	1.8e-06 (sig)	1.9e-05 (sig)	yes