

Linear Regression with One Variable

TOTAL POINTS 5

1. Consider the problem of predicting how well a student does in her second year of college/university, given how well she did in her first year. 1 pt

Specifically, let x be equal to the number of "A" grades (including A-, A and A+ grades) that a student receives in their first year of college (freshmen year). We would like to predict the value of y , which we define as the number of "A" grades they get in their second year (sophomore year).

Here each row is one training example. Recall that in linear regression, our hypothesis is $h_{\theta}(x) = \theta_0 + \theta_1 x$, and we use m to denote the number of training examples.

x	y
5	4
3	4
0	1
4	3

For the training set given above (note that this training set may also be referenced in other questions in this quiz), what is the value of m ? In the box below, please enter your answer (which should be a number between 0 and 10).

4

2. Many substances that can burn (such as gasoline and alcohol) have a chemical structure based on carbon atoms; for this reason they are called hydrocarbons. A chemist wants to understand how the number of carbon atoms in a molecule affects how much energy is released when that molecule combusts (meaning that it is burned). The chemist obtains the dataset below. In the column on the right, "kJ/mol" is the unit measuring the amount of energy released. 1 point

Name of molecule	Number of hydrocarbons in molecule (x)	Heat release when burned (kJ/mol) (y)
methane	1	-890
ethene	2	-1411
ethane	2	-1560
propane	3	-2220
cyclopropane	3	-2091
butane	4	-2878
pentane	5	-3537
benzene	6	-3268
cyclohexane	6	-3920
hexane	6	-4163
octane	8	-5471
naphthalene	10	-5157

You would like to use linear regression ($h_{\theta}(x) = \theta_0 + \theta_1 x$) to estimate the amount of energy released (y) as a function of the number of carbon atoms (x). Which of the following do you think will be the values you obtain for θ_0 and θ_1 ? You should be able to select the right answer without actually implementing linear regression.

☐ $\theta_0 = -569.6, \theta_1 = 530.9$

☒ $\theta_0 = -569.6, \theta_1 = -530.9$

☐ $\theta_0 = -1780.0, \theta_1 = 530.9$

☐ $\theta_0 = -1780.0, \theta_1 = -530.9$


3. Suppose we set $\theta_0 = 0, \theta_1 = 1.5$. What is $h_\theta(2)$?

3

4. In the given figure, the cost function $J(\theta_0, \theta_1)$ has been plotted against θ_0 and θ_1 , as shown in 'Plot 2'. The contour plot for the same cost function is given in 'Plot 1'. Based on the figure, choose the correct options (check all that apply).

1 point

Plots for Cost Function $J(\theta_0, \theta_1)$



- ☐ Point P (The global minimum of plot 2) corresponds to point C of Plot 1.
- ☒ Point P (the global minimum of plot 2) corresponds to point A of Plot 1.
- ☐ If we start from point B, gradient descent with a well-chosen learning rate will eventually help us reach at or near point A, as the value of cost function $J(\theta_0, \theta_1)$ is maximum at point A.
- ☐ If we start from point B, gradient descent with a well-chosen learning rate will eventually help us reach at or near point C, as the value of cost function $J(\theta_0, \theta_1)$ is minimum at point C.
- ☒ If we start from point B, gradient descent with a well-chosen learning rate will eventually help us reach at or near point A, as the value of cost function $J(\theta_0, \theta_1)$ is minimum at A.

5. Suppose that for some linear regression problem (say, predicting housing prices as in the lecture), we have some training set, and for our training set we managed to find some θ_0, θ_1 such that $J(\theta_0, \theta_1) = 0$.

1 point

Which of the statements below must then be true? (Check all that apply.)

- ☐ Gradient descent is likely to get stuck at a local minimum and fail to find the global minimum.
- ☐ For this to be true, we must have $y^{(i)} = 0$ for every value of $i = 1, 2, \dots, m$.
- ☒ Our training set can be fit perfectly by a straight line.
i.e., all of our training examples lie perfectly on some straight line.
- ☐ For this to be true, we must have $\theta_0 = 0$ and $\theta_1 = 0$
so that $h_\theta(x) = 0$