

Condition/Medicat	Also Known As	Description	Type	Facial/Body/Massage	Nails	Fitness	Barber/Styling
ACSM Pre-Exercise Screening Algorithm	Exercise Preparticipation Screening, Medical Clearance for Exercise	American College of Sports Medicine algorithm using current exercise status, known cardiovascular/metabolic/renal disease, and symptoms to determine medical clearance requirements before moderate or vigorous exercise.	SCREENING PROTOCOL - Not a Condition	Not applicable.	Not applicable.	MANDATORY SCREENING PROTOCOL: Implement structured pre-participation screening aligned with ACSM algorithm. THREE KEY FACTORS: (1) Current regular exercise participation status, (2) Known cardiovascular/metabolic/renal disease (CVD/metabolic/renal), (3) Presence of signs/symptoms suggestive of disease. MEDICAL CLEARANCE NEEDED: Individuals with known CVD/metabolic/renal disease who want to begin vigorous exercise, OR any individual with concerning signs/symptoms (chest pain, unusual shortness of breath, dizziness, syncope, etc.) REGARDLESS of exercise history. SYMPTOMS = STOP and seek medical evaluation. Staff training essential to recognize warning signs. Document all screening.	Not applicable.
Acne (Active/Severe)	Acne Vulgaris, Breakouts, Blemishes	Inflammatory condition with comedones, papules, pustules, cysts; bacteria, oil, dead cells involved	RELATIVE - Modified treatment	Severe/cystic acne: gentle treatment only. No picking/squeezing cysts - scarring risk. Can perform extractions on whiteheads/blackheads only. Sanitize thoroughly.	Safe.	Safe and beneficial. Sweat can temporarily worsen - cleanse immediately after. Don't touch face during workout.	Safe. Use clean towels. Avoid comedogenic hair products near face.
Acrylate Allergy (Occupational & Client)	Methacrylate Sensitization, Gel Manicure Allergy, Occupational Contact Dermatitis	Acrylates are significant cause of allergic contact dermatitis; increasing prevalence with popularity of gel/acrylic nails affecting both clients and nail technicians; can cause permanent sensitization	TOTAL once sensitized + PREVENTIVE controls essential	Not applicable.	ONCE SENSITIZED: ABSOLUTE contraindication - continued exposure worsens dermatitis and triggers broader sensitization. Refer for medical evaluation and patch testing. PREVENTION FOR ALL: Engineering controls (ventilation systems meeting NIOSH recommendations), PPE (nitrile gloves - not latex/vinyl), proper product handling (avoid skin contact with uncured product), training on hazard communication. Staff developing hand dermatitis: immediate medical referral and exposure assessment. This is an OCCUPATIONAL HEALTH PRIORITY.	Not applicable.	Not applicable.
Acute Musculoskeletal Injuries	Acute Sprains, Strains, Fractures, Dislocations	Recent injury (typically <72 hours) with pain, swelling, inflammation; acute phase requires rest	LOCAL - Acute phase, then Medical Clearance	AVOID area for 48-72 hours (acute inflammatory phase). After acute phase: medical clearance for rehabilitation massage. Ice initially, heat later.	Avoid area if hand/foot injury until cleared.	TOTAL contraindication for affected area during acute phase. Medical/PT clearance for return. Rehabilitation protocol required.	Safe unless neck injury.
Anemia (Severe)	Iron Deficiency, Low Hemoglobin	Insufficient red blood cells or hemoglobin; causes fatigue, weakness, dizziness	Mild safe, Severe contraindicated	Mild anemia safe. Severe anemia contraindicated - increases cardiovascular stress. Client may tire easily.	Safe. Client may need breaks. Monitor for dizziness.	Severe anemia requires medical clearance. May need iron supplementation first. Low intensity initially. Monitor for excessive fatigue.	Safe. Client may tire easily.
Angina Pectoris	Chest Pain, Cardiac Angina	Chest pain from reduced blood flow to heart muscle; may indicate coronary artery disease	RELATIVE - Medical Clearance Required	Unstable angina is TOTAL contraindication. Stable angina requires medical permission. Gentle massage only. Keep nitroglycerin accessible.	Safe with medical clearance. Keep appointment duration reasonable. Have emergency protocol ready.	Requires medical clearance and stress testing. Exercise prescription from cardiologist. Emergency action plan required.	Safe with medical clearance. Keep sessions short. Avoid stressful situations.
Blood Thinners/Anticoagulants	Warfarin (Coumadin), Apixaban (Eliquis), Rivaroxaban (Xarelto), Dabigatran (Pradaxa), Enoxaparin (Lovenox), Aspirin, Clopidogrel (Plavix)	Medications preventing blood clotting; increases bleeding time and bruising risk from even minor trauma	RELATIVE - Special Precautions Required	Light to moderate pressure only - NO deep tissue. Increased bruising risk. Gentle techniques essential. Document medication.	Extra care required. Avoid aggressive cuticle work. No cutting - push back only. Use gentle techniques. Have styptic pencil ready.	Safe with precautions. Avoid contact sports. Wear protective gear. Alert trainer. Emergency protocol for bleeding episodes.	ELECTRIC RAZOR STRONGLY RECOMMENDED. If blade shaving: use extreme care, new blade, light pressure, styptic pencil ready. Apply pressure 10-15 min for any nicks. High dose (Lovenox, Warfarin) may require avoiding blade shaving entirely.
Bloodborne Pathogen Exposure Risk (Barber Services)	HIV, Hepatitis B, Hepatitis C Transmission Risk	HIV not transmitted through saliva or unbroken skin. Hepatitis B and C CAN spread through blood and via contaminated personal items (razors, nail clippers). Any nick/cut creates potential exposure.	PREVENTIVE - OSHA Standard Compliance Required	Any service that may cause bleeding requires standard precautions and immediate cleanup protocols.	CRITICAL SAFETY PROTOCOL: Any bleeding incident requires immediate response. Never share implements between clients without proper disinfection. Hepatitis C can survive on surfaces for up to 3 weeks.	Not applicable to equipment use, but relevant if offering any services that could cause skin breaks.	OSHA BLOODBORNE PATHOGENS STANDARD COMPLIANCE REQUIRED: Exposure Control Plan (written), Staff training (annual), Hepatitis B vaccination offered, Post-exposure protocols, Immediate cleanup of any blood with EPA-registered disinfectant, Contaminated tool protocols (immersion in disinfectant per label, proper storage), Single-use items (dispose immediately), Personal Protective Equipment available. Shaving services carry highest risk - new blade for each client, proper disposal of used blades in sharps container.
Bone Metastases	Cancer Spread to Bones, Skeletal Metastases	Cancer that has spread to bones; extremely fragile bones with high fracture risk from minimal pressure	TOTAL for pressure/manipulation	ABSOLUTE CONTRAINDICATION for massage. Even gentle pressure can cause fractures. Oncologist clearance required for any touch therapy. Light touch only if	Medical clearance required. Extremely gentle - bone fragility.	Medical clearance required. Non-weight bearing only. Physical therapy guided. High fracture risk.	Safe. Gentle positioning.
COPD (Chronic Obstructive Pulmonary Disease)	Emphysema, Chronic Bronchitis	Progressive lung disease causing breathing difficulty; severe cases struggle with humidity, heat, exertion	Severe = TOTAL for heat/humidity, Moderate = Medical Clearance	Severe: AVOID steam rooms, hot humid treatments. Moderate: medical clearance for heat. Position semi-reclined (not flat) - easier breathing. Monitor for respiratory distress.	Safe. May need breaks. Ensure client comfortable in positioning.	Severe: TOTAL contraindication without pulmonary rehabilitation program. Moderate: medical clearance required. Low-intensity only. Pulmonary rehab preferred. Emergency oxygen protocol.	Safe. Position for comfortable breathing. Avoid prolonged supine positioning.
COVID-19 or Fever	Coronavirus, Elevated Body Temperature, Active Infection	Any fever (>100.4°F/38°C) indicates active infection; massage/exercise increases	TOTAL until fever-free 24-48 hours	TOTAL contraindication. Reschedule appointment. Client/staff safety.	TOTAL contraindication. Reschedule.	TOTAL contraindication. Increases infection complications. Rest essential.	TOTAL contraindication. Reschedule.
Cancer (Active Treatment)	Malignancy, Undergoing Chemotherapy/Radiation	Malignant growth; treatment causes immunosuppression, fragile skin, fatigue, lymphedema risk	RELATIVE - Medical Permission Required	Medical clearance required. Gentle massage only. AVOID lymphatic drainage if lymphedema risk. NO massage near tumors or radiation sites. Immune compromised - perfect sanitation. Oncology massage certification ideal.	Medical permission advised. Immune compromised - extreme sanitation. Nails may be brittle from chemotherapy. Gentle work only.	Oncologist clearance required. Light activity beneficial for fatigue. May need to avoid gym during active chemotherapy (infection risk). Listen to body.	Safe with precautions. May have hair loss. Wig consultation/styling services appropriate.
Cold Sores (Herpes Simplex)	Fever Blisters, Oral Herpes, HSV 1	Viral infection causing fluid-filled blisters around mouth; highly contagious during	LOCAL - Active lesions	Active cold sores: AVOID face. Can treat body. Wait until crusted over before facial services.	Safe if not on hands.	Safe. Use personal towel.	Active outbreak: POSTPONE facial services. Cannot shave over lesions. Wait until healed.
Corticosteroids (Topical)	Hydrocortisone, Betamethasone, Clobetasol, Steroid Creams	Anti-inflammatory medications thinning skin with prolonged use; increased bruising risk	LOCAL - Avoid treated areas	Thin fragile skin where applied. Avoid deep pressure. Increased bruising risk.	AVOID waxing areas using topical steroids - skin tearing risk. Wait 3 months after stopping.	Safe.	Safe but gentle on affected areas.
Deep Vein Thrombosis (DVT)	Blood Clot in Leg, Venous Thrombosis	Blood clot in deep vein (usually leg); risk of pulmonary embolism if clot dislodges	TOTAL for 3-6 months post-diagnosis	TOTAL contraindication for 3-6 months. After clearance, gentle massage only. Never massage legs with suspected DVT. Client/therapist safety critical.	AVOID pedicures for 3-6 months. After clearance, gentle foot massage only. Never massage calves if DVT history.	AVOID exercise during acute phase. Medical clearance required before resuming. Light activity only initially.	Safe for head/facial services.
Diabetes Mellitus (Type 1 & 2)	Blood Sugar Disease, Insulin Resistance	Impaired glucose regulation; complications include neuropathy, poor circulation, slow wound healing, increased infection risk	RELATIVE - Special Precautions	Generally safe. Monitor blood glucose before treatment. Have fast-acting carbs available. Gentle pressure - increased bruising risk. Watch for neuropathy.	High risk services. Inspect feet thoroughly for wounds/infections (can be painless due to neuropathy). Absolutely no cutting calluses. Gentle filing only. Perfect sanitation critical - slow healing. Physician clearance recommended for foot services.	Medical clearance recommended but not required for low-moderate intensity. Monitor blood glucose before/after. Should not exercise if glucose >250 mg/dL with ketones. Have glucose tablets available. Type 1: higher hypoglycemia risk - may need carb adjustment or insulin reduction.	Safe. Extra care with heat tools near face. Scalp checks for wounds.

Diabetic Foot Risk - High Priority	Diabetes with Peripheral Arterial Disease (PAD), Diabetic Neuropathy, High-Risk Diabetic Foot	ADA explains PAD reduces blood flow + neuropathy reduces sensation = dramatically increased ulcer, infection, and amputation risk. APMA advises IMMEDIATE evaluation if drainage or excessive redness present.	RELATIVE - Conservative Care ONLY + High Referral Threshold	Conservative approach. Check visible skin integrity. Refer ANY open wounds or infection signs. Document modifications. Avoid high-heat treatments on feet.	HIGHEST RISK DEPARTMENT FOR DIABETIC COMPLICATIONS: "Minor" pedicure trauma can become serious ulcer. PROTOCOL: Visual foot inspection BEFORE service - any wounds, ulcers, drainage, excessive redness, swelling, warmth, discoloration = DEFER and URGENT REFERRAL. If cleared to proceed: CONSERVATIVE COSMETIC CARE ONLY (no blades, no aggressive filing, no callus cutting), gentle techniques, avoid heat on feet (no hot paraffin), proper footwear emphasis. CLIENT EDUCATION: daily foot checks, proper footwear, immediate medical attention for any wounds. DOCUMENT thoroughly: foot risk screening, modifications made, any findings. Many jurisdictions require physician clearance for diabetic foot services. Consider this HIGH LIABILITY - when in doubt, defer and refer.	Proper footwear ESSENTIAL. Foot inspection before/after each session. Avoid activities requiring fine foot sensation if neuropathy severe. Swimming/cycling (non-weight bearing) may be safer than running. Medical clearance recommended. Emergency protocol for altered mental status (hypoglycemia risk with exercise).	Not applicable.
Diabetic Neuropathy	Nerve Damage from Diabetes, Peripheral Neuropathy	Nerve damage causing numbness, tingling, pain; impaired sensation means injuries may go unnoticed	RELATIVE - Modified Treatment	Client may not feel pain - reduce pressure. Check skin integrity before/after. Avoid hot stones - cannot sense temperature.	HIGH RISK. Client cannot feel injuries. Visual inspection critical. Gentle filing only - NO cutting. Physician referral for any lesions. Document condition.	Proper footwear essential. Check feet before/after. Non-weight-bearing exercise preferred if severe. Balance training important.	Safe. Extra caution with heat.
Eczema/Atopic Dermatitis	Atopic Eczema, Inflammatory Skin Condition	Chronic inflammatory condition causing dry, itchy, red patches; impaired barrier function	RELATIVE - Modified treatment	AVOID active flares. During remission: gentle pressure, hypoallergenic products, avoid fragrances. Can trigger reaction.	Avoid active areas. Use gentle, fragrance-free products. May have nail pitting/ridging.	Safe. Sweat may irritate - shower immediately after. Moisturize. Wear soft clothing.	Avoid affected scalp areas. Use gentle, fragrance-free products.
Epilepsy/Seizure Disorders	Convulsions, Fits	Neurological condition causing seizures; certain triggers include flashing lights, stress, fatigue	RELATIVE - Safety precautions	Generally safe if controlled. Avoid triggers (strong scents, flashing lights). Have seizure protocol. Relaxation beneficial.	Safe. Have emergency protocol. Remove tools if seizure occurs.	Medical clearance if uncontrolled. Avoid activities where seizure would cause injury (climbing, swimming alone). Buddy system. Staff trained on seizure response.	Safe. Avoid flashing lights. Have emergency protocol.
Fibromyalgia	Chronic Pain Syndrome, FM	Chronic widespread musculoskeletal pain with tender points; extreme sensitivity to pressure	RELATIVE - Light Pressure Only	Medical clearance advised. Client extremely sensitive to pressure - communicate constantly. Light to medium pressure only. What feels light to therapist may feel intense. Heat generally beneficial. Very slow, gentle techniques.	Safe. Client may be sensitive to pressure - be gentle.	Medical clearance recommended. Low-impact exercise beneficial (water aerobics, walking, gentle yoga). Avoid overexertion - pacing essential. Start very gently.	Safe. Client may be sensitive to prolonged positioning - offer breaks.
Formaldehyde Exposure (Hair Smoothing)	Keratin Treatment Formaldehyde Release, Brazilian Blowout Exposure	OSHA documented that air testing in some salons during smoothing treatments exceeded OSHA short-term exposure limit; FDA warns that some products release formaldehyde when heated despite "formaldehyde-free" marketing claims	OCCUPATIONAL HEALTH HAZARD - Controls Required	Not applicable.	Not applicable.	Not applicable.	CRITICAL WORKER SAFETY ISSUE: Formaldehyde is a known carcinogen. OSHA requires: Hazard Communication Program (SDS availability, product labeling, staff training), Air monitoring when exposure suspected, Engineering controls (local exhaust ventilation at styling stations), PPE when controls insufficient. IMMEDIATE: Check product labels and SDS. Consider product substitution to truly formaldehyde-free alternatives. NEVER rely on marketing claims - verify with SDS. Symptoms (eye/nose/throat irritation, breathing difficulty, headaches) indicate overexposure. CLIENT CONSIDERATIONS: Inform of potential exposure. Ensure ventilation. Those with asthma/respiratory conditions at higher risk.
Fungal Nail Infection (Onychomycosis)	Nail Fungus, Tinea Unguium, Ringworm of Nail	Fungal infection causing nail thickening, discoloration, crumbling; highly contagious through contaminated tools	TOTAL until cleared by physician	Generally not applicable unless systemic fungal infection.	ABSOLUTE CONTRAINDICATION. Cannot perform services. High cross-contamination risk. Requires antifungal treatment (2-6 months). Refer to physician/podiatrist. Can spread to other nails and clients through tools.	Safe - not transmitted through equipment. Keep affected area covered.	Not applicable.
Head Lice Infestation	Pediculosis Capitis, Lice, Nits	Parasitic insects feeding on human blood, found mostly on scalp; spread via close contact and shared items (brushes, combs, capes, headrests)	TOTAL - Highly Contagious	Not typically applicable unless client rests head on treatment table.	Not applicable.	Low risk unless shared headgear/helmets used.	ABSOLUTE CONTRAINDICATION: Defer ALL services immediately. High transmission risk via combs, brushes, capes, chairs. PROTOCOL: Instruct client to seek CDC-recommended treatment. Launder ALL capes/linens (hot water/high heat dry). Disinfect ALL tools per regulations (immersion in EPA-registered disinfectant). Disinfect chair, headrest, cape clips. Store tools properly post-disinfection. DOCUMENT deferral. Do not proceed until client shows proof of completed treatment and no live lice/nits present on inspection.
Heart Failure (Congestive Heart Failure)	CHF, Cardiac Failure	Heart unable to pump sufficient blood to meet body needs; causes fluid buildup in lungs and extremities	RELATIVE - Medical Clearance Required	Medical permission required. Gentle massage only to avoid stressing weak heart. Monitor for shortness of breath. No deep pressure.	Safe with precautions. Elevate legs if swelling present. Watch for respiratory distress.	Requires cardiology clearance. Cardiac rehabilitation preferred. Low intensity only with close monitoring.	Safe. Position comfortably to ease breathing. Keep sessions reasonable duration.
Hemophilia	Bleeding Disorder, Clotting Factor Deficiency	Genetic disorder preventing normal blood clotting due to missing clotting factors	SEVERE = TOTAL, MILD = Medical Clearance	Severe hemophilia is TOTAL contraindication. Mild forms require medical clearance and factor replacement before treatment. Extremely gentle techniques only.	Severe hemophilia contraindicated. Mild requires medical consultation. Absolutely no cutting. Visual services only.	Requires hematologist clearance. Non-contact activities only. Swimming, cycling safe. Must have emergency plan.	Severe hemophilia: electric razor only or beard growth. Mild: medical consultation required. Have hemostatic agent available.
Hydrophobia (Fear of Water)	Aquaphobia, Water Phobia	Exaggerated, irrational fear of water causing extreme psychological distress	TOTAL for water immersion	Safe for dry treatments. Can use hydrotherapy tubs if client comfortable and in control.	May be uncomfortable with foot soaks - offer dry pedicure option.	ABSOLUTE CONTRAINDICATION for pools/hydrotherapy. Forcing immersion causes severe distress - not therapeutic. Dry land exercise only.	Safe. May be anxious during shampooing - communicate.
Hypertension (High Blood Pressure)	HBP, Elevated Blood Pressure	Blood pressure consistently above 140/90 mmHg; uncontrolled hypertension increases risk of cardiovascular events during	RELATIVE	Avoid deep abdominal massage. Require medical clearance if uncontrolled. Light to moderate pressure only. Avoid hot stone therapy.	Generally safe. Monitor for dizziness. Avoid prolonged heat exposure (paraffin).	Medical clearance required if symptomatic or uncontrolled. Moderate intensity safe if controlled. Monitor blood pressure before/after sessions.	Generally safe. Avoid excessive heat from dryers in confined spaces.
Hypotension (Low Blood Pressure)	Low BP, Postural Hypotension	Blood pressure below 90/60 mmHg; increased risk of dizziness, fainting, and falls	RELATIVE	Extra care needed. Assist client when standing. Avoid sudden position changes. Keep head elevated during treatment.	Safe with precautions. Ensure client eats before appointment. Assist with position changes.	Medical consultation recommended. Start with low intensity. Monitor for dizziness. Avoid rapid position changes.	Safe. Assist client when standing from reclined position. Avoid sudden movements.
Impetigo	School Sores, Bacterial Skin Infection	Highly contagious bacterial infection with honey-colored crusted sores	TOTAL until treated	TOTAL contraindication. Highly contagious. Requires antibiotic treatment. Wait until cleared.	TOTAL contraindication.	TOTAL contraindication.	TOTAL contraindication.
Incontinence (Urinary/Fecal)	Loss of Bladder/Bowel Control	Inability to control urination or bowel movements; strict hygiene concern for water facilities	TOTAL for water-based services	Safe for dry treatments. Ensure client comfortable. Waterproof protection may be needed for massage tables.	Safe.	ABSOLUTE CONTRAINDICATION for pools/hydrotherapy. Hygiene regulation - prevents water contamination. Dry land exercise safe.	Safe.
Isotretinoin (Accutane) - Updated Guidance	Accutane, Roaccutane, Oral Retinoid Therapy	Powerful oral retinoid causing extreme skin thinning and fragility. UPDATED: Recent dermatologic surgery evidence indicates superficial chemical peels and nonablative lasers may not require historic 6-12 month delay, but risk tolerance and operator training matter critically.	RELATIVE - Timing depends on procedure depth	DURING TREATMENT: Skin extremely fragile - gentle techniques only, no aggressive exfoliation. WAIT TIMES POST-TREATMENT: Superficial peels/nonablative procedures: emerging evidence suggests shorter delays possible with proper training and conservative approach. Medium-deep peels, ablative lasers, dermabrasion: 6-12 months post-treatment still recommended. Waxing: AVOID during treatment and 6 months after. Document medication, treatment dates, and obtain informed consent for any resurfacing.	Skin may be fragile if using on body. Avoid aggressive cuticle work. No waxing.	Safe. Sun protection critical - photosensitivity increased.	Safe for cutting. NO waxing during treatment or for 6 months after completion. Chemical services: use caution - scalp may be sensitive.

Legionella Risk (Hot Tub/Water Features)	Legionnaires' Disease Risk, Legionella Pneumophila	CDC documented hot tub-associated Legionnaires' disease outbreaks; proper disinfectant residual and pH reduce spread of this potentially fatal pneumonia	PREVENTIVE - Water Management Program Required	FACILITY PROTOCOL: Water management program for hot tubs/steam rooms. Regular monitoring. Immediate response to detection. Counsel vulnerable clients (immunocompromised, elderly, smokers) about risks.	Not applicable unless facility has water features.	CRITICAL FACILITY SAFETY: Hot tubs associated with Legionnaires outbreaks. Implement water management program. Monitor disinfectant and pH. Vulnerable populations at higher risk. Emergency protocol if suspected exposure - clients should seek medical care for respiratory symptoms post-exposure.	Not applicable.
Leukemia	Blood Cancer, Hematological Malignancy	Cancer of blood-forming tissues; impaired immune function and clotting	RELATIVE - Medical Permission Required	Require medical permission. Gentle massage only. Immune compromised - strict hygiene essential. Watch for easy bruising.	Medical clearance required. Immune compromised - perfect sanitation critical. Gentle techniques only.	Oncologist clearance required. May need to avoid gym during chemotherapy. Light activity. Fatigue management.	Medical clearance advised. Extra sanitation. Gentle techniques. May have hair loss from treatment.
Lupus (Systemic Lupus Erythematosus)	SLE, Autoimmune Disease	Autoimmune disease affecting multiple organs; extreme UV sensitivity, fatigue, inflammation, "butterfly rash" on face	RELATIVE - Medical Clearance Required	Medical clearance required. Gentle treatment only during remission. Avoid active flares. No aggressive exfoliation. HIGH SPF essential post-treatment. Be aware of photosensitivity.	Medical clearance advised. Gentle techniques. May have Raynaud's phenomenon (avoid temperature extremes).	Medical clearance required. Low-impact beneficial. Avoid overexertion (triggers flares). Sun protection essential for outdoor activity. Listen to fatigue levels.	Safe with precautions. Be aware of hair loss (common). Gentle scalp care. Sun protection if spending time outdoors post-appointment.
Lymphedema	Lymphatic Swelling	Chronic swelling from impaired lymphatic drainage; common after lymph node removal	RELATIVE - Specialized treatment only	Requires certified lymphedema therapist. AVOID deep tissue on affected limb. Can worsen condition if improper technique.	Extra infection risk in affected limb. Gentle work only. Perfect sanitation. Blood pressure/injections avoided in limb - same for aggressive nail work.	Exercise beneficial with compression garments. Progressive resistance training safe. Start slowly.	Safe.
Lymphedema - Specialized Protocol	Secondary Lymphedema, Post-Cancer Lymphedema, Lymphatic Insufficiency	International Society of Lymphology notes vigorous classical massage may damage lymphatic vessels or attachments. Requires specialized manual lymphatic drainage (MLD) training - NOT standard massage.	RELATIVE - Specialized Training REQUIRED	STANDARD DEEP TISSUE CONTRAINDICATED: Can worsen swelling and damage fragile lymphatic system. ONLY certified lymphedema therapists should perform MLD. For regular massage therapists: AVOID affected limb entirely OR use only very gentle, superficial techniques with medical clearance. Heat: AVOID on affected limb - worsens swelling. Screen for recent cellulitis (common complication) - if suspected infection, urgent medical referral. Never use compression without proper training.	Affected limb has extra infection risk (compromised lymphatic clearance). GENTLE techniques only. Perfect sanitation critical. Avoid blood pressure cuffs, injections, blood draws, IV placement, tight jewelry AND aggressive nail work on affected limb. Document which limb affected.	Exercise BENEFICIAL with proper protocol: compression garment required during exercise, progressive resistance training safe, start slowly and build gradually. Avoid overheating affected limb. Swimming excellent option. Medical/PT clearance recommended.	Safe. If arm affected: comfortable positioning, avoid prolonged arm elevation.
Multiple Sclerosis (MS)	Demyelinating Disease	Autoimmune disease affecting central nervous system; heat sensitivity common (Uhthoff's phenomenon - heat worsens symptoms), impaired sweating, fatigue, mobility issues	RELATIVE - Medical Clearance, Temperature Restrictions	Medical clearance recommended. AVOID heat therapies (hot stone, sauna, steam) - can trigger symptom flare. Cool/lukewarm temperatures only. Gentle pressure. Very beneficial for relaxation and symptom management when properly modified.	Safe. Avoid paraffin wax (heat). Accommodate mobility limitations.	Medical clearance required. AVOID overheating - exercise in cool environment, use cooling vest/towels. Aquatic therapy excellent (cool water). Beneficial with proper modifications. Monitor for fatigue.	Safe. Avoid excessive heat from dryers. Cool water preferred. Accommodate positioning for mobility issues.
Oncology/Cancer Treatment - Comprehensive Protocols	Active Cancer, Chemotherapy, Radiation Therapy, Immunotherapy	Society for Oncology Massage (\$40M) defines oncology massage as modification of existing massage to safely work with cancer-affected individuals. Requires advanced training, careful pressure decisions, tumor site awareness, lymph node removal/radiation history consideration.	RELATIVE - Specialized Training Required	REQUIRES ONCOLOGY-TRAINED STAFF: Never massage over tumor sites. Bone metastases: extreme fracture risk from even light pressure - ABSOLUTE contraindication or extremely light touch only with oncologist clearance. Post-mastectomy/lymph node removal: AVOID vigorous massage on affected limb - can worsen lymphedema. Requires certified lymphedema therapy for those areas. Chemotherapy: immunosuppressed - perfect sanitation essential. Radiation sites: skin may be fragile/burned - avoid area. Pressure decisions critical - some clients need very light touch, others can tolerate moderate. COORDINATE with medical team when possible. Positioning adaptations essential.	Medical clearance recommended. Immunosuppressed - extreme sanitation protocols. Chemotherapy causes nail changes (brittleness, ridging, discoloration, detachment). Gentle techniques only. NO aggressive cutting/filing. Bleeding risk may be increased (thrombocytopenia). Have hemostatic agents ready.	Oncologist clearance REQUIRED. Light activity generally beneficial for fatigue management. May need to avoid gym during active chemotherapy (infection risk in immunosuppressed state). Physical therapy coordination preferred. Listen to fatigue levels - energy unpredictable. Bone metastases: non-weight bearing only, high fracture risk.	Safe with compassion. Hair loss common with chemotherapy - may need wig consultation/styling services, gentle scalp care, or head covering options. Scalp may be sensitive. Chemical services: consult oncologist first - skin may be fragile from treatment.
Open Wounds/Cuts	Abrasions, Lacerations, Broken Skin	Break in skin integrity; infection risk	LOCAL - Avoid area completely	Avoid area completely until healed. Infection risk.	CANNOT service if near/on nail area. Reschedule.	Cover completely. May need to avoid certain exercises.	Avoid area completely. Cannot shave over cuts.
PPD Hair Dye Allergy	Para-Phenylenediamine Allergy, Permanent Hair Dye Contact Dermatitis	PPD is frequent contact sensitizer in permanent hair dyes; presentations may be acute, subacute, or chronic. Re-exposure produces faster, more severe reactions with possible severe facial swelling in sensitized individuals - OPERATIONALLY CRITICAL	TOTAL once sensitized	Not applicable.	Not applicable.	Not applicable.	ONCE SENSITIZED: ABSOLUTE contraindication to re-exposure. Risk of severe allergic reaction including facial swelling, respiratory distress. Refer for dermatologic evaluation and formal patch testing. DOCUMENT: exact product used, reaction timeline, severity. Future risk management requires avoiding ALL PPD-containing products and related aromatic amines. Cross-reactivity possible with related chemicals. Patch test MANDATORY before any future color services if client has ANY history of dye reaction.
Pacemaker/ICD (Implantable Cardioverter	Cardiac Device, Heart Device	Implanted electronic device regulating heart rhythm or providing emergency shocks	RELATIVE - Equipment Restrictions	Avoid vigorous massage over device site. Hot stone magnets and RF devices contraindicated - can interrupt pacing.	Safe. No restrictions for nail services.	Safe with medical clearance. Avoid contact sports. Some equipment restrictions. Technogym equipment generally safe.	Safe. No restrictions.
Paralysis	Loss of Motor Function, Hemiplegia, Paraplegia, Quadriplegia	Loss of muscle function in part of body; may have altered sensation, circulation issues, pressure sore risk	RELATIVE - Positioning & Pressure Precautions	Medical clearance advised. Extra care with positioning - pressure sore prevention. May have altered sensation. Can be very beneficial. Specialized training helpful.	Safe. Accommodate mobility limitations. Extra care if sensation impaired.	Medical clearance required. Adaptive equipment essential. Physical therapy coordination. Can be highly beneficial.	Safe. Accommodate positioning needs. Transfer assistance may be needed.
Parkinson's Disease	PD, Parkinsonian Syndrome	Progressive neurological disorder causing tremors, rigidity, slow movement, balance problems	RELATIVE - Safety & Positioning Precautions	Medical clearance advised. Can be very beneficial for rigidity. Extra time for positioning. Fall prevention. May have difficulty with face-down positioning.	Safe. Tremors may make services challenging - work around them. Extra time needed. Ensure client comfortable.	Medical clearance recommended. Exercise highly beneficial. Balance exercises important. Fall prevention protocols. Avoid activities requiring quick reactions.	Safe. Tremors may require extra time. Head support may be helpful. Patient, gentle approach.
Paronychia	Nail Fold Infection, Cuticle Infection	Bacterial/fungal infection around nail causing redness, swelling, pus, pain	TOTAL for affected digit	Local contraindication for hands/feet if present.	ABSOLUTE CONTRAINDICATION for affected nail. Can spread infection. May require antibiotics. Refer to physician. Can service other nails with extreme caution and perfect sanitation.	Safe if wound covered.	Not applicable.
Pedicure-Associated Nontuberculous Mycobacterial Infection Risk	Mycobacterium fortuitum, Rapidly Growing Mycobacteria (RGM), Footbath Furunculosis	Bacterial infections from rapidly growing mycobacteria found in inadequately cleaned whirlpool footbaths; CDC-documented outbreak investigations linked to spa pedicure equipment	PREVENTIVE - Equipment Safety Protocol Required	Not applicable to this department.	CRITICAL SAFETY PROTOCOL: Rigorous, auditable footbath cleaning/disinfection required. Clean and disinfect after EACH client per manufacturer instructions. Daily filter/screen removal and cleaning. Date/time-stamped documentation logs MANDATORY in many jurisdictions. Equipment must be removed from service if documentation lapses. High proportion of footbaths have tested positive for mycobacteria in outbreak investigations.	Not applicable unless facility has pedicure services.	Not applicable.
Peripheral Neuropathy	Nerve Damage, Loss of Sensation	Nerve damage causing numbness, tingling, or complete loss of sensation; cannot feel pain, temperature, or pressure	RELATIVE - Cannot provide feedback	Client cannot feel if pressure too deep or temperature too hot. Reduce pressure significantly. Avoid heat treatments - burn risk. Constant visual skin checks. Document condition.	HIGH RISK - especially feet. Client cannot feel injuries. Visual inspection critical. Gentle techniques only. No callus removal. Physician referral for any lesions.	Proper footwear essential. Check feet before/after for injuries. Avoid activities requiring fine sensation. Balance may be affected.	Safe. Extra caution with heat tools.
Photosensitizing Medications	Doxycycline, Ciprofloxacin, Hydrochlorothiazide, NSAIDs	Medications increasing UV sensitivity and sunburn risk	RELATIVE - Sun exposure precautions	High SPF essential post-treatment. Avoid treatments with extended UV exposure risk.	Be aware for outdoor post-appointment exposure.	Outdoor exercise: high SPF sunscreen essential. May need to adjust to indoor activities.	Educate client on sun protection.

Post-Partum & Breastfeeding	Postpartum Period, Lactation, Nursing Mother	Period after childbirth including while breastfeeding; ongoing hormonal changes, risk of infection if stitches present (C-section/episiotomy), treatment chemicals may pass into breast milk	RELATIVE - Modified Treatment	Avoid abdominal massage until cleared (6-8 weeks post-birth, longer for C-section). Light massage only. Avoid essential oils that may affect milk supply or pass to baby (peppermint, sage, parsley). Position comfortably for nursing mothers.	Safe with precautions. Ensure good ventilation - minimize fume exposure. Avoid strong chemical treatments.	Medical clearance for exercise resumption (6 weeks vaginal birth, 8-12 weeks C-section). Pelvic floor considerations. Start gently. Stay hydrated for milk supply.	Safe. Ensure good ventilation with chemical services. Heat treatments on chest may affect milk supply/letdown.
Pregnancy	Expecting, Prenatal, Gravid	Increased relaxin hormone softens tissues; higher infection risk; specific trimester considerations	RELATIVE - Modified treatments	AVOID 1st trimester (high miscarriage risk). 2nd/3rd: specialized prenatal massage only. Side-lying position. Avoid deep abdominal work. No essential oils (clary sage, rosemary, juniper). Avoid pressure points (SP6, LI4). Heat contraindicated (hot stone, sauna, steam).	Safe with precautions. Pregnancy hormones may affect polish wear. Avoid strong fumes - ensure ventilation. No waxing - skin more sensitive. Gentle filing only on brittle nails. Comfortable positioning essential. Avoid paraffin (heat).	20-30 min moderate intensity most days recommended. Benefits: reduced gestational diabetes risk, better cardiovascular health, easier delivery. Avoid supine position after 1st trimester. No contact sports. Stay hydrated. Stop if bleeding, dizziness, chest pain, contractions. Pregnancy-specific training preferred.	Safe. Ensure comfortable positioning. Minimize strong chemical fumes. Avoid overheating under dryers.
Pseudomonas ("Greenies")	Green Nail Syndrome, Bacterial Nail Infection	Bacterial infection under nail enhancement causing green discoloration; trapped moisture creates breeding ground	TOTAL - Removal required	Not applicable.	REMOVE all enhancements immediately. Clean and dry nail. Do not reapply product until greenie grows out completely (2-4 months). Not contagious but indicates service issue. Educate client on moisture prevention.	Safe.	Not applicable.
Pseudomonas Folliculitis (Hot Tub Rash)	Hot Tub Folliculitis, Whirlpool Dermatitis	Pruritic follicular skin eruption from Pseudomonas aeruginosa exposure in poorly maintained hot tubs/whirlpools; prevention requires maintaining proper disinfectant levels and pH	PREVENTIVE - Water Quality Protocol	If client has active rash: defer wet wraps and heavy occlusion - may worsen discomfort and risk secondary infection. Hydrotherapy services can be the exposure source. For facilities with hot tubs: follow CDC healthy-swimming parameters for disinfectant and pH monitoring. If outbreak suspected, suspend use and review maintenance logs.	Defer services if active rash on hands/feet. Can worsen with occlusion.	FACILITY PROTOCOL: Monitor hot tub/spa water quality continuously. CDC parameters: proper chlorine/bromine levels and pH. Maintenance logs required. If outbreak suspected, close facility and investigate. Symptomatic clients: refer to medical care if systemic symptoms develop.	Not applicable.
Psoriasis	Plaque Psoriasis, Autoimmune Skin Condition	Autoimmune condition causing thick, scaly, red patches; Koebner phenomenon (trauma triggers new lesions)	RELATIVE - Avoid active plaques	Avoid active plaques - trauma can trigger new lesions. Can massage around areas gently.	Psoriatic nails common (pitting, onycholysis, subungual hyperkeratosis). Very gentle work. No aggressive techniques.	Safe and beneficial. May improve with exercise. Sweat may irritate - shower after.	Scalp psoriasis: avoid affected areas. Use medicated shampoos if physician-prescribed.
Recent Botox/Dermal Fillers	Injectable Neurotoxins, Facial Injectables, Cosmetic Injectables	Recent cosmetic injections; massage can displace product from intended area, reducing efficacy or causing unwanted migration	LOCAL - Time Restrictions	Botox: avoid massage/manipulation for 24-48 hours, no lying flat for 4 hours. Dermal Fillers: avoid massage for 2-4 weeks. No facial massage, facials, or heat treatments during restriction period. Can displace product.	Safe.	Botox: avoid strenuous exercise 24 hours. No inverted positions (yoga headstands). Otherwise safe.	Safe but avoid face-down positioning during restriction periods.
Recent Heart Attack (Myocardial Infarction)	MI, Cardiac Arrest	Blocked blood flow to heart causing muscle damage; high recurrence risk in first 2-3 months	TOTAL for 2-3 months, then Medical Clearance	TOTAL contraindication for 2-3 months due to high recurrence risk. After period, require medical clearance. Gentle massage only.	Generally safe after acute period. Keep relaxing. Monitor for chest pain.	TOTAL contraindication until cardiac rehabilitation clearance. Supervised cardiac rehab program required initially.	Safe after acute period. Low-stress environment.
Retinoids (Topical)	Tretinoin (Retin-A), Adapalene (Differin), Tazarotene (Tazorac), Isotretinoin (Accutane)	Vitamin A derivatives thinning stratum corneum; increased sensitivity, fragility, peeling	RELATIVE - Wait periods required	Topical: discontinue 3-7 days before exfoliating facials, chemical peels, waxing, microdermabrasion. Accutane: wait 6-12 months after completion before ANY resurfacing. Extreme fragility/sensitivity.	If using on hands: very thin skin - avoid waxing. Otherwise safe.	Safe.	Safe for haircuts. Avoid facial waxing.
Rheumatoid Arthritis	RA, Inflammatory Arthritis	Autoimmune disease causing joint inflammation, pain, stiffness; joints may be swollen, hot, extremely tender	RELATIVE - Gentle Treatment Required	Medical clearance advised. AVOID massage directly on inflamed joints during flare. Gentle massage around area beneficial. Heat during remission, cold during flares. Very light pressure only.	Hand/foot arthritis common. Extra gentle with joint manipulation. Accommodate limited mobility. Warm water soaks beneficial (not during flare).	Medical clearance recommended. Low-impact exercise beneficial (swimming, water aerobics, tai chi). Avoid high-impact during flares. Range of motion exercises important.	Safe. Accommodate neck mobility limitations for shampooing/styling.
Ringworm (Tinea)	Fungal Skin Infection, Dermatophytosis	Fungal infection causing circular, red, scaly patches; contagious	LOCAL contraindication	Avoid affected area. Can spread through contact. High sanitation required.	Avoid affected area completely. If widespread, postpone.	Cover lesions. Use personal towel/mat. Clean equipment after use.	Scalp ringworm (tinea capitis) prevents services. Body ringworm: avoid area.
Rosacea	Acne Rosacea, Adult Acne	Chronic inflammatory condition causing facial redness, visible blood vessels, papules/pustules; triggers include heat,	RELATIVE - Gentle treatment required	Gentle techniques only. Avoid heat, steam, hot towels. No aggressive exfoliation. Use cooling products. Avoid trigger ingredients (fragrance, alcohol, menthol).	Generally safe.	Exercise triggers flushing. Cool environment. Use cooling towel. Lower intensity. Still beneficial overall.	Avoid hot tools near face. Use gentle products. Cool water.
Scleroderma/Morphea	Systemic Sclerosis, Localized Scleroderma	Autoimmune disease causing skin hardening and tightening; Raynaud's phenomenon common	RELATIVE - Resurfacing Contraindicated	Medical clearance required. CONTRAINDICATED for aggressive resurfacing treatments (microdermabrasion, chemical peels). Gentle massage may help with skin pliability. Avoid temperature extremes (Raynaud's).	Extra gentle - poor circulation. Raynaud's: avoid cold, ensure warmth. May have tight skin around fingers.	Medical clearance required. Gentle stretching beneficial for skin pliability. Avoid temperature extremes.	Safe. May have facial skin tightness affecting expressions.
Severe Chlorine Allergy	Pool Chemical Hypersensitivity	Severe allergic reaction to chlorine or pool chemicals causing respiratory distress or skin reactions	TOTAL for chlorinated pools	Safe for non-pool services.	Safe.	ABSOLUTE CONTRAINDICATION for chlorinated pools/hydrotherapy. Can cause severe respiratory distress. Salt water or non-chlorinated alternatives may be safe with testing.	Safe.
Stroke/CVA	Brain Attack, Cerebrovascular Accident	Disrupted blood flow to brain causing cell death; high recurrence risk in first 6 months	TOTAL for 6 months, then Medical Clearance	TOTAL contraindication for 6 months post-stroke due to recurrence risk. After period, medical clearance required. Be aware of paralysis/weakness.	Safe after acute period. Accommodate mobility limitations. Extra support may be needed.	Medical clearance with progressive rehabilitation required. May need adaptive equipment. Physical therapy guided.	Safe after acute period. Accommodate head/neck positioning limitations.
Sunburn (Recent)	UV Damage, Erythema	Acute skin inflammation from UV exposure; impaired barrier, increased sensitivity	RELATIVE - Wait until healed	AVOID affected areas until healed. Can apply aloe gel gently.	Wait until skin healed for waxing/exfoliation.	Safe but may be uncomfortable. Avoid further sun exposure.	Avoid affected areas. Gentle products only.
Thrombophlebitis	Superficial Vein Inflammation	Inflammation of superficial vein with possible clot formation	LOCAL - Avoid affected area	Avoid affected area completely. Can massage other areas with medical clearance.	Avoid area if in hands/feet. Otherwise safe.	Medical clearance required. Avoid exercises stressing affected area.	Safe.
Tuberculosis (Active)	TB, Mycobacterium Tuberculosis	Highly contagious airborne bacterial infection affecting lungs	TOTAL - Highly Contagious	ABSOLUTE CONTRAINDICATION. Airborne transmission risk to staff and clients. Reschedule until cleared by physician with negative sputum cultures.	ABSOLUTE CONTRAINDICATION.	ABSOLUTE CONTRAINDICATION.	ABSOLUTE CONTRAINDICATION.
Uncontrolled Blood Glucose (Diabetes)	Hyperglycemia (High), Hypoglycemia (Low)	Blood glucose >250 mg/dL with ketones OR <70 mg/dL with symptoms; immediate medical concern	TOTAL until stabilized	Do not proceed if client reports very high or very low blood sugar with symptoms. Risk of diabetic emergency during treatment. Reschedule.	Do not proceed with unstable blood sugar. Reschedule.	ABSOLUTE CONTRAINDICATION. Exercise with glucose >250 mg/dL + ketones = diabetic ketoacidosis risk. Exercise with severe hypoglycemia = seizure/coma risk. Must stabilize first.	Do not proceed if client symptomatic. Risk of medical emergency.
Uncontrolled Epilepsy	Seizure Disorder (Active), Convulsive Disorder	Frequent seizures not controlled by medication; triggers include heat, flashing lights, overexertion, stress	RELATIVE - Medical Clearance & Safety Protocols	Requires medical clearance. Avoid known triggers: flashing lights (contraindicated for LED therapy), strong scents, overstimulation. Have seizure response protocol. Remove sharp objects from treatment room. Generally relaxation beneficial if controlled.	Safe with clearance. Staff trained on seizure response. Remove tools if seizure occurs.	Medical clearance required. Avoid overexertion and overheating (seizure triggers). Avoid activities where seizure would cause injury (swimming alone, climbing). Buddy system. Emergency protocol essential.	Medical clearance advised. Avoid flashing lights/strobes. Have emergency protocol. Remove sharp implements if seizure occurs.
Varicose Veins	Spider Veins, Venous Insufficiency	Enlarged, twisted veins with impaired valve function; usually in legs	LOCAL - Avoid direct pressure	Do not massage directly over varicose veins. Can massage around area with gentle pressure. Elevate legs during treatment.	Avoid waxing over varicose veins. Massage around area only. Otherwise safe.	Exercise beneficial. Avoid prolonged standing. Wear compression garments. Swimming excellent.	Safe.
Warts (Verruca)	HPV, Human Papillomavirus	Viral infection causing raised growths; contagious through direct contact	LOCAL - Avoid area	TOTAL facial contraindication. Can massage body avoiding wart areas.	Avoid area completely. Cannot file over warts - spreads virus. Refer to physician for removal.	Cover completely. Wear shoes in shared areas. Use personal mat.	Facial warts prevent services.

MEDICAL CONTR

ENHANCED COLOR CODING SYSTEM

RED ROWS
AMBER ROWS
PURPLE ROWS
MINT ROWS

CRITICAL EVIDENCE-BASED PROTOCOLS

FOOTBATH MYCOBACTERIA PREVENTION: CDC-
ACRYLATE EXPOSURE CONTROLS: NIOSH-recom
FORMALDEHYDE MONITORING: OSHA requires
BLOODBORNE PATHOGEN COMPLIANCE: OSHA
HOT TUB LEGIONELLA PREVENTION: Water mar
ACSM EXERCISE SCREENING: Structured pre-par

FORBES FIVE-STAR OPERATIONAL STAN

1. SCREENING IS PARAMOUNT: Detailed health h
2. MEDICATIONS MATTER: Document blood thin
3. KNOW YOUR SCOPE: Beautify, don't diagnose.
4. RED FLAGS REQUIRE ACTION: Oozing lesions, s
5. INFECTION CONTROL: EPA-registered disinfect
6. STANDARD PRECAUTIONS: Treat all blood/boc
7. CHEMICAL SAFETY: SDS availability, hazard cor
8. DOCUMENTATION PROTECTS: Record conditic
9. STAFF COMPETENCY: Role-based training with
10. EMERGENCY READINESS: Protocols for chest
11. VULNERABLE POPULATIONS: Extra vigilance f
12. WHEN IN DOUBT: Medical clearance or defer

HIGHEST PRIORITY CONDITIONS BY DEF

NAILS - TOP RISKS: Onychomycosis (fungal nails), Green nail syndrome (Pseudomonas)
FACIAL/BODY - TOP RISKS: Mpox, Impetigo, MRS.

BARBER - TOP RISKS: Head lice, Tinea capitis (scalp ringworm)

*FITNESS - TOP RISKS: Cardiovascular symptoms (heart disease, high blood pressure, diabetes, etc.)
Communicable skin infections (MRSA, ringworm)*

CONTRAINDICATIONS - COMPREHENSIVE CLINICAL REFERENCE

TOTAL/ABSOLUTE CONTRAINDICATION - Service MUST be refused. Proceeding could cause serious harm to client or transmission to others. Reschedule after medical clearance/resolution.
RELATIVE/LOCAL CONTRAINDICATION - Service may proceed WITH MODIFICATIONS or requires written medical clearance. Specific areas may need avoidance. Document all modifications.
PREVENTIVE PROTOCOLS & SAFETY SYSTEMS - Critical infection control, chemical safety, or screening protocols required for safe operations. These are FACILITY RESPONSIBILITIES, not individual client conditions.
Standard information with routine precautions apply

CONTRAINDICATIONS (Purple Rows)

documented outbreak risk. Date/time-stamped cleaning logs MANDATORY.
 mended ventilation + PPE essential for nail services.
 hazard communication + air monitoring for smoothing treatments.
 standard compliance required for barber/shaving services.
 vancement program + monitoring required.
 participation screening protocol mandatory for fitness.

CONTRAINDICATIONS

history required before first treatment, updated annually
 ners, Accutane, retinoids, immunosuppressants, photosensitizers
 . Refer suspicious lesions/conditions to physicians
 spreading redness+fever, DVT symptoms, uncontrolled diabetes
 ants, proper contact time, documented cleaning protocols
 ly fluid exposure as potentially infectious
 mmunication training, proper ventilation systems
 ns, modifications, referrals, product reactions, incidents
 documented competencies, annual updates essential
 pain, syncope, seizures, severe allergic reactions, bleeding
 for immunocompromised, diabetic, pregnant, anticoagulated clients
 ral - client safety over service revenue ALWAYS

CONTRAINDICATIONS

*Paronychia (nail fold infection), Diabetic foot complications, Acrylate allergy, Footbath mycobacteria,
 A/cellulitis, Scabies, Suspected DVT, Cancer/lymphedema, Bleeding disorders*

lip ringworm), PPD dye allergy, Formaldehyde exposure (smoothing), Bloodborne pathogens (shaving)

chest pain, syncope), Uncontrolled diabetes/hypoglycemia, Acute injuries, Legionella (hot tubs),

,