

Distributing of this exam and the solution without the consent of the instructor is prohibited, as this exam is protected by copyright laws.

1. Determine whether each of the following statements is True or False. If True, prove it, if False, provide a counterexample.

- (a) Consider the probability space $(\Omega, \mathcal{F}, \mathbf{P}(\cdot))$. Let $X, Y : \Omega \rightarrow \mathbb{R}$, be such that $X + Y$ is a random variable. Then X, Y are random variables on this probability space.

Solution: : False. For any mapping X , let $Y = -X$, then $X + Y$ is random variable but we can choose the mapping X such that X, Y are not random variables.

- (b) If A_1, A_2, \dots are events such that $\mathbf{P}(A_n \text{ i.o.}) = 0$, then $\sum_{n=1}^{\infty} \mathbf{P}(A_n) < \infty$.

Solution: False. Let X be a random variable with uniform distribution over $[0, 1]$. Let $A_n = \{0 < X < \frac{1}{n}\}$. Then, we have $\sum_{n=1}^{\infty} \mathbf{P}(A_n) = \infty$ and $\mathbf{P}(A_n \text{ i.o.}) = 0$.

- (c) If \mathcal{F}_1 and \mathcal{F}_2 are both σ -algebras on a sample space Ω , then $\mathcal{F}_1 \cup \mathcal{F}_2$ is a σ -algebra on Ω .

Solution: False. $\Omega = \{1, 2, 3\}$, $\mathcal{F}_1 = \{\emptyset, \{1\}, \{2, 3\}, \Omega\}$, $\mathcal{F}_2 = \{\emptyset, \{2\}, \{1, 3\}, \Omega\}$. $\mathcal{F}_1 \cup \mathcal{F}_2$ is not a σ -algebra.

2. Let Θ be a uniformly distributed random variable over $[0, 2\pi]$. Let

$$R = \begin{cases} \cos(\Theta) & \text{if } \Theta \geq \frac{\pi}{3} \\ 0 & \text{otherwise} \end{cases}.$$

(a) Find the Cumulative Distribution Function (CDF) of R .

Solution: For $0.5 \leq r \leq 1$, we have

$$P(R > r) = P(\cos \Theta > r) = P(\Theta \in (2\pi - \arccos r, 2\pi]) = \frac{\arccos r}{2\pi}.$$

For $-1 \leq r < 0$, we have

$$\begin{aligned} P(R \leq r) &= P(\cos \Theta \leq r) \\ &= P(\cos \Theta \leq r, \Theta \in [0, \pi]) + P(\cos \Theta \leq r, \Theta \in [\pi, 2\pi]) \\ &= P(\Theta \in [\arccos r, \pi]) + P(\Theta \in [\pi, 2\pi - \arccos r]) \\ &= \frac{\pi - \arccos r}{\pi}. \end{aligned}$$

For $0 \leq r \leq 0.5$, we have

$$\begin{aligned} P(R \leq r) &= P(\cos \Theta \leq r) + P(\Theta \in [0, \pi/3]) \\ &= P(\cos \Theta \leq r, \Theta \in [0, \pi]) + P(\cos \Theta \leq r, \Theta \in [\pi, 2\pi]) + P(\Theta \in [0, \frac{\pi}{3}]) \\ &= P(\Theta \in [\arccos r, \pi]) + P(\Theta \in [\pi, 2\pi - \arccos r]) + \frac{1}{6} \\ &= \frac{\pi - \arccos r}{\pi} + \frac{1}{6}. \end{aligned}$$

Therefore, we have

$$P(R \leq r) = \begin{cases} 0 & r < -1 \\ 1 - \frac{\arccos r}{\pi} & -1 \leq r < 0 \\ 1 - \frac{\arccos r}{\pi} + \frac{1}{6} & 0 \leq r < 0.5 \\ 1 - \frac{\arccos r}{2\pi} & 0.5 \leq r < 1 \end{cases}$$

(b) Is R a continuous or a discrete random variable?

Solution: R is not continuous as the CDF is discontinuous at 0. It is not discrete as $P(R = 0) = \frac{1}{6}$ and $P(Y = \alpha) = 0$ for any other $\alpha \neq 0$.

(c) Find $\mathbb{E}[\cdot]$. **Solution:**

$$\begin{aligned} \mathbb{E}[R] &= \int_{\frac{\pi}{3}}^{2\pi} \frac{1}{2\pi} \cos \theta d\theta \\ &= \frac{1}{2\pi} [\sin \theta]_{\frac{\pi}{3}}^{2\pi} = -\frac{\sqrt{3}}{4\pi} \end{aligned}$$

3. For each of the following processes, determine whether the **partial sum sequence** $S_k = \sum_{i=1}^k X_i$ converges almost surely or not, i.e., $\lim_{k \rightarrow \infty} S_k$ exists almost surely or not.

- (a) $\{X_k\}$ is an independent process such that X_k takes three values $-k, 0, k^k$ with

$$\mathbf{P}(X_k = k^k) = \mathbf{P}(X_k = -k) = \frac{1}{2k^2}$$

and $\mathbf{P}(X_k = 0) = 1 - \frac{1}{k^2}$.

- (b) $\{X_k\}$ is given by $X_k = \frac{Y_k - 1}{k}$ where $\{Y_k\}$ is an i.i.d. Gaussian random process with $Y_k \sim \mathcal{N}(0, 1)$ (i.e., a Gaussian with zero mean and unit variance).

Solution:

- (a) It is convergence almost surely. To show this, we invoke Kolmogorov's three series theorem with $\alpha = 0.5$. Note that $Y_k = 1_{|X_k| < \alpha} X_k = 0$ surely. Therefore, we have

- i. $\sum_{k=1}^{\infty} P(|X_k| > \alpha) = \sum_{k=1}^{\infty} \frac{1}{k^2} < \infty$,
- ii. $\sum_{k=1}^{\infty} \mathbb{E}[Y_k] = 0$, and
- iii. $\sum_{k=1}^{\infty} \text{Var}[Y_k] = 0$.

Therefore, by Kolmogorov's Three Series Theorem, the sequence is convergent almost surely.

Alternatively, you could argue that $P(\{X_k \neq 0 \text{ i.o.}\}) = 0$ and hence, $\lim_{k \rightarrow \infty} X_k = 0$ almost surely.

- (b) This series is not convergent. Let $Z_k = \frac{Y_k}{k}$. Therefore, $X_k = Z_k - \frac{1}{k}$. Note that $\mathbb{E}[Z_k] = 0$ and $\text{Var}(Z_k) = \frac{1}{k^2}$. Therefore, since the process is independent, the series $\sum_{k=1}^{\infty} Z_k$ is convergent almost surely. But the sequence $-\frac{1}{k}$ is not convergent and hence, $\sum_{k=1}^{\infty} \frac{Y_k - 1}{k} = \sum_{k=1}^{\infty} Z_k - \frac{1}{k}$ is not convergent.

4. Three players P1, P2, and P3 taking turns on flipping a fair coin (starting from P1, then P2, then P3, and then P1 again, etc.). The person who gets the first head is the winner.
- (a) Let Ω be the set of all the binary sequences of the form $1, 01, 001, 0001, \dots$. Provide the set of events \mathcal{F} and a probability measure $\mathbf{P}(\cdot)$, such that the probability space $(\Omega, \mathcal{F}, \mathbf{P}(\cdot))$ models this experiment.
 - (b) Determine the event B where P2 is winning and find $\mathbf{P}(B)$.
 - (c) Let A be the event that P1 is winning. Is A and B independent?

Solution:

- (a) Let's denote a sequence of coin flip with a binary sequence where tail is represented with 0 and head with 1. It is clear that our experiment ends when we see a head. Therefore, a binary sequence $\omega_k = \underbrace{0 \dots 0}_{k-1 \text{ times}} 1$ represent the experiment that we have $k - 1$ tails and then a head at the k th flip. With the set of such sequences being Ω , then we can simply set $\mathcal{F} = \mathcal{P}(\Omega)$. Note that the probability of the event of a single outcome ω_k is exactly $P(\{\omega_k\}) = \frac{1}{2^k}$. Let us denote this with p_{ω_k} . Once we know the probability of these singleton sets, we can set $P(E) = \sum_{\omega \in E} p_{\omega}$ for any $E \in \mathcal{F}$.
- (b) Note that the winner is P2 if $k \bmod 3 = 2$, i.e.,

$$B = \{\omega_2, \omega_5, \dots\} = \{\omega_{3k+2} \mid k = 0, 1, 2, \dots\}.$$

Therefore,

$$\begin{aligned} P(B) &= \sum_{k=0}^{\infty} p_{\omega_{3k+2}} = \sum_{k=0}^{\infty} \frac{1}{2^{3k+2}} \\ &= \frac{1}{4} \sum_{k=0}^{\infty} \frac{1}{2^{3k}} \\ &= \frac{1}{4} \sum_{k=0}^{\infty} \frac{1}{8^k} \\ &= \frac{1}{4} \times \frac{1}{1 - \frac{1}{8}} = \frac{2}{7}. \end{aligned}$$

- (c) Using the same argument as above, you can show that $P(A) = 2P(B) = \frac{4}{7} \neq 0$. But $A \cap B = \emptyset$ and hence, $0 = P(A \cap B) \neq P(A)P(B)$. Therefore, the two events are not independent.