

Index

Page numbers followed by *f* indicate figure, *t* indicate table

A

Abundance. *See also* Population size

apparent, 14t

covariate relationship with, 395, 395f

estimation, 15, 384

analysis of real data, 396–409

binomial mixture model with
overdispersion in, 404–409

covariates in, 390–396

ecological process in, 392f

generation and analysis of simulated
data, 388–396

GLMM and, 383

introduction to, 383–388

observation process in, 392f

open-population binomial mixture
model in, 396–409

program R in, 396

p-values in, 397

silver-washed fritillary example of,
396–409, 397f, 408f

simple Poisson model for, 398–401

summary of, 409–410

ZIP binomial mixture model in,
401–404

false negatives and, 384

importance of, 20

as key population descriptor, 409–410

latent, 384

modeled, 14t

overdispersion in, 404–409

parameters, 387

Poisson distribution for, 385

spatial variation in, 384

in species–abundance distribution, 5

as state variable, 410, 414

study of, 1–6

true, 14t

Age effects

models with, 208–212

time effects combined with, 212

Age-dependent models, multinomial

likelihood and, 227–231, 252–255

Age-specific probability, of first breeding
model

analysis of, 290–294

description of, 288–289

generation of simulated data, 289–290

survival and movement estimation with,
288–294

Akaike's information criterion (AIC), 42,
469, 470

Algebra

BUGS language and, 50–51, 84

modeling in, 25, 267

multistate models and, 267

ANCOVA linear model

GLM, 50, 52

GLMM, 74–77, 106

R code for, 50, 52

random-effects, 75–76

ANOVA, 100–106

Apparent abundance, 14t

Apparent distribution, 14t, 415

Asp viper, 17–18, 18–19f, 74–77, 76f

Atlas, distribution, 4

B

Band-recovery models. *See* Mark-recovery
models

Batched effects, pooling of, 81

Bayes rule

information combined with, 35

learning formalized in, 34–35

paraphrased, 34

priors and, 35–37, 44

quantities in, 33

Bayesian computation, in Bayesian statistical
modeling, 38

Bayesian *p*-value

Bayesian statistical modeling

Bayesian computation in, 38

Bayesian statistical modeling (*Cont.*)

- challenges of, 41
- hierarchical, 43–44
- inference in, 44
- introduction to, 23–24, 28–38
- overview of, 464
- parameter nonidentifiability and, 217
- posterior sampling and, 41–43
- p*-values in, 222–223, 226f
- role of, 24–27
- summary of, 44–45

Begon, M., 2

Behavioral effects

- defined, 136
- in generation and analysis of simulated data, 148–150
- importance of, 138

Bernoulli distribution

- defined, 67
- site-occupancy models and, 417

Bernoulli trials, 176, 244

Beta-binomial distribution, 12

Bias

- descriptor, 7
- estimation error and, 366
- simulations and, 16–20
- systematic, 121–126

Binomial distribution

- beta-binomial, 12
- capture–recapture models and, 385
- observation process and, 9–13, 20

Binomial GLM

- analysis of real data, 70–71
- generation and analysis of simulated data, 68–70
- for modeling bounded counts or proportions, 67–71
- Poisson GLM compared to, 48
- summary of, 71–72

Binomial mixture model

- abundance estimation and, 15, 384
- analysis of real data, 396–409
- binomial mixture model with overdispersion in, 404–409
- covariates in, 390–396
- ecological process in, 392f
- generation and analysis of simulated data, 388–396
- GLMM and, 383
- introduction to, 383–388
- observation process in, 392f

- open-population binomial mixture model in, 396–409
- program R in, 396
- p*-values in, 397
- silver-washed fritillary example of, 396–409, 397f, 408f
- simple Poisson model for, 398–401
- summary of, 409–410
- ZIP binomial mixture model, 401–404

assumptions, 387–388

benefits, 385

defined, 384

detection probability in, 387

for dynamic situations, 387

false positives and, 388

Poisson-binomial, 385–386

for static situations, 387

Binomial response, 15

Biodiversity, 4–5

Black grouse, 329, 329f, 334f

Blue bug. *See* *Rosalia alpina* beetle

Brooks–Gelman–Rubin statistic, 39

BRT. *See* Bugs run time

bugs function, 87

BUGS language

- algebraic description in, 50–51, 84
- as hierarchical model translation, 25
- history of, 38
- likelihood defined in, 358
- logit stabilized in, 430
- value of, 25, 39, 41, 54–55

Bugs run time (BRT), 93

Burnin, 39

C

Capture–recapture data

- for estimating closed population size
 - analysis of real data, 157–162
 - behavioral effects in, 148–150
 - CJS model and, 135, 137–138
 - generation and analysis of simulated data, 139–157
 - individual effects in, 150–153
 - introduction to, 134–139
 - model $M_{t_{bh}}$ for, 157–162
 - model M_{t+x} for, 162–168
 - pen shell example of, 166–168, 167f
 - summary of, 169–170
 - time effects in, 145–147
 - trap response example of, 136–137, 148–157

- in integrated population model example, 357–363
- joint analysis of
 - generation of simulated data, 296–297
 - model for, 295–300
 - survival and movement estimation from, 295–300
- mark-recovery data compared to, 295
- overdispersion and, 196
- for recruitment, survival, and
 - population size estimation using JS model
 - analysis of real data, 341–345
 - assumptions of, 316
 - black grouse example of, 329, 329f, 334f
 - connections between parameters in, 339–341
 - constant survival and time-dependent entry models in, 328–335
 - data augmentation in, 319–328, 338
 - GLM and, 328, 345
 - grey-headed woodpecker example of, 335–339, 336f
 - identifiability in, 339–341
 - introduction to, 316–317
 - Leisler's bat example of, 341–345, 344f
 - models with individual capture heterogeneity, 335–339
 - parameterizations of, 317, 325–328
 - priors in, 326
 - summary of, 345–346
- for survival and movement estimation
 - using multistate models
 - age-specific probability of first breeding model, 288–294
 - fire salamander example of, 282–284, 283f
 - introduction to, 264–268
 - little ringed plover example of, 270–274, 271f, 289–290, 302–303
 - model for joint analysis of capture–recapture and mark-recovery data, 295–300
 - movement among three sites, 300–306
 - movement between two sites, 268–281
 - real-data example, 307–311
 - showy lady's slipper example, 307f, 307–311
 - state-space likelihood and, 274
 - summary of, 311–312
 - temporal effects and, 280
 - temporary emigration model, 281–287
- for survival estimation using CJS model
 - age effect models and, 208–212
 - age-dependent models and, 227–231
 - analysis of real data, 231–237
 - constant parameters and, 177–183
 - fixed group and random time effect models in, 204–208
 - fixed group and time effect models in, 199–204
 - fixed group effect models in, 192–194
 - fixed time effect models in, 184
 - GLM and, 199, 211, 237
 - individual group effects in, 195–199
 - introduction to, 172–175
 - latent state variable in, 181–183
 - Leisler's bat example, 231–237, 232f, 235–236f
 - little owl example of, 177–183, 178f, 185, 189, 196, 209, 227–231
 - models with individual variation and, 192–199
 - multinomial likelihood and, 220–231, 237
 - parameter identifiability in, 216–220
 - random group effects in, 194–195
 - random time effect models in, 184–188
 - recapture probability and, 212–216
 - red-backed shrike example, 213, 214f
 - summary of, 237–238
 - temporal covariates in, 188–192
 - temporal variability, 204
 - time and group effect models in, 199–208
 - time-dependent models and, 222–227
 - time-variation models and, 183–192
 - trap response models and, 212–216
 - understanding of, 465
- Capture–recapture models
 - advantages of, 169
 - binomial distribution and, 385
 - classical, 135–136
 - data augmentation and, 139–140, 152
 - detection probability and, 136–137, 137t
 - GLM and, 138
 - with individual covariates, 162–168

- Capture–recapture models (*Cont.*)
 - introduction to, 134
 - SECR, 135, 475
 - simulations with, 139–157
 - behavioral effects, 148–150
 - individual effects, 150–153
 - time effects, 145–147
- Chance, 27
- Cheap data, 385, 409–410
- CI. *See* Confidence interval
- CJS model. *See* Cormack–Jolly–Seber model
- Classical capture–recapture models, 135–136
- Cleaner thinking, hierarchical models for, 467
- Closed population size, capture–recapture
 - data for estimating
 - analysis of real data, 157–162
 - behavioral effects in, 148–150
 - CJS model and, 135, 137–138
 - generation and analysis of simulated data, 139–157
 - individual effects in, 150–153
 - introduction to, 134–139
 - model M_{tbh} , 157–162
 - model M_{t+X} , 162–168
 - pen shell example of, 166–168, 167f
 - summary of, 169–170
 - time effects in, 145–147
 - trap response example of, 136–137, 148–157
- Closure assumption
 - relaxing of, 475–476
 - temporary emigration and, 416
- Coal tit, 95f, 95–110, 97f
- Coefficient of variation (CV), 129
- Common terns, 244–246, 245f, 252–255
- Community
 - models, for metapopulation designs, 474–475
 - as population, 5
- Confidence interval (CI), 37
- Conservation, population analysis for, 477–478
 - restricted occupancy, 331
 - superpopulation formulation and, 333
- Conventional Poisson GLMM, for count data
 - analysis of real data, 88–90, 95–110
 - fixed site and fixed year effects, 100–102
 - fixed site effects, 99–100
 - full model, 108–110
 - generation and analysis of simulated data, 84–88, 92–95
 - introduction to, 73–82
 - overdispersion in, 83
 - with random effects for variability among groups, 90–110
 - random site and random year effects, 103–105
 - random site effect, random year effect, and first-year fixed observer effect, 105–106
 - random site effect, random year effect, first-year fixed observer effect, and overall linear time trend, 106–108
 - random site effects, 102–103
- Convergence, checking for, 62–63, 67, 70–71, 87, 89
- Cormack–Jolly–Seber (CJS) model
 - assumptions, 173–174
 - closed population size and, 135, 137–138
 - emigration and, 371
 - false negatives and, 173
 - false positives and, 173
 - in integrated population models, 371
 - JS model compared to, 237, 345
 - mark-recovery model compared to, 243–244, 248
 - m-array and, 44, 220–231
 - multinomial likelihood and, 173, 175, 220–231, 237
 - multistate model compared to, 267, 278
 - state-space models and, 173, 175–177, 181, 237, 465, 466t
 - survival estimation using
 - age effect models and, 208–212
 - age-dependent models and, 227–231
 - analysis of real data, 231–237
 - constant parameters and, 177–183
 - fixed group and random time effect models in, 204–208
 - fixed group and time effect models in, 199–204
 - fixed group effect models in, 192–194
 - fixed time effect models in, 184
 - GLM and, 199, 211, 237
 - individual group effects in, 195–199
 - introduction to, 172–175
 - latent state variable in, 181–183
 - Leisler’s bat example, 231–237, 232f, 235–236f
 - little owl example of, 177–183, 178f, 185, 189, 196, 209, 227–231
 - models with individual variation and, 192–199

- multinomial likelihood and, 220–231, 237
 - parameter identifiability in, 216–220
 - random group effects in, 194–195
 - random time effect models in, 184–188
 - recapture probability and, 212–216
 - red-backed shrike example, 213, 214f
 - summary of, 237–238
 - temporal covariates in, 188–192
 - temporal variability, 204
 - time and group effect models in, 199–208
 - time-dependent models and, 222–227
 - time-variation models and, 183–192
 - trap response models and, 212–216
- Correlation, temporal, 205
- Count data
- in integrated population model, 357–363
 - likelihood of, 352–354
 - Poisson GLMM for
 - analysis of real data, 88–90, 95–110
 - fixed site and fixed year effects, 100–102
 - fixed site effects, 99–100
 - full model, 108–110
 - generation and analysis of simulated data, 84–88, 92–95
 - introduction to, 73–82
 - overdispersion in, 83
 - with random effects for variability among groups, 90–110
 - random site and random year effects, 103–105
 - random site effect, random year effect, and first-year fixed observer effect, 105–106
 - random site effect, random year effect, first-year fixed observer effect, and overall linear time trend, 106–108
 - random site effects, 102–103
 - uses of, 111
 - replicated, 385, 409
- Covariates
- in abundance estimation, 390–396
 - abundance's relationship with, 395, 395f
 - flexible modeling of, 476
 - in individual covariate models
 - capture–recapture models, 162–168
 - for population size estimation, 166–168
 - for species richness estimation, 163–166
 - in binomial mixture model, 390–396
 - single-season occupancy model with, 422–427
 - site, 390, 386, 411, 418
 - temporal, 188–192
- Covariates, sampling, 390, 411, 418
- Covariates, survey, 386, 390, 418
- Credible interval (CRI)
- defined, 37
 - increase in, 369
- CV. *See* coefficient of variation
- ## D
- DA. *See* Data augmentation
- DAG. *See* Directed acyclic graph
- Data augmentation (DA)
- capture–recapture models and, 139, 140, 152
 - defined, 140
 - generation and analysis of simulated data with, 139–157
 - behavioral effects, 148–150
 - individual effects, 150–153
 - introduction to, 139–145
 - time effects, 145–147
 - introduction to, 139
 - JS model and, 319–328, 338
 - as multistate model, 322–325
 - as restricted dynamic occupancy model, 320–322
 - superpopulation parameterization and, 325–328
 - in random-effects models, 152
 - in recruitment, survival, and population size estimation, 319–328, 338
- Data cloning, 36
- Data dredging, 469–470
- Dead-recovery matrix, 247
- Dead-recovery models. *See* Mark-recovery models
- Demographic parameters, temporal variability of, 373
- Demographic rates
- integrated population models estimating analysis of, 358–363
 - CJS model in, 371
 - fecundity estimated with, 363–366, 377–378
 - hoopoe example of, 371f, 371–379, 378f

Demographic rates (*Cont.*)

- introduction to, 348–350
 - ortolan bunting example of, 349, 350–351f, 357–363, 366–370
 - for population viability analysis, 366–370
 - without productivity data, 363–366
 - real data example of, 371–379
 - simple example of, 357–363
 - summary of, 379–380
 - population size and, 350–352
- Detection. *See also* Detection/nondetection data
- binomial mixture model with
 - overdispersion with, 404–409
 - linear predictor for, 386
 - probability, 6, 11
 - in binomial mixture models, 387
 - capture–recapture models and, 137t, 136–137
 - date related to, 433
 - model selection view of, 473
 - state-space models and, 124–125, 131
 - time of day related to, 433
 - within-capture-history dependence of, 148
 - of silver-washed fritillary, 408f

Detection error. *See* False negatives

Detection/nondetection data

- defined, 414
- introduction to, 414–419
- occurrence and species distribution
 - estimation from
 - analysis of real data, 427–436, 445–450
 - dynamic models for, 436–450, 459
 - generation and analysis of simulated data in, 420–427, 439–445
 - goodness-of-fit and, 418
 - introduction to, 414–418
 - long-eared owl example of, 439–445, 440f, 454, 456
 - multistate occupancy models, 450–458
 - p*-values and, 419–420
 - Rosalia alpina* example of, 427–436, 428–429f, 432f, 434f
 - single-season occupancy analysis, 420–427
 - six-spot burnet example of, 445–450, 445f, 450f

summary of, 459–460

WinBUGS in, 419–436, 439–450

Deterministic mechanisms, 6

Deviance information criterion (DIC), 42, 469

Directed acyclic graph (DAG), 356f, 467

Dirichlet distribution, 300, 302, 308, 454

Dispersion, extra-Poisson, 386, 405

Distance sampling, 135, 169

Distribution. *See also* specific distributions

apparent, 14t, 415

atlas, 4

catalog, 49

importance of, 20

organism, 4

study of, 1–6

true, 14t, 415

Double-counting, 125

Dynamic multistate occupancy model, 458–459

Dynamic occupancy models, 436–450

Dynamic occurrence and species distribution models, 436–450, 459

E

Ecological process, 8, 392f

Ecology

definition of, 1–6

hierarchical view of, 6, 7f

metapopulation, 3–4

state changes in, 2, 2f

Emigration. *See also* Temporary emigration

CJS model and, 371

permanent, 295

Entry probability, 317, 323, 329, 333, 340, 345

Errors. *See also* False negatives; False

positives

estimation, 366

misclassification, 476

Monte Carlo, 144

nondetection, 7, 10

observation, 11

hierarchical models correcting, 11, 116–117, 117f, 384

Poisson distribution for, 352

Estimated population size, 121–126

Estimation error, bias and, 366

Estimator, 17, 20

E-SURGE (software), 470–471

Exchangeability, assumption of, 77, 82

Expected count, 111, 134

Explanation, objective of, 26

Explicit models
 defined, 26
 hierarchical, 43, 111, 384, 472
 of misclassification error, 476
 SECR, 135, 475
Extra-Poisson dispersion, 386, 405
Extrinsic nonidentifiability, 216

F

False negatives
 abundance and, 384
 accounting for, 476
 CJS model and, 173
 false positives canceling out, 125
 introduction to, 11–13, 20
 multistate models and, 265
 site-occupancy model and, 417
False positives
 accounting for, 476
 binomial mixture models and, 388
 CJS model and, 173
 false negatives canceling out, 125
 introduction to, 12–13, 20
 multistate models and, 265
 sample size and, 13
 site-occupancy model and, 416–417
Fecundity
 integrated population models estimating,
 363–366, 377–378
 Poisson GLM modeling, 66–67
Fidelity, 295
Finite-population standard deviation, 79
Fire salamander, 282–284, 283f
First breeding model, age-specific
 probability of
 analysis of, 290–294
 description of, 288–289
 generation of simulated data, 289–290
 survival and movement estimation with,
 288–294
First-year fixed observer effect, 105–108
Fisher Scoring, 58
Fixed effects
 additive, 154
 in fixed group and random time effect
 models, 204–208
 in fixed group and time effect models,
 199–204
 in fixed group effect models in, 192–194
 in fixed site and fixed year effects model,
 100–102
 in fixed site effects model, 99–100

 in fixed time effect models, 184
 random effects compared to, 76, 82
 in random site effect, random year effect,
 and first-year fixed observer effect
 model, 105–106
 time effects, 184, 199–204
 in WinBUGS, 76

Frailty, 208

Freeman–Tukey statistic, 224

Frequentist analysis, of statistical models,
 28–38

Freuler, Reto, 126

G

GAM. *See* Generalized additive model

Gamma prior, 120

Generalized additive model (GAM)

 defined, 71

 development of, 476

 smoothing of, 131

Generalized linear mixed model (GLMM)

 abundance estimation and, 383

 ANCOVA and, 74–77, 106

 ANOVA and, 100–106

 asp viper example, 17–18, 18–19f, 74–77,
 76f

 coal tit example of, 95f, 95–110, 97f

 complex estimation in, 95

 conventional Poisson, for count data

 analysis of real data, 88–90, 95–110

 fixed site and fixed year effects,
 100–102

 fixed site effects, 99–100

 full model, 108–110

 generation and analysis of simulated
 data, 84–88, 92–95

 introduction to, 73–82

 overdispersion in, 83

 with random effects for variability
 among groups, 90–110

 random site and random year
 effects, 103–105

 random site effect, random year
 effect, and first-year fixed observer
 effect, 105–106

 random site effect, random year
 effect, first-year fixed observer effect,
 and overall linear time trend,
 106–108

 random site effects, 102–103

 uses of, 111

 defined, 71

Generalized linear mixed model (GLMM)
(*Cont.*)

GLM compared to, 88
introduction to, 73–82
latent effects and, 73
peregrine falcon example of, 84–90, 86f,
92–95
summary of, 110–112

Generalized linear models (GLM)

advantages of, 54
ANCOVA, 50, 52
binomial
analysis of real data, 70–71
generation and analysis of simulated
data, 68–70
for modeling bounded counts or
proportions, 67–71
Poisson GLM compared to, 48
summary of, 71–72
capture–recapture models and, 138
GLMM compared to, 88
introduction to, 48
movement among three sites model
and, 301
peregrine falcon example of, 56–71,
56–57f, 65f

Poisson

analysis of real data, 64–66
binomial GLM compared to, 48
fecundity modeling by, 66–67
generation and analysis of
simulated data, 56–64
overdispersion in, 83
in R, 55–66
summary of, 71–72
random effects in, 73
in recruitment, survival, and population
size estimation, 328, 345
response components in
link function and, 54–55
noise, 48–55
signal, 48–55
summary of, 71–72
in survival estimation using CJS model,
199, 211, 237
with WinBUGS, 48
binomial GLM and, 68–71
Poisson GLM and, 55–66
time series of counts modeled with,
55–66
undefined real result trap in, 60–61

Gibbs sampling, 38

GLM. *See* Generalized linear models

GLMM. *See* Generalized linear mixed
model

Goodness-of-fit (GOF)

for integrated population models, 380
m-array and, 222, 224
in occurrence and species distribution
estimation, 418

Gregg, Kathy, 308

Grey-headed woodpecker, 335–339, 336f

Group effects, 138

fixed

in fixed group and random time
effect models, 204–208

in fixed group and time effect
models, 199–204

in fixed group effect models in,
192–194

individual, 195–199

models with, 199–208

random, 194–195

H

Hendriks, Iris, 166

Heterogeneity model, 150–153, 335–339

Hierarchical models. *See also specific models*

Bayesian statistical modeling and,
43–44

binomial mixture, 44

BUGS language translation of, 25

defined, 91

explicit, 43, 111, 384, 472

implicit, 43, 111, 384, 472

importance of, 8

observation error corrected by, 11,
116–117, 117f, 384

partitioning in, 13

power of, 464–472

for cleaner thinking, 467

for fitting of complex statistical
models, 464–465

for primary model selection,
468–469

for secondary model selection,
469–470

for step-up approach to problems,
467

for study design, 471–472

for synthetic understanding of
models, 466t, 465–467

random variables in, 43

stochastic parts of, 27

systematic parts of, 27

variable selection and, 469–470

Hierarchical scales of organization, 2
 ecology and, 6, 7f
 modeling of, 5
Home range, 135
Hoopoe, 371f, 371–379, 378f
House martin, 126–130, 126f, 130f

I

Ibex, 118–121, 118f, 121f, 124f
Identifiability, 216–220, 339–341
Immigration, 371, 377
Implicit models
 defined, 26
 hierarchical, 43, 111, 384, 472
Incidence. *See also* Occupancy
 defined, 437
 in metapopulation, 3
Independence assumption, of integrated
 population models, 355
Individual capture heterogeneity models,
 335–339
Individual covariate models
 capture–recapture, 162–168
 for population size estimation, 166–168
 for species richness estimation, 163–166
Individual effects, 136, 138
 in generation and analysis of simulated
 data with data augmentation,
 150–153
Individual variation, models with
 with fixed group effects, 192–194
 with individual group effects, 195–199
 with random group effects, 194–195
 survival estimation and, 192–199
Inference, scope of, 78–79
Information combination
 with Bayes rule, 35
 development of, 474
 as motivation for random-effects
 model, 81
Inits function, 128
Integrated population models
 capture–recapture data in, 357–363
 count data in, 357–363
 defined, 348
 demographic rates, population size, and
 projection matrices estimation using
 analysis of, 358–363
 CJS model in, 371
 fecundity estimated with, 363–366,
 377–378
 hoopoe example of, 371f, 371–379,
 378f

 introduction to, 348–350
 ortolan bunting example of, 349,
 350–351f, 357–363, 366–370
 for population viability analysis,
 366–370
 without productivity data, 363–366
 real data example of, 371–379
 simple example of, 357–363
 summary of, 379–380
development of, 349
 first step, 350–352
 second step, 352–354
 third step, 354–357
fecundity estimated by, 363–366,
 377–378
goodness-of-fit and, 380
independence assumption of, 355
for population viability analysis,
 366–370
reproduction in, 357–363
simulation with, 349
state-space likelihood and, 354
Intercepts
 random, 75, 78
 varying, 78
Intrinsic nonidentifiability, 216
Inverse Wishart distribution, 206–207

J

Joint analysis, of capture–recapture and
 mark–recovery data
 generation of simulated data, 296–297
 model for, 295–300
 survival and movement estimation from,
 295–300
Joint likelihood, formulation of, 354–357
Jolly–Seber (JS) model
 binomial mixture, 475
 CJS model compared to, 237, 345
 data augmentation and, 319–328, 338
 as multistate model, 322–325
 as restricted dynamic occupancy
 model, 320–322
 superpopulation parameterization
 with, 325–328
as multistate model, 322–325, 332
observation process in, 318, 318f
recruitment, survival, and population size
 estimation using
 analysis of real data, 341–345
 assumptions of, 316
 black grouse example of, 329, 329f,
 334f

Jolly–Seber (JS) model (*Cont.*)

- connections between parameters in, 339–341
 - data augmentation in, 319–328, 338
 - GLM and, 328, 345
 - grey-headed woodpecker example of, 335–339, 336f
 - identifiability in, 339–341
 - introduction to, 316–317
 - Leisler's bat example of, 341–345, 344f
 - models with constant survival and time-dependent entry, 328–335
 - models with individual capture heterogeneity, 335–339
 - parameterizations of, 317, 325–328
 - priors in, 326
 - summary of, 345–346
 - as restricted occupancy model, 331
 - state process in, 318, 318f
 - as state-space model, 317–319
 - superpopulation formulation and, 333
- JS model. *See* Jolly–Seber model

L

- Lack-of-fit ratio, 401
- Latent abundance, 384
- Latent effects, GLMMs and, 73
- Latent state variable, 181–183, 247, 276, 318
- Latent suitability indicators, 402
- Learning, formalized, 34–35
- Leisler's bat
 - in recruitment, survival, and population size estimation using JS, 341–345, 344f
 - in survival estimation using CJS model, 176f, 231–237, 232f, 235–236f
- Leslie-matrix modeling, 14t, 362, 379
- Levels, 78
- Likelihood. *See also* Maximum likelihood estimate; Multinomial likelihood
 - BUGS language defining, 358
 - of capture–recapture data, 354
 - of count data, 352–354
 - function, 29, 33, 36
 - joint, 354–357
 - principle, 223
 - of reproductive success data, 354
 - state-space
 - integrated population models and, 354
 - in survival and movement estimation using multistate models, 274

Linear models. *See also specific models*

- mean structure described by, 49
- themes in, 15
- Linear predictor, 51
- Link function, 54–55, 422
- Little owl, 177–183, 178f, 185, 189, 196, 209, 227–231
- Little ringed plover, 270–274, 271f, 289–290, 302–303
- Log-function, 386, 394f
- Logistic normal, 150
- Logit, 148, 154, 186, 430
- Logit-link function, 386, 394f
- Long-eared owl, 439–445, 440f, 454, 456

M

- Magden, 126–130
- Marginal probability, 33
- MARK (software), 470–471
- Markov chain Monte Carlo (MCMC)
 - benefits of, 358
 - development of, 38
 - sampling techniques, 38
 - WinBUGS blackbox of, 38–41, 44
- Mark-recovery data
 - capture–recapture data compared to, 295
 - joint analysis of
 - generation of simulated data, 296–297
 - model for, 295–300
 - survival and movement estimation from, 295–300
 - simulations with, 244–246, 296–297
 - survival estimation using
 - common terns example of, 244–246, 245f, 252–255
 - introduction to, 241–242
 - latent state variable in, 247
 - m-array for, 249, 251–253, 255
 - MCMC and, 261
 - multinomial likelihood and, 248–255, 261
 - priors in, 258, 260
 - real-data example of, 255–261
 - red kite example, 255–261, 256f, 259–260f
 - simulation of, 244–246
 - state-space models and, 242–248, 261
 - summary of, 261
- Mark-recovery models
 - CJS model compared to, 243–244, 248

- multinomial likelihood fitting, 248–255
 - model with age-dependent parameters, 252–255
 - model with constant parameters, 248–252
- as state-space model
 - advantages of, 242–248, 261
 - analysis of model with constant parameters, 246–248
 - simulation of mark-recovery data, 244–246
 - state and observation processes, 243f, 243
- M-array
 - CJS model fitted via, 44, 220–231
 - goodness-of-fit and, 222–224
 - mark-recovery data and, 249, 251–253, 255
- Maximum likelihood estimate (MLE)
 - benefits of, 31
 - from data cloning, 36
 - defined, 29
 - mode and, 28, 37
 - posterior mean and, 28
 - tadpole example of, 29, 30f, 33
- MCMC. *See* Markov chain Monte Carlo
- Memory, 160
- Metacommunity, 3t, 5
- Metapopulation, 15, 96
 - ecology, 3–4
 - extent of, 4
 - incidence in, 3
- Metapopulation designs
 - abundance estimation and, 15, 384
 - analysis of real data, 396–409
 - binomial mixture model with overdispersion in, 404–409
 - covariates in, 390–396
 - ecological process in, 392f
 - generation and analysis of simulated data, 388–396
 - GLMM and, 383
 - introduction to, 383–388
 - observation process in, 392f
 - open-population binomial mixture model in, 396–409
 - posterior distributions in, 394f, 397, 401f
 - program R in, 396
 - p*-values in, 397
 - silver-washed fritillary example of, 396–409, 397f, 408f
 - simple Poisson model for, 398–401
 - simplest case, 388–390
 - summary of, 409–410
 - ZIP binomial mixture model, 401–404
- community models for, 474–475
- in occurrence and species distribution estimation
 - analysis of real data, 427–436, 445–450
 - dynamic models for, 436–450, 459
 - generation and analysis of simulated data in, 420–427, 439–445
 - goodness-of-fit and, 418
 - introduction to, 414–418
 - long-eared owl example of, 439–445, 440f, 454, 456
 - multistate occupancy models, 450–458
 - p*-values and, 419–420
 - Rosalia alpina* example of, 427–436, 428–429f, 432f, 434f
 - single-season occupancy analysis, 420–427
 - six-spot burnet example of, 445–450, 445f, 450f
 - summary of, 459–460
 - WinBUGS in, 419–436, 439–450
- population models for, 474–475
- Misclassification error, 476
- Mixed models. *See* Generalized linear mixed model
- MLE. *See* Maximum likelihood estimate
- Mode, 28, 37
- Modeling. *See also specific types of modeling*
 - in algebra, 25, 267
 - flexible, of covariates, 476
 - of hierarchical scales of organization, 5
 - objectives, 26
 - of parameter correlations, 79–80
 - role of, 24–27
 - sayings about, 24
 - summary of, 44–45
 - synthetic understanding of, 465–467, 466t
 - themes, 15
- Modeling bounded counts or proportions
 - analysis of real data, 70–71
 - binomial GLM for, 67–71
 - generation and analysis of simulated data, 68–70
- Models. *See also specific models*
 - M_b*, 148–150

Models (*Cont.*) M_{hr} 150–153 M_{tr} 145–147 M_{tthr} 157–162 M_{thr} 154–157 M_{t+x} for population size estimation,
162–168

for species richness, 163–166

selection of, 37

detection probability and, 473

primary, 468–469

secondary, 469–470

Monneret, R.-J., 64

Monte Carlo error, 144

Movement among three sites, model of

analysis of, 304–306

description of, 300–302

generation of simulated data, 302–303

GLM and, 301

survival and movement estimation using,
300–306

Movement and survival estimation, from

capture–recapture data using

multistate models

age-specific probability of first breeding
model, 288–294fire salamander example of, 282–284,
283f

introduction to, 264–268

little ringed plover example of, 270–274,
271f, 289–290, 302–303

model for joint analysis of

capture–recapture and mark-
recovery data, 295–300

movement among three sites, 300–306

movement between two sites, 268–281

real-data example, 307–311

showy lady's slipper example, 307f,
307–311

state-space likelihood and, 274

summary of, 311–312

temporal effects and, 280

temporary emigration model, 281–287

Movement between two sites, estimated

analysis of, 274–281

described, 268–270

generation of simulated data, 270–274

using multistate models, 268–281

Movement probability, 14t

Multidimensional arrays, in WinBUGS,

396

Multievent models, 312

Multinomial likelihood

age-dependent models, 227–231, 252–255

CJS model fitted using, 173, 175, 220–231,
237

constant parameters and, 248–252

introduction to, 220–222

mark-recovery models fitted with,
248–255with age-dependent parameters,
252–255

with constant parameters, 248–252

survival estimation and, 220–231, 237,
248–255, 261

time-dependent models, 222–227

Multiseason site-occupancy models,

436–450

Multistate models

algebraic description of, 267

applications of, 265

CJS models compared to, 267, 278

as constant survival and time-dependent
entry models, 332

false negatives and, 265

false positives and, 265

JS model as, 322–325, 332

observation process and, 265, 266f

occupancy

dynamic, 458, 459

extensions of, 458, 459

importance of, 459

in occurrence and species

distribution estimation, 450–458

parameterizations of, 453

state process and, 265, 266f

in survival and movement estimation

from capture–recapture data

age-specific probability of first

breeding model, 288–294

fire salamander example of,
282–284, 283f

introduction to, 264–268

little ringed plover example of,
270–274, 271f, 289–290, 302–303

model for joint analysis of

capture–recapture and mark-
recovery data, 295–300movement among three sites,
300–306movement between two sites,
268–281

real-data example, 307–311

showy lady's slipper example, 307f, 307–311
state-space likelihood and, 274
summary of, 311–312
temporal effects and, 280
temporary emigration model, 281–287
Mutually exclusive events, 32, 32t

N

Negative binomial distribution, 12, 386
N-mixture model. *See also* Binomial mixture model
Noise
 link function and, 54–55
 as response component, 48–55
Nondetection data. *See* Detection/
 nondetection data
Nondetection error, 7, 10
Nonidentifiability, parameter, 216–217
Nonparametric assumptions, 476
Nuisance parameter, 320
Null model, 98–99

O

Observation
 equations, 116
 error
 hierarchical models correcting, 11, 116–117, 117f, 384
 Poisson distribution for, 352
 genesis of, 6–9
 process, 7f
 in abundance estimation, 392f
 accounted for, 13
 binomial distribution and, 9–13, 20
 importance of, 472–474
 in JS models, 318, 318f
 in mark-recovery models, 243f, 243
 in multistate models, 265, 266f
 population analysis and, 8
 systematic bias in, 121–126
 true state and, 8
 variance, 123
Observed population size, 121–126
Occam's razor, 24
Occupancy. *See also* Site-occupancy models
 defined, 437
 as state variable, 4
Occupancy and species distribution
 estimation, using site-occupancy models
 analysis of real data, 427–436, 445–450

dynamic models for, 436–450, 459
generation and analysis of simulated data
 in, 420–427, 439–445
goodness-of-fit and, 418
introduction to, 414–418
long-eared owl example of, 439–445, 440f, 454, 456
multistate occupancy models, 450–458
p-values and, 419–420
Rosalia alpina example of, 427–436, 428–429f, 432f, 434f
single-season occupancy analysis, 420–427
six-spot burnet example of, 445–450, 445f, 450f
summary of, 459–460
WinBUGS in, 419–436, 439–450
OpenBUGS, 390, 439
Open-population model
 binomial mixture
 in abundance estimation, 396–409
 simple Poisson model, 398–401
 understanding of, 465
Organism distribution, 4
Ortolan bunting, 349, 350–351f, 357–363, 366–370
Overdispersion
 binomial mixture model with, 404–409
 capture-recapture data subject to, 196
 correction for, 386, 404
 defined, 82
 in Poisson GLM, 83
 in Poisson GLMM, 83
 in random-effects models, 82–90
Overparameterization, 100

P

Parameterization
 effects, 52–53
 of JS recruitment, survival, and population size modeling, 317, 325–328
 means, 53
 of multistate occupancy models, 450–458
 overparameterization and, 100
 superpopulation, 325–328
 treatment contrast, 53
Parameters
 abundance, 387
 age-dependent, 252–255

Parameters (*Cont.*)

- correlations, modeling of, 79–80, 205
- defined, 20
- demographic, 373
- estimation and inference, 29
- fixed, 28–29
- identifiability, 216–220, 339–341
- nonidentifiability, 216–217
- nuisance, 320
- variance, 37

Pen shell, 166–168, 167f

Peregrine falcon

- GLM and, 56–71, 56–57f, 65f
- GLMM and, 84–90, 86f, 92–95

Permanent emigration, 295

Poisson distribution, 12

- for abundance, 385
- for observation error, 352
- zero-inflated, 386

Poisson models

- for abundance estimation, 398–401

GLM

- analysis of real data, 64–66
- binomial GLM compared to, 48
- fecundity modeling by, 66–67
- generation and analysis of simulated data, 56–64
- overdispersion in, 83
- in R, 55–66
- summary of, 71–72

GLMM

- analysis of real data, 88–90, 95–110
- fixed site and fixed year effects, 100–102
- fixed site effects, 99–100
- full model, 108–110
- generation and analysis of simulated data, 84–88, 92–95
- introduction to, 73–82
- overdispersion in, 83
- with random effects for variability among groups, 90–110
- random site and random year effects, 103–105
- random site effect, random year effect, and first-year fixed observer effect, 105–106
- random site effect, random year effect, first-year fixed observer effect, and overall linear time trend, 106–108
- random site effects, 102–103
- uses of, 111

PLN, 83–84

Poisson-binomial mixture model, 385–386

Poisson random variable, 55

Poisson response, 15

Poisson–log-normal (PLN) model, 83–84

Population(s). *See also* Integrated

- population models; Metapopulation; Open-population model; Population size; Superpopulation

abundance as key descriptor of, 409–410

analysis

- for conservation, 477–478
- defined, 6
- methods for, 15
- observation process and, 8
- recurring themes in, 477
- viability, 366–370
- for wildlife management, 477–478

classic, 4

community as, 5

counts, state-space models for

- analysis of simulated data, 119
- framework of, 117
- house martin example, 126–130, 126f, 130f
- ibex example, 118f, 118–121, 121f, 124f
- introduction to, 115–118
- MCMC and, 120, 130
- observation equations in, 116
- simple, 118–121, 131
- state-process equations in, 116
- systematic bias and, 121–126

descriptors, 7

in finite-population standard deviation, 79

likelihood of count data, 352–354

models, for metapopulation designs, 474–475

unobserved, 366

Population size. *See also* Abundance

closed, capture–recapture data for estimating

- analysis of real data, 157–162
- behavioral effects in, 148–150
- CJS model and, 135, 137–138
- generation and analysis of simulated data, 139–157
- individual effects in, 150–153
- introduction to, 134–139
- model M_{tbh} for species richness estimation, 157–162

- model M_{t+x} 162–168
- pen shell example of, 166–168, 167f
- summary of, 169–170
- time effects in, 145–147
- trap response example of, 136–137, 148–157
- demographic rates and, 350–352
- individual covariate models estimating, 166–168
- integrated population models estimating
 - analysis of, 358–363
 - CJS model in, 371
 - fecundity estimated with, 363–366, 377–378
 - hoopoe example of, 371f, 371–379, 378f
 - introduction to, 348–350
 - ortolan bunting example of, 349, 350–351f, 357–363, 366–370
 - for population viability analysis, 366–370
 - without productivity data, 363–366
 - real data example of, 371–379
 - simple example of, 357–363
 - summary of, 379–380
- JS model estimating
 - analysis of real data, 341–345
 - assumptions of, 316
 - black grouse example of, 329, 329f, 334f
 - connections between parameters in, 339–341
 - data augmentation in, 319–328, 338
 - GLM and, 328, 345
 - grey-headed woodpecker example of, 335–339, 336f
 - identifiability in, 339–341
 - introduction to, 316–317
 - Leisler's bat example of, 341–345, 344f
 - models with constant survival and time-dependent entry, 328–335
 - models with individual capture heterogeneity, 335–339
 - parameterizations of, 317, 325–328
 - summary of, 345–346
- observed, 121–126
- of occupied patches, 13
- as smoothed index, 380
- true, 121–126
- of unobserved populations, 366
- Population viability analysis, 366–370
- Positional matching, 87
- Posterior distributions
 - introduction to, 33–34, 36–37
- Posterior mean, MLE and, 28
- Posterior predictive check, 175, 222–226, 237, 388, 397–398, 400–401
- Posterior sampling, 41–43
- Precision, 16–20
- Predator control, 477
- Prediction, objective of, 26
- PRESENCE (software), 470–471
- Presence/absence data, 414, 416. *See also* Detection/nondetection data
- Primary model selection, 468–469
- Priors
 - Bayes rule and, 35–37, 44
 - Dirichlet distribution, 300, 302, 308, 454
 - gamma, 120
 - uniform, 120
 - for variance parameters, 37
- Probability
 - age-specific, of first breeding model
 - analysis of, 290–294
 - description of, 288–289
 - generation of simulated data, 289–290
 - survival and movement estimation with, 288–294
 - defined, 28
 - detection, 6, 11
 - in binomial mixture models, 387
 - capture–recapture models and, 137t, 136–137
 - model selection view of, 473
 - state-space models and, 124–125, 131
 - time of day related to, 433
 - within-capture-history dependence of, 148
 - entry, 317, 323, 329, 333, 340, 345
 - marginal, 33
 - movement, 14t
 - recapture, 212–216
 - removal entry, 320, 323, 333
 - survival, 14t, 15
- Process
 - observation, 243, 243f, 265, 266f, 318, 318f, 439
 - state, 8–13, 20, 318, 318f, 243, 243f, 265, 266f, 392f
- Process variance, 123
- Productivity data, estimation without, 363–366

Program R. *See also* R code
 distribution catalog in, 49
 simulations with, 9, 10

Projection matrices, integrated population models and
 analysis of, 358–363
 CJS model in, 371
 fecundity estimated with, 363–366, 377–378
 hoopoe example of, 371f, 371–379, 378f
 introduction to, 348–350
 ortolan bunting example of, 349, 350–351f, 357–363, 366–370
 for population viability analysis, 366–370
 without productivity data, 363–366
 real data example of, 371–379
 reproduction and, 357–363
 summary of, 379–380

Pseudo-individuals, 319, 331

Pseudoreplication
 avoidance of, 80
 in site-occupancy models, 437

p-values
 in abundance estimation, 397
 in Bayesian statistical modeling, 222–223, 226f
 criticisms of, 222–223
 in site-occupancy model, 419–420

R

R code
 for ANCOVA linear model, 50, 52
 mark-recovery data and, 249
 Poisson GLM in, 55–66
 presentation of, 16
 random-effects modeling in, 82–90
 for survival estimation using CJS model, 178
 treatment contrast parameterization and, 53

Random effects
 annual rates as, 377
 correlated, 205
 costs of, 82
 defined, 73, 77–78
 fixed effects compared to, 76, 82
 flexibility from, 15
 GLM and, 73
 group, 194–195
 in linear predictor for detection, 386
 for overdispersion correction, 404
 priors and, 77

time effects, 184–188, 204–208
 for variability among groups, 90–110
 in WinBUGS, 76, 82–90

Random factor, 78

Random processes, accounting for, 80

Random site and random year effects model, 103–105

Random site effect, random year effect, and first-year fixed observer effect model, 105–106

Random site effect, random year effect, first-year fixed observer effect, and overall linear time trend model, 106–108

Random site effects model, 102–103

Random temporary emigration, 436

Random time effect models, 184–188

Random variables, 27
 in fitting of complex statistical models, 464–465
 in hierarchical models, 43
 Poisson, 55

Random-coefficients models, 78

Random-effects models
 ANCOVA, 75–76
 data augmentation in, 152
 difficulty fitting, 191
 motivations, 78–81
 accounting for all random processes, 80
 borrowed strength, 80–81, 110
 information combination, 81
 modeling parameter correlations, 79–80
 partitioning of variability, 79
 pooling of batched effects, 81
 pseudoreplication avoidance, 80
 scope of inference, 78–79
 variability assessment, 79
 overdispersion accounted for in, 82–90
 in R, 82–90
 temporal correlation in, 205
 terminology, 78
 in WinBUGS, 82–90

Random-intercepts models, 75, 78

Range, 4

Recapture probability, survival estimation and, 212–216

Recruitment, survival, and population size estimation using JS model
 analysis of real data, 341–345

- assumptions of, 316
 - black grouse example of, 329, 329f, 334f
 - connections between parameters in, 339–341
 - data augmentation in, 319–328, 338
 - GLM and, 328, 345
 - grey-headed woodpecker example of, 335–339, 336f
 - identifiability in, 339–341
 - introduction to, 316–317
 - Leisler's bat example of, 341–345, 344f
 - models with constant survival and time-dependent entry, 328–335
 - models with individual capture heterogeneity, 335–339
 - parameterizations of, 317, 325–328
 - priors in, 326
 - summary of, 345–346
 - Recruitment probability. *See* Entry probability
 - Red kite, 255–261, 256f, 259–260f
 - Red-backed shrike, 213, 214f
 - Regression model, 74
 - Reif, Jiri, 157
 - REML method, 75
 - Removal entry probability, 320, 323, 333
 - Replicated counts, 385, 409
 - Reproduction, in integrated population model, 357–363
 - Reproductive success data, 354
 - Response components, in GLM
 - link function and, 54–55
 - noise, 48–55
 - signal, 48–55
 - Restricted occupancy model
 - dynamic, 320–322
 - JS model as, 331
 - MCMC samples for, 343
 - speed of, 341
 - Rhat statistic, 39, 62, 63f, 87
 - Robust design, 387, 472
 - Rosalia alpina* beetle, 427–436, 428–429f, 432f, 434f
- ## S
- Sample size
 - false positives and, 13
 - population quantities and, 20
 - Sampling
 - of capture–recapture data, 472
 - covariate, 390, 411, 418
 - design, 468
 - distance, 135, 169
 - Gibbs, 38
 - for mark-recovery data, 242
 - MCMC, 38
 - posterior, 41–43
 - repeated, 17
 - variation, 13, 19–20, 422
 - Schorcht, Wigbert, 231
 - Science, modeling's role in, 24–27
 - Seasons, as grouping factor, 437
 - Secondary model selection, 469–470
 - SECR models. *See* Spatially explicit capture–recapture models
 - Showy lady's slipper, 307f, 307–311
 - Shrinkage, 80, 377
 - Signal
 - link function and, 54–55
 - as response component, 48–55
 - Silver-washed fritillary, 396–409, 397f, 408f
 - Simulations
 - asp viper, 17–18, 18–19f, 74–77, 76f
 - benefits of, 16–20
 - bias and, 16–20
 - from inside out, 17
 - with program R, 9–10
 - Single-season occupancy analysis
 - model with covariates, 422–427
 - in occurrence and species distribution estimation, 420–427
 - Rosalia alpina* beetle example of, 427–436, 428–429f, 432f, 434f
 - simplest model for, 420–422
 - Site covariate, 390
 - Site effects, 90–110
 - Site-occupancy models
 - assumptions of, 416
 - Bernoulli distributions and, 417
 - defined, 414
 - dynamic, 436–450, 459
 - false negatives and, 417
 - false positives and, 416–417
 - metapopulation designs and, 474–475
 - multiseason, 436–450
 - multistate
 - dynamic, 458–459
 - extensions of, 458–459
 - importance of, 459
 - in occurrence and species distribution estimation, 450–458
 - parameterizations of, 453
 - uses of, 451

Site-occupancy models (*Cont.*)

occurrence and species distribution

estimation using

analysis of real data, 427–436,
445–450

dynamic models for, 436–450, 459

generation and analysis of simulated
data in, 420–427, 439–445

goodness-of-fit and, 418

introduction to, 414–418

long-eared owl example of, 439–445,
440f, 454, 456multistate occupancy models,
450–458*p*-values and, 419–420*Rosalia alpina* example of, 427–436,
428–429f, 432f, 434fsingle-season occupancy analysis,
420–427six-spot burnet example of, 445–450,
445f, 450f

summary of, 459–460

WinBUGS in, 419–436, 439–450

pseudoreplication in, 437

model with covariates, 422–427

in occurrence and species

distribution estimation, 420–427

Rosalia alpina beetle example of,
427–436, 428–429f, 432f, 434f

simplest model for, 420–422

for species distributions, 15, 419–420

state process of, 439

temporary emigration in, 436–437

Six-spot burnet, 445–450, 445f, 450f

Smoothed estimate, 131

Sparrows, 9–11, 11f

Spatial models

development of, 475

SECR, 135, 475

for study design, 475

Spatial variation, in abundance, 384

Spatially explicit capture–recapture (SECR)

models, 135, 475

Species distribution

estimation

analysis of real data, 427–436,
445–450

dynamic models for, 436–450, 459

generation and analysis of simulated
data in, 420–427, 439–445

goodness-of-fit and, 418

introduction to, 414–418

long-eared owl example of, 439–445,
440f, 454, 456multistate occupancy models,
450–458*p*-values and, 419–420*Rosalia alpina* example of, 427–436,
428–429f, 432f, 434fsingle-season occupancy analysis,
420–427six-spot burnet example of, 445–450,
445f, 450f

summary of, 459–460

WinBUGS in, 419–436, 439–450

site-occupancy model for, 15, 419–420

species–abundance, 5

Species richness, 139

biodiversity and, 5

model M_{tbh} for estimating, 157–162model M_{t+x} for estimating, 162–166

Species–abundance distribution, 5

Spline modeling, 476

Spotted owl, 414

Standard error, of estimator, 20

State

process

equations, 116, 265

in JS model, 318, 318f

in mark-recovery model, 243f, 243

multistate models and, 265, 266f

in site-occupancy model, 439

true, 8

variables, 6

abundance, 410, 414

latent, 181–183, 247, 276, 318

occupancy, 4

State-space likelihood

integrated population models and, 354

in survival and movement estimation

using multistate models, 274

State-space models

CJS models and, 173, 175–177, 181, 237,
465, 466t

defined, 111, 115

detection probability and, 124–125,
131

JS model as, 317–319

mark-recovery models as

advantages of, 242–248, 261

analysis of model with constant
parameters, 246–248simulation of mark-recovery data,
244–246

- state and observation processes, 243, 243f
- for population counts
 - analysis of simulated data, 119
 - framework of, 117
 - house martin example, 126–130, 126f, 130f
 - ibex example, 118f, 118–121, 121f, 124f
 - introduction to, 115–118
 - MCMC and, 120, 130
 - observation equations in, 116
 - simple, 118–121, 131
 - state-process equations in, 116
 - systematic bias and, 121–126
- priors and, 127
- summary of, 131
- survival estimation using mark-recovery data and, 242–248, 261
- Statistical models. *See also* Bayesian statistical modeling
 - complex, fitting of, 464–465
 - frequentist analysis of, 28–38
 - introduction to, 27
 - organic approach to, 469
 - response components in, 48–55
- Step-up approach, to problems, 467
- Strength, borrowing of, 80–81, 110
- Study design
 - hierarchical models and, 471–472
 - spatial models for, 475
- Suitability indicators, latent, 402
- Superpopulation
 - in constant survival and time-dependent entry models, 333
 - parameterization, 325–328
- Survey covariate, 386, 390, 418
- Survival estimation
 - using CJS model
 - age effect models and, 208–212
 - age-dependent models and, 227–231
 - analysis of real data, 231–237
 - constant parameters and, 177–183
 - fixed group and random time effect models in, 204–208
 - fixed group and time effect models in, 199–204
 - fixed group effect models in, 192–194
 - fixed time effect models in, 184
 - GLM and, 199, 211, 237
 - individual group effects in, 195–199
 - introduction to, 172–175
 - latent state variable in, 181–183
 - Leisler's bat example, 231–237, 232f, 235–236f
 - little owl example of, 177–183, 178f, 185, 189, 196, 209, 227–231
 - models with individual variation and, 192–199
 - multinomial likelihood and, 220–231, 237
 - parameter identifiability in, 216–220
 - random group effects in, 194–195
 - random time effect models in, 184–188, 204–208
 - recapture probability and, 212–216
 - red-backed shrike example, 213, 214f
 - summary of, 237–238
 - temporal covariates in, 188–192
 - temporal variability, 204
 - time and group effect models in, 199–208
 - time-dependent models and, 222–227
 - time-variation models and, 183–192
 - trap response models and, 212–216
 - using JS model
 - analysis of real data, 341–345
 - assumptions of, 316
 - black grouse example of, 329, 329f, 334f
 - connections between parameters in, 339–341
 - data augmentation in, 319–328, 338
 - GLM and, 328, 345
 - grey-headed woodpecker example of, 335–339, 336f
 - identifiability in, 339–341
 - introduction to, 316–317
 - Leisler's bat example of, 341–345, 344f
 - models with constant survival and time-dependent entry, 328–335
 - models with individual capture heterogeneity, 335–339
 - parameterizations of, 317, 325–328
 - priors in, 326
 - summary of, 345–346
 - using mark-recovery data
 - common terns example of, 244–246, 245f, 252–255
 - introduction to, 241–242
 - latent state variable in, 247
 - m-array for, 249, 251–253, 255

Survival estimation (*Cont.*)

MCMC and, 261
 multinomial likelihood and, 248–255, 261
 priors in, 258, 260
 real-data example of, 255–261
 red kite example, 255–261, 256f, 259–260f
 sampling for, 242
 simulation of, 244–246
 state-space models and, 242–248, 261
 summary of, 261

using multistate models

age-specific probability of first breeding model, 288–294
 fire salamander example of, 282–284, 283f
 introduction to, 264–268
 little ringed plover example of, 270–274, 271f, 289–290, 302–303
 model for joint analysis of capture–recapture and mark–recovery data, 295–300
 movement among three sites, 300–306
 movement between two sites, 268–281
 real-data example, 307–311
 showy lady’s slipper example, 307f, 307–311
 state-space likelihood and, 274
 summary of, 311–312
 temporal effects and, 280
 temporary emigration model, 281–287

Synthetic understanding, of models, 466t, 465–467

Systematic bias, state-space models and, 121–126

T

Tadpole example, 29, 30f, 33, 38

Temporal effects, 280

Temporal variability

of demographic parameters, 373
 in survival estimates, 204

Temporary emigration

challenge of dealing with, 135
 closure and, 416
 model
 analysis of, 284–287
 description of, 281–282

generation of simulated data, 282–284
 survival and movement estimation from, 281–287

random, 436

in site-occupancy models, 436–437

Territory mapping method, 95

tiefschutz and probefahrt error, 486

Time and group effect models, 199–208

Time effects, 136, 138

age effects combined with, 212

fixed, 184, 199–204

in generation and analysis of simulated data with data augmentation, 145–147

random, 184–188, 204–208

in time and group effect models, 199–208

Time series, of counts, 55–66

Time-variation, models with

with fixed time effects, 184
 with random time effects, 184–188
 survival estimation with, 183–192
 with temporal covariates, 188–192

Trap response

closed population size estimation and, 136–137, 148–157
 survival estimation and, 212–216

Trap-happy effect, 137, 213

Trap-shyness, 137

Treatment contrast parameterization, 53

Trend, 5

True abundance, 14t

True distribution, 14t, 415

True population size, 121–126

True state, observed state compared to, 8

U

Ultrastructural modeling, 189

Unbiased estimator, 17

Uncertainty, evaluation of, 28

Unmarked, 470–471

Unobserved populations, estimates of, 366

V

Variability

assessment, random-effects models and, 79

among groups, 90–110

partitioning of, 79

temporal, 204, 373

Variance

decomposition, 8

observation, 123
parameters, 37
process, 123
Varying-intercepts models, 78
Varying-slopes models, 78
Vital rates, 14t, 15

W

White storks, 252
Wildlife management, population analysis
for, 477–478
WinBUGS. *See also* BUGS language
Bayesian statistical modeling and,
38–40
cleaner thinking in, 467
fixed effects in, 76
flexibility of, 41
idiosyncrasies of, 40

MARK compared to, 471
as MCMC blackbox, 38–41, 44
multidimensional arrays in, 396
presentation of, 16
primary model selection in, 468–469
random effects in, 76, 82–90
secondary model selection in, 469–470
as standalone software, 40
in step-up approach, 467
Wryneck, 158f

Y

Year effects, 90–110

Z

Zero-inflated Poisson distribution, 386
Zero-inflated Poisson (ZIP) binomial
mixture model, 401–404

This page intentionally left blank