

1

2 **Strange particle in jets and underlying events with different models**

3 **1 Simulate with PYTHIA 8 sQCD with CR1 and rope**

4 **Parameters**

5 Beams:idA = 2212
6 Beams:idB = 2212
7 Main:numberOfEvents = 1001
8 Beams:eCM = 13000.
9 SoftQCD:all = on

10

11 **CR**

12 MultiPartonInteractions:pT0Ref = 2.15
13 BeamRemnants:remnantMode = 1
14 BeamRemnants:saturation = 5
15 ColourReconnection:reconnect = on
16 ColourReconnection:mode = 1
17 ColourReconnection:allowDoubleJunRem = off
18 ColourReconnection:m0 = 0.3
19 ColourReconnection:allowJunctions = on
20 ColourReconnection:junctionCorrection = 1.2
21 ColourReconnection:timeDilationMode = 2
22 ColourReconnection:timeDilationPar = 0.18

23

24 **Rope**

25 Ropewalk:RopeHadronization = on
26 Ropewalk:doShoving = on
27 Ropewalk:tInit = 1.5
28 Ropewalk:deltat = 0.05
29 Ropewalk:tShove = 0.1
30 Ropewalk:gAmplitude = 0.

31

32 Ropewalk:doFlavour = on
33 Ropewalk:r0 = 0.5
34 Ropewalk:m0 = 0.2
35 Ropewalk:beta = 0.1

36

37 ///
38 PartonVertex:setVertex = on
39 PartonVertex:protonRadius = 0.7
40 PartonVertex:emissionWidth = 0.1




Figure 1: Baryon-to-meson(top) and baryon-to-baryon ratio with PYTHIA sQCD + CR1 + rope.

2 Simulate with PYTHIA 8 sQCD with CR1 and rope




Figure 2: Track η distribution.

References




Figure 3: Integrated yields ratios of strange particle to π with $dN_{ch}/d\eta$.




Figure 4: Inclusive baryon-to-meson ratio(top) and Baryon-to-meson ratio(bottom) with p_T distribution.




Figure 5: Baryon-to-meson ratio(top) and Baryon-to-meson ratio(bottom) in jets with p_T distribution.