

1

2 **Multiplicity dependence of strange and multi-strange particle in jets in pp**  
3 **collisions at  $\sqrt{s} = 7$  TeV**

4

authors

5

**Abstract**

6

## 1 Introduction

In heavy-ion collisions at ultra-relativistic energies, it is well established that a strongly coupled Quark-Gluon-Plasma (QGP) is formed [1–5]. Recent measurements in high multiplicity pp, p–A and d–A collisions at different energies have revealed strong flow-like effects even in these small systems [6–14]. The origin of these phenomena is debated [? ? ? ? ? ? ? ].

The multi-strange baryons,  $\Omega$  (sss) and  $\Xi$  (dss), are particularly important in high energy particle and nuclear physics due to their dominant strange quark (s-quark) content. The initial state colliding projectiles contain no strange valence quark, therefore all particles with non-zero strangeness quantum number are created in the course of the collision.

## References

- [1] **STAR** Collaboration, J. Adams *et al.*, “Experimental and theoretical challenges in the search for the quark gluon plasma: The STAR Collaboration’s critical assessment of the evidence from RHIC collisions”, *Nucl. Phys. A* **757** (2005), [arXiv:nuc1-ex/0501009](#).
- [2] **PHENIX** Collaboration, K. Adcox *et al.*, “Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: Experimental evaluation by the PHENIX collaboration”, *Nucl. Phys. A* **757** (2005), [arXiv:nuc1-ex/0410003](#).
- [3] **BRAHMS** Collaboration, I. Arsene *et al.*, “Quark gluon plasma and color glass condensate at RHIC? The Perspective from the BRAHMS experiment”, *Nucl. Phys. A* **757** (2005), [arXiv:nuc1-ex/0410020](#).
- [4] **PHOBOS** Collaboration, B. B. Back *et al.*, “The PHOBOS perspective on discoveries at RHIC”, *Nucl. Phys. A* **757** (2005), [arXiv:nuc1-ex/0410022](#).
- [5] **ALICE** Collaboration, J. Schukraft, “Heavy Ion physics with the ALICE experiment at the CERN LHC”, *Phil. Trans. Roy. Soc. Lond. A* **370** (2012), [arXiv:1109.4291](#) [[hep-ex](#)].
- [6] **ALICE** Collaboration, B. Abelev *et al.*, “Transverse sphericity of primary charged particles in minimum bias proton-proton collisions at  $\sqrt{s} = 0.9, 2.76$  and 7 TeV”, *Eur. Phys. J. C* **72** (2012), [arXiv:1205.3963](#) [[hep-ex](#)].
- [7] **CMS** Collaboration, S. Chatrchyan *et al.*, “Study of the Production of Charged Pions, Kaons, and Protons in pPb Collisions at  $\sqrt{s_{NN}} = 5.02$  TeV”, *Eur. Phys. J. C* **74** no. 6, (2014), [arXiv:1307.3442](#) [[hep-ex](#)].
- [8] **CMS** Collaboration, V. Khachatryan *et al.*, “Observation of Long-Range Near-Side Angular Correlations in Proton-Proton Collisions at the LHC”, *JHEP* **09** (2010), [arXiv:1009.4122](#) [[hep-ex](#)].
- [9] **CMS** Collaboration, S. Chatrchyan *et al.*, “Observation of Long-Range Near-Side Angular Correlations in Proton-Lead Collisions at the LHC”, *Phys. Lett. B* **718** (2013), [arXiv:1210.5482](#) [[nucl-ex](#)].
- [10] **ALICE** Collaboration, B. Abelev *et al.*, “Long-range angular correlations on the near and away side in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV”, *Phys. Lett. B* **719** (2013), [arXiv:1212.2001](#) [[nucl-ex](#)].
- [11] **ATLAS** Collaboration, G. Aad *et al.*, “Observation of Associated Near-Side and Away-Side Long-Range Correlations in  $\sqrt{s_{NN}}=5.02$  TeV Proton-Lead Collisions with the ATLAS Detector”, *Phys. Rev. Lett.* **110** no. 18, (2013), [arXiv:1212.5198](#) [[hep-ex](#)].

- 48 [12] **ATLAS** Collaboration, G. Aad *et al.*, “Measurement with the ATLAS detector of multi-particle  
49 azimuthal correlations in p+Pb collisions at  $\sqrt{s_{NN}}=5.02$  TeV”, *Phys. Lett. B* **725** (2013) ,  
50 [arXiv:1303.2084 \[hep-ex\]](#) .
- 51 [13] **CMS** Collaboration, S. Chatrchyan *et al.*, “Multiplicity and Transverse Momentum Dependence  
52 of Two- and Four-Particle Correlations in pPb and PbPb Collisions”, *Phys. Lett. B* **724** (2013) ,  
53 [arXiv:1305.0609 \[nucl-ex\]](#) .
- 54 [14] **STAR** Collaboration, J. Adams *et al.*, “Identified hadron spectra at large transverse momentum in  
55 p+p and d+Au collisions at  $s(NN)^{1/2} = 200$ -GeV”, *Phys. Lett. B* **637** (2006) ,  
56 [arXiv:nucl-ex/0601033](#) .