



Clemente Bretti, Gabriele Lando, Demetrio Milea 16 September 2021

Nickel(II)

Reaction	Feitknecht et al., 1963	Baes and Mesmer, 1976	NIST46	Gamsjäger et al., 2005	Thoenen et al., 2014	Brown and Ekberg, 2016
$Ni^{2+} + H_2O = NiOH^+ + H^+$		-9.86	-9.9	-9.54 ± 0.14	-9.54 ± 0.14	-9.90 ± 0.03
$Ni^{2+} + 2 H_2O = Ni(OH)_2 + 2 H^+$		-19	-19		<-18	-21.15 ± 0.06
$Ni^{2+} + 3 H_2O = Ni(OH)_3^- + 3 H^+$		-30	-30	-29.2 ± 1.7	-29.2 ± 1.7	
$Ni^{2+} + 4 H_2O = Ni(OH)_4^{2-} + 4 H^+$		<-44				
$2 \text{ Ni}^{2+} + \text{H}_2\text{O} = \text{Ni}_2(\text{OH})^{3+} + \text{ H}^+$		-10.7		-10.6 ± 1.0	-10.6 ± 1.0	-10.6 ± 1.0
$4 \text{ Ni}^{2+} + 4 \text{ H}_2\text{O} = \text{Ni}_4(\text{OH})_4^{4+} + 4 \text{ H}^+$		-27.74	-27.7	-27.52 ± 0.15	-27.52 ± 0.15	-27.9 ± 0.6

β -Ni(OH) ₂ (s) + 2 H ⁺ = Ni ²⁺ + 2 H ₂ O		10.8			11.02 ± 0.20	10.96 ± 0.20 11.75 ± 0.13 (microcr)
$Ni(OH)_2(s) = Ni^{2+} + 2 OH^-$	-17.2 (inactive)		-17.2	$-16.97 \pm 0.20 (\beta)$ $-17.2 \pm 1.3 (cr)$		
$Ni(OH)_2(s) + OH^- = Ni(OH)_3^-$	-4.2 (inactive)					
$NiO(cr) + 2 H^{+} = Ni^{2+} + H_2O$				12.38 ± 0.06		12.48 ± 0.15

C.F. Baes and R.E. Mesmer, The Hydrolysis of Cations. Wiley, New York, 1976.

P.L. Brown and C. Ekberg, Hydrolysis of Metal Ions. Wiley, 2016, pp. 632-649.

W. Feitknecht and P. Schindler, Solubility constants of metal oxides, metal hydroxides and metal hydroxide salts in aqueous solution. Pure and Applied Chemistry, 6, 126-199 (1963).

H. Gamsjäger, J. Bugajski, T. Gajda, R.J. Lemire and W. Preis, Chemical Thermodynamics of Nickel, Chemical Thermodynamics, Volume 6, OECD, Paris, 2005.

NIST46, NIST Critically Selected Stability Constants of Metal Complexes: Version 8.0. Available at: www.nist.gov/srd/nist46

T. Thoenen, W. Hummel, U. Berner and E. Curti, The PSI/Nagra Chemical Thermodynamic Database 12/07, 2014.