

Antimony(III)

Reaction	Baes and Mesmer, 1976	Lothenbach et al., 1999; Kitamura et al., 2010	Filella and May, 2003
$\text{Sb(OH)}_3 + \text{H}^+ = \text{Sb(OH)}_2^+ + \text{H}_2\text{O}$	1.41	1.30	1.371
$\text{Sb(OH)}_3 + \text{H}_2\text{O} = \text{Sb(OH)}_4^- + \text{H}^+$	-11.82	-11.93	-11.70
$0.5 \text{ Sb}_2\text{O}_3(\text{s}) + 1.5 \text{ H}_2\text{O} = \text{Sb(OH)}_3$	-4.24		
$\text{Sb}_2\text{O}_3(\text{rhombic}, \text{s}) + 3 \text{ H}_2\text{O} = 2 \text{ Sb(OH)}_3$		-8.72	-10.00
$\text{Sb}_2\text{O}_3(\text{cubic}, \text{s}) + 3 \text{ H}_2\text{O} = 2 \text{ Sb(OH)}_3$			-11.40

C.F. Baes and R.E. Mesmer, The Hydrolysis of Cations. Wiley, New York, 1976, p. 375.

M. Filella and P.M. May, Computer simulation of the low-molecular-weight inorganic species distribution of antimony(III) and antimony(V) in natural waters. *Geochim. Cosmochim. Acta* 67, 4013-4031 (2003). doi:10.1016/S0016-7037(03)00095-4

A. Kitamura, K. Fujiwara, R. Doi, Y. Yoshida, M. Mihara, M. Terashima and M. Yui, JAEA Thermodynamic Database for Performance Assessment of Geological Disposal of High-Level Radioactive and TRU-Wastes. Report JAEA-Data/Code 2009-024, Japan Atomic Energy Agency (2010).

B. Lothenbach, M. Ochs, H. Wanner and M. Yui, Thermodynamic Data for the Speciation and Solubility of Pd, Pb, Sn, Sb, Nb and Bi in Aqueous Solution. Japan Nuclear Cycle Development Institute (JNC), TN8400 99-011 (1999).

Antimony(V)

Reaction	Baes and Mesmer, 1976	Lothenbach et al., 1999; Kitamura et al., 2010
$\text{Sb}(\text{OH})_5^0 + \text{H}_2\text{O} = \text{Sb}(\text{OH})_6^- + \text{H}^+$	-2.72	-2.72
$12 \text{ Sb}(\text{OH})_5^0 + 4 \text{ H}_2\text{O} = \text{Sb}_{12}(\text{OH})_{64}^{4-} + 4 \text{ H}^+$	20.34	20.34
$12 \text{ Sb}(\text{OH})_5^0 + 5 \text{ H}_2\text{O} = \text{Sb}_{12}(\text{OH})_{65}^{5-} + 5 \text{ H}^+$	16.72	16.72
$12 \text{ Sb}(\text{OH})_5^0 + 6 \text{ H}_2\text{O} = \text{Sb}_{12}(\text{OH})_{66}^{6-} + 6 \text{ H}^+$	11.89	11.89
$12 \text{ Sb}(\text{OH})_5^0 + 7 \text{ H}_2\text{O} = \text{Sb}_{12}(\text{OH})_{67}^{7-} + 7 \text{ H}^+$	6.07	6.07

$0.5 \text{ Sb}_2\text{O}_5(\text{s}) + 2.5 \text{ H}_2\text{O} = \text{Sb}(\text{OH})_5$	-3.7	
$\text{Sb}_2\text{O}_5(\text{am}) + 5 \text{ H}_2\text{O} = 2 \text{ Sb}(\text{OH})_5$		-7.400

C.F. Baes and R.E. Mesmer, The Hydrolysis of Cations. Wiley, New York, 1976, p. 375.

A. Kitamura, K. Fujiwara, R. Doi, Y. Yoshida, M. Mihara, M. Terashima and M. Yui, JAEA Thermodynamic Database for Performance Assessment of Geological Disposal of High-Level Radioactive and TRU-Wastes. Report JAEA-Data/Code 2009-024, Japan Atomic Energy Agency (2010).

B. Lothenbach, M. Ochs, H. Wanner and M. Yui, Thermodynamic Data for the Speciation and Solubility of Pd, Pb, Sn, Sb, Nb and Bi in Aqueous Solution. Japan Nuclear Cycle Development Institute (JNC), TN8400 99-011 (1999).