



Clemente Bretti, Gabriele Lando, Demetrio Milea, 16 September 2021

## Tin(II)

Reaction	Baes and Mesmer, 1976	Feitknecht, 1963	Hummel et al., 2002	NIST46	Cigala et al., 2012	Gamsjäger et al, 2012	Brown and Ekberg, 2016
$Sn^{2+} + H_2O = SnOH^+ + H^+$	-3.40		$-3.8 \pm 0.2$	-3.4	$-3.52 \pm 0.05$	$-3.53 \pm 0.40$	$-3.53 \pm 0.40$
$Sn^{2+} + 2 H_2O = Sn(OH)_2 + 2 H^+$	-7.06		$-7.7 \pm 0.2$	-7.1	$-6.26 \pm 0.06$	$-7.68 \pm 0.40$	$-7.68 \pm 0.40$
$Sn^{2+} + 3 H_2O = Sn(OH)_3^- + 3 H^+$	-16.61		$-17.5 \pm 0.2$	-16.6	$-16.97 \pm 0.17$	$-17.00 \pm 0.60$	$-17.56 \pm 0.40$
$2 \operatorname{Sn}^{2+} + 2 \operatorname{H}_2 O = \operatorname{Sn}_2(OH)_2^{2+} + 2 \operatorname{H}^+$	-4.77			-4.8	$-4.79 \pm 0.05$		
$3 \operatorname{Sn}^{2+} + 4 \operatorname{H}_2 O = \operatorname{Sn}_3(OH)_4^{2+} + 4 \operatorname{H}^+$	-6.88		$-5.6 \pm 1.6$	-6.88	$-5.88 \pm 0.05$	$-5.60 \pm 0.47$	$-5.60 \pm 0.47$
$Sn(OH)_2(s) = Sn^{2+} + 2 OH^{-}$				-25.8	$-26.28 \pm 0.08$		

$SnO(s) + 2 H^{+}= Sn^{2+} + H_2O$	1.76		2.5± 0.5		$1.60 \pm 0.15$
$SnO(s) + H_2O = Sn^{2+} + 2 OH^-$		-26.2			
$SnO(s) + H_2O = Sn(OH)_2$		-5.3			
$SnO(s) + 2 H_2O = Sn(OH)_3^- + H^+$		-0.9			

C.F. Baes and R.E. Mesmer, The Hydrolysis of Cations. Wiley, New York, 1976.

P.L. Brown and C. Ekberg, Hydrolysis of Metal Ions. Wiley, 2016, pp. 836-842.

R.M. Cigala, F. Crea, C. De Stefano, G. Lando, D. Milea and S. Sammartano, The inorganic speciation of tin(II) in aqueous solution. Geochim. Cosmochim. Acta, 87, 1-20 (2012).

W. Feitknecht and P. Schindler, Solubility constants of metal oxides, metal hydroxides and metal hydroxide salts in aqueous solution. Pure and Applied Chemistry, 6, 126-199 (1963).

H. Gamsjäger, T. Gajda, J. Sangster, S. K. Saxena and W. Voigt, Chemical Thermodynamics of Tin. Chemical Thermodynamics Volume 12. OECD, Paris, 2012.

W. Hummel, U. Berner, E. Curti, F.J. Pearson and T. Thoenen, Nagra / PSI Chemical Thermodynamic Data Base 01/01, July 2002.

NIST46, NIST Critically Selected Stability Constants of Metal Complexes: Version 8.0. Available at: www.nist.gov/srd/nist46





Clemente Bretti, Gabriele Lando, Demetrio Milea 16 September 2021

## Tin(IV)

Reaction	Hummel et al., 2002	Gamsjäger et al, 2012	Brown and Ekberg, 2016
$Sn^{4+} + 4 H_2O = Sn(OH)_4 + 4 H^+$			$7.53 \pm 0.12$
$Sn^{4+} + 5 H_2O = Sn(OH)_5^- + 5 H^+$			$-1.07 \pm 0.42$
$Sn^{4+} + 6 H_2O = Sn(OH)_6^{2-} + 6 H^+$			$-11.14 \pm 0.32$
$Sn(OH)_4 + H_2O = Sn(OH)_5^- + H^+$	$-8.0 \pm 0.3$	$-8.60 \pm 0.40$	
$Sn(OH)_4 + 2 H_2O = Sn(OH)_6^{2-} + 2 H^+$	$-18.4 \pm 0.3$	$-18.67 \pm 0.30$	
$SnO_2(cr) + 2 H_2O = Sn(OH)_4$	$-8.0 \pm 0.2$	$-8.06 \pm 0.11$	

$SnO_2(am) + 2 H_2O = Sn(OH)_4$	$-7.3 \pm 0.3$	$-7.22 \pm 0.08$	
$SnO_2(s) + 4 H^+ = Sn^{4+} + 2 H_2O$			$-15.59 \pm 0.04$

- P.L. Brown and C. Ekberg, Hydrolysis of Metal Ions. Wiley, 2016, pp. 836–42.
- W. Hummel, U. Berner, E. Curti, F.J. Pearson and T. Thoenen. Nagra / PSI Chemical Thermodynamic Data Base 01/01, July 2002.
- H. Gamsjäger, T. Gajda, J. Sangster, S. K. Saxena and W. Voigt. Chemical Thermodynamics of Tin. Chemical Thermodynamics Volume 12. OECD, Paris, 2012.