## Useful statistical and mathematical results

This is a collection of useful statistical and mathematical results.

## 1.1 Probability

**Definition 1.1 (Mean and Variance - Discrete random variable).** Let X be a discrete random variable with support  $x_1, ..., x_K$  and probability mass function (pmf) p(x), then

$$\mathbb{E}(X) = \sum_{k=1}^{K} x_k p(x_k). \tag{1.1}$$

and

$$\mathbb{V}(X) = \sum (x_i - \mathbb{E}(X))^2 p(x_i) dx. \tag{1.2}$$

Definition 1.2 (Mean and Variance - Continuous random variable). Let X be a continuous random variable with probabilty density function (pdf) p(x), then

$$\mathbb{E}(X) = \int x p(x) dx. \tag{1.3}$$

and

$$\mathbb{V}(X) = \int (x - \mathbb{E}(X))^2 p(x) dx. \tag{1.4}$$

**Lemma 1.3 (The law of iterated expectations).** Let X and Y be two random variables. Let  $\mathbb{E}_X$  denote the expectation with respect to the marginal distribution for X and  $\mathbb{E}_{Y|X}$  the expectation with respect to the conditional distribution of Y given X. Then

$$\mathbb{E}_{Y}(Y) = \mathbb{E}_{X} \mathbb{E}_{Y|X}(Y|X). \tag{1.5}$$

**Lemma 1.4 (Law of total variance).** Let X and Y be two random variables. Then

$$\mathbb{V}_Y(Y) = \mathbb{E}_X \mathbb{V}_{Y|X}(Y|X) + \mathbb{V}_X \mathbb{E}_{Y|X}(Y|X). \tag{1.6}$$

**Definition 1.5 (Marginal distribution - two variables).** Let X and Y be a two random variables with joint distribution p(x, y).

The marginal distribution of x when both variables are discrete is

$$p(x) = \sum_{y} p(x, y), \tag{1.7}$$

where the sum runs over all the values in the support of Y. When both variables are continuous

$$p(x) = \int p(x, y)dy. \tag{1.8}$$

**Definition 1.6 (Marginal distribution - n variables).** Let  $X_1, \ldots, X_n$  be a set of n random variables with joint distribution  $p(x_1, \ldots, x_n)$ . The marginal distribution of  $X_i$  when all variables are discrete is

$$p(x_i) = \sum_{x_1} \sum_{x_{i-1}} \cdots \sum_{x_{i+1}} \cdots \sum_{x_n} p(x_1, \dots, x_n),$$
 (1.9)

where the sums runs over all the values in each variable's support. The marginal distribution of  $X_i$  when all variables are continuous is

$$p(x_i) = \int \cdots \int p(x_1, \dots, x_n) dx_1 \dots dx_{i-1} dx_{i+1} \dots dx_n.$$
 (1.10)

**Definition 1.7 (Conditional distribution).** Let X and Y be two random variables with joint distribution p(x,y) and marginal distributions  $p_X(x)$  and  $p_Y(y)$ . Then the conditional distribution of X given Y is

$$p(x|y) = \frac{p(x,y)}{p_Y(y)}. (1.11)$$

**Lemma 1.8 (Product rule - two variables).** Let X and Y be two random variables with joint distribition p(x, y). Then

$$p(x,y) = p(y)p(x|y) = p(y|x)p(x).$$
 (1.12)

Lemma 1.9 (Product rule - n variables). Let  $X_1, \ldots, X_n$  be a set of n random variables with joint distribution  $p(x_1, \ldots, x_n)$ . Then

$$p(x_1, \dots, x_n) = p(x_1)p(x_2|x_1) \cdots p(x_n|x_1, \dots x_{n-1}).$$
(1.13)

Lemma 1.10 (Law of total probability for events). Let A and B be two events, and let  $B^c$  be the complement to B. Then

$$p(A) = p(A|B)p(B) + p(A|B^c)p(B^c). (1.14)$$

Lemma 1.11 (Bayes theorem for events). Let A and B be two events. Then

$$p(A|B) = \frac{p(B|A)p(A)}{p(B)} = \frac{p(B|A)p(A)}{p(A|B)p(B) + p(A|B^c)p(B^c)}.$$
 (1.15)

Lemma 1.12 (Bayes theorem for continuous variables). Let X denote the data and  $\theta$  a continuous parameter. Then

$$p(\theta|x) = \frac{p(x|\theta)p(\theta)}{p(x)} = \frac{p(x|\theta)p(\theta)}{\int p(x|\theta)p(\theta)d\theta}.$$
 (1.16)

## 1.2 Standard mathematical functions

Definition 1.13 (The Gamma function).

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt \tag{1.17}$$

$$\Gamma(x+1) = x\Gamma(x) \tag{1.18}$$

$$\Gamma(n) = (n-1)!$$
 if  $n$  is an integer (1.19)

$$\Gamma(1/2) = \sqrt{\pi} \tag{1.20}$$

Definition 1.14 (The Beta function).

$$B(x,y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt$$
 (1.21)

$$B(x,y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$$
 (1.22)

(1.23)