

Supplement to ‘Common or Distinct Attention Mechanisms for Contrast and Assimilation?’

Hope K. Snyder¹, Sean M. Rafferty¹, Julia M. Haaf¹, & Jeffery N. Rouder^{2,1}

¹ University of Missouri

² University of California, Irvine

Author Note

This document was written in R-Markdown with code for data analysis integrated into the text. The Markdown script is open and freely available at

<https://github.com/PerceptionAndCognitionLab/ctx-flanker/tree/public/papers/current>.

The data were *born open* (Rouder, 2016) and are freely available at

<https://github.com/PerceptionCognitionLab/data1/tree/master/ctxIndDif/flankerMorph4>

Correspondence concerning this article should be addressed to Hope K. Snyder, 205 McAlester Hall, University of Missouri, Columbia, MO 65211. E-mail:

hks7w2@mail.missouri.edu

Supplement to ‘Common or Distinct Attention Mechanisms for Contrast and Assimilation?’

This document is the supplement to “Common or Distinct Attention Mechanisms for Contrast and Assimilation?”. It provides the specification and analysis of a hierarchical Bayesian probit model for assessing the correlation across individuals’ abilities to inhibit distractors in assimilation and contrast contexts.

Model Specification

Let $Y = 0, 1$ denote whether a response is “A” or “H”, respectively. Further, let $Y_{ijk\ell m}$ denote the response for the i th participant, $i = 1, \dots, I$, in the j th context type ($j = 1, 2$ for word and letter contexts, respectively), for the k th context direction ($k = 1, 2$, for contexts that promote “A” and “H” responses, respectively), for the ℓ th target, $\ell = 1, \dots, L$, and for the m th replicate, $m = 1, \dots, M_{ijk\ell}$. The context assignments are displayed in the following table:

		k=1	k=2
		A-promoting	H-promoting
j=1	word	C_T	T_E
j=2	letter	H	A

Observations $Y_{ijk\ell m}$ are dichotomous. To model the effect of covariates in dichotomous observations, we use a probit-regression specification:

$$Y_{ijk\ell m} \stackrel{ind}{\sim} \text{Bernoulli} [\Phi(\mu_{ijk\ell})].$$

Here, Φ , the cumulative distribution function of the standard normal, is the link, and $\mu_{ijk\ell} \in (-\infty, \infty)$ is the combined effect of people, conditions, and the target.

To model individual inhibition effects, we additively decompose $\mu_{ijk\ell}$ into the effect of the target for a particular participant, $\gamma_{i\ell}$, the individual’s *assimilation* effect when the background context is a word, α_i and the individual’s *contrast* effect when the background

context is a letter frame, β_i . The decomposition is:

$$\mu_{ijkl} = \gamma_{il} + v_j x_k \alpha_i + (1 - v_j) x_k \beta_i.$$

The quantities v_j and x_k are indicators of the context type and direction, as follows:

$$v_j = \begin{cases} 0 & \text{if } j = 2 \text{ (letters)} \\ 1 & \text{if } j = 1 \text{ (word)} \end{cases} \quad x_k = \begin{cases} \frac{-1}{2} & \text{if } k = 1 \text{ (A-promoting)} \\ \frac{1}{2} & \text{if } k = 2 \text{ (H-promoting)} \end{cases}$$

Prior distributions are needed for α_i, β_i , and γ_{il} . We start with γ_{il} , the effect of the target for a particular participant. These parameters are not of primary concern, and we use a broad hierarchical prior given by: $\gamma_{il} \sim N(\nu_\ell, \delta_\ell)$ with hyper priors of $\nu_\ell \sim N(0, 1)$ and $\delta_\ell \sim \text{Inverse Gamma}(.5, .01)$.

The effects of interest are an individual's assimilation effect, α_i , and an individual's contrast effect, β_i . We follow the development in Rouder et al. (2007) who describe Bayesian analyses in these settings. The joint prior over these parameter is

$$\begin{pmatrix} \alpha_i \\ \beta_i \end{pmatrix} = N(\boldsymbol{\lambda}, \boldsymbol{\Sigma}),$$

where $\boldsymbol{\lambda}$ is the vector (μ_α, μ_β) and $\boldsymbol{\Sigma} = \begin{pmatrix} \Sigma_{1,1} & \Sigma_{1,2} \\ \Sigma_{2,1} & \Sigma_{2,2} \end{pmatrix}$ is a covariance matrix.

Following Rouder et al. (2007), the hyperpriors for μ_α and μ_β are given by

$$\mu_\alpha \sim N(0, 1),$$

$$\mu_\beta \sim N(0, 1).$$

The hyperprior for $\boldsymbol{\Sigma}$ is

$$\boldsymbol{\Sigma} \sim \text{Inverse Wishert}(3, \Omega),$$

where $\Omega = \begin{vmatrix} .05 & 0 \\ 0 & .05 \end{vmatrix}$. A key property of the above specification is that population covariance in $\boldsymbol{\Sigma}$ is an explicit parameter given by $\Sigma_{1,2}$. Hence, a population correlation is well defined, and subsequent inferences generalize to new data from new participants.

The choices we make above are all reasonable given the expected degree of variability in proportions. The standard normal prior in the probit space on μ_β and μ_α corresponds to flat priors in the space of proportions, and the shape value of 3 in the Inverse Wishert corresponds to a flat prior on the population-level correlation coefficient. The only substantive choice are the values .05 in the scale of the Inverse Wishert. This small value of precision corresponds to a vaguely informative prior on effects. Additional details are provided in Rouder et al. (2007).

Analyses and Results

The estimation of probit-transformed parameters follows from Albert and Chib (1995). Accordingly, a new set of latent variables is introduced for each observation, and we denote them w_{ijklm} . These variables are defined as follows:

$$y_{ijklm} = \begin{cases} 0 & \text{then } w_{ijklm} < 0 \\ 1 & \text{then } w_{ijklm} \geq 0 \end{cases}$$

Without loss of generality, the variable w_{ijklm} is assumed to be distributed as a normal with a variance of one and the mean of μ_{ijkl} . Thus, the parameters of interest at this level are w_{ijklm} and μ_{ijkl} . It is easy to compute the posterior of \mathbf{w} given the vector of $\boldsymbol{\mu}$ and, conversely, it is easy to compute the posterior of $\boldsymbol{\mu}$ given the \mathbf{w} 's. Hence, the marginals of each may be found by Markov chain Monte Carlo (MCMC) sampling. The conditional posterior distribution of \mathbf{w} is a truncated normal:

$$w_{ijklm} | y_{ijklm} \sim \begin{cases} N_-(\mu_{ijkl}, 1) & \text{if } y_{ijklm} = 0 \\ N_+(\mu_{ijkl}, 1) & \text{if } y_{ijklm} = 1, \end{cases}$$

where N_+ and N_- denote normals distributions truncated at zero from below and above, respectively.

Posterior distributions for remaining parameters may also be sampled with MCMC. In all cases, conditional posterior distributions of parameters may be derived directly from the

proportional form of Bayes rule (Jackman, 2009; Rouder & Lu, 2005). Priors were chosen to be conjugate, and consequently, posterior distributions may be sampled from known distributions in Gibbs steps.

Critical parameters are individual estimates of contrast and assimilation, and the population-level correlation of these abilities. Individual estimates are parameters α_i and β_i . Posterior means and posterior standard deviations of these parameters are shown as points and ellipses, respectively, in Figure 4A in the main paper. The most critical parameter is the population correlation. On each iteration of the MCMC chain, we calculated $\rho = \Sigma_{1,2} / \sqrt{\Sigma_{1,1} \times \Sigma_{2,2}}$. The prior and posterior distribution of ρ is shown in Figure 4B. A full description of the results may be found in the main paper.

References

- Albert, J. H., & Chib, S. (1995). Bayesian residual analysis for binary response regression models. *Biometrika*, *82*, 747–759.
- Jackman, S. (2009). *Bayesian analysis for the social sciences*. Chichester, United Kingdom: John Wiley & Sons.
- Rouder, J. N. (2016). The what, why, and how of born-open data. *Behavioral Research Methods*, *48*, 1062–1069. Retrieved from [10.3758/s13428-015-0630-z](https://doi.org/10.3758/s13428-015-0630-z)
- Rouder, J. N., & Lu, J. (2005). An introduction to Bayesian hierarchical models with an application in the theory of signal detection. *Psychonomic Bulletin and Review*, *12*, 573–604.
- Rouder, J. N., Lu, J., Sun, D., Speckman, P. L., Morey, R. D., & Naveh-Benjamin, M. (2007). Signal detection models with random participant and item effects. *Psychometrika*, *72*, 621–642.