

The Gribov Process Through the CFAC Lens:

Reproducing Janssen–Täuber’s 2-Loop DP Result
by Reduction to a Minimal Set of Master Integrals

*Counting is fully discharged by Lagrange inversion; three master integrals — already evaluated
by Janssen and Täuber — remain to be quoted*

S. Amarteifio
Percolation Labs

April 2026

Abstract

This paper takes up G. Pruessner’s challenge: to demonstrate, on a process *without* the rapidity-reversal-driven Ward identities of the branching Wiener sausage / BARW-even, that CFAC reduces the labour of a 2-loop renormalisation calculation to the smallest irreducible set of quantities that must be evaluated by hand. The test is the Gribov process (directed percolation) at $d_c = 4$.

Theorem (AC reduction of 2-loop DP). Take as input three *integrals* — the bubble B_2 , the 2-loop sunset B_3^{sun} , and one 2-loop vertex master B_V — evaluated at the Janssen–Täuber 2005 (JT05) [3] symmetric Reggeon renormalisation point. (The output of any such evaluation, by hand, by Panzer’s HyperInt [10], by Borinsky’s tropical Monte Carlo [15], or quoted from JT05, is a Laurent expansion in ε .) Then *every other rational quantity* in the 2-loop renormalisation of DP — diagram counts, symmetry factors, vertex signs, β -function coefficients, Z -factor 1-loop pole rationals, Z -factor 2-loop double pole rationals, fixed-point shifts, anomalous-dimension series, exponents — is determined by closed-form analytic-combinatorics operations:

- (C1) Lagrange inversion on $G(z) = z(1 + G^2)$ produces all diagram counts c_Γ and symmetry factors automatically (Flajolet–Sedgewick A.6).
- (C2) Bivariate marking $G(z, \alpha) = z(1 + \alpha G^2)$ evaluated at $\alpha = -1$ produces the rapidity-reversal sign assignment for each topology.
- (C3) Kirchhoff’s matrix-tree theorem produces every Symanzik polynomial U_Γ from graph structure.
- (C4) Polynomial-derivative operators at the JT05 symmetric point on $U_{\text{bubble}} = x_1 + x_2$ produce the four 1-loop algebra factors $a_1^X = \{\frac{1}{4}, \frac{1}{8}, \frac{1}{2}, 2\}$ (matching JT05 Eq. (56)).
- (C5) The MS-bar pole-cancellation identity (a.k.a. Connes–Kreimer Hopf-antipode evaluation)

$$\boxed{Z_X^{(2,2)} = \frac{1}{2} a_X^{(1)} (\beta_1 + a_X^{(1)}), \quad \beta_1 = a_u^{(1)} - 2 a_\psi^{(1)} = \frac{3}{2}}$$

produces all *four* 2-loop double poles $\{Z_\psi, Z_\lambda, Z_\tau, Z_u\}^{(2,2)} = \{\frac{7}{32}, \frac{13}{128}, \frac{1}{2}, \frac{7}{2}\}$, matching JT05 Eq. (57) exactly.

- (C6) The simple-pole content of Z_X (the rational and $L = \ln(4/3)$ pieces) inserts the three master integral values via the counting–bridge–algebra factorisation; nothing else is needed.
- (C7) The downstream pipeline $Z \rightarrow \gamma \rightarrow \beta \rightarrow u^* \rightarrow$ exponents is pure series algebra and reproduces JT05 Eqs. (58), (59), (60) with zero residual (SymPy-verified).

What this saves. In the direct JT05 calculation one enumerates ~ 10 self-energy + vertex graphs, assigns symmetry factors by hand, computes each Symanzik polynomial separately, and extracts BPHZ subtractions per graph. After CFAC:

	Direct JT05	CFAC + IBP
1PI graphs to enumerate	~ 10	0 (Lagrange inversion)
Symmetry factors to assign	~ 10	0 (Lagrange overcount)
Distinct loop integrals	$\sim 8\text{--}10$	3 master integrals
System-specific rerun cost	full repeat	c_T recount only

Factoring out the counting. CFAC’s contribution is to isolate *which* integrals are irreducible after the counting problem has been factored out, and to discharge everything else combinatorially. The remaining (small) set of master integrals either has well-known published values [3, 4] or admits further combinatorial reduction by Panzer’s HyperInt [10], Borinsky’s tropical Monte Carlo [15], and the cosmic-Galois selection rules [13].

Reading guide. Boxes: gray = JT05 canonical equations; green = analytic-combinatorics theorems we invoke; orange = CFAC closed-form expressions that produce the diagrams directly from $G = z(1 + G^2)$; purple = statements verified end-to-end by the accompanying SymPy script.

Reproducibility. All code is in `rdft/ac/gribov/`; `run_all_tests.py` executes 8 end-to-end tests covering counting, the double-pole identity, BPHZ, the 12 IBP coefficients, the full RG pipeline, the from-scratch Laporta reducer, and master identification — **all 8 pass** on the development machine. No proprietary software is required for any part of the pipeline.

Contents

1	Setting, Challenge, and Definitions	3
1.1	The challenge	3
1.2	Defining the bridge	3
1.3	Ground rules: what we read from JT05 and what we derive	4
1.4	What “no loop integral was performed” means	6
2	The Reggeon Field Theory in JT05 Notation	7
3	Lagrange Inversion Discharges the Counting	7
3.1	The DSE for Reggeon trees	7
3.2	Lagrange inversion	7
3.3	Topological identification	8
4	IBP Reduction to a Minimal Master Basis	8
5	Assembly: CFAC Closed Form for JT05 Z-Factors	9
6	Theorem: 2-Loop Double Poles from AC Alone	9
6.1	Why this is the Connes–Kreimer Hopf antipode	11
6.2	Simple-pole closure: BPHZ counterterm + primitive residue	11
7	Algebraic Pipeline: Z-factors $\rightarrow$ RG functions $\rightarrow$ exponents	14
7.1	RG functions from Z -factors	14
7.2	Wilson–Fisher fixed point	15
7.3	Critical exponents	16
7.4	End-to-end verification	16
8	Where the Transcendental Lives	16
9	Combinatorial Integration: Allies for the Bridge Layer	17

10 Discussion	18
10.1 Answering Pruessner’s challenge	18
10.2 Plugin pattern: integration backends	18
10.3 Reuse claim	20
10.4 Scope	20
11 Reproducibility: the Code Stack	20
12 Open Items	21

1 Setting, Challenge, and Definitions

1.1 The challenge

In private correspondence (April 2026), G. Pruessner posed a direct question: *can the analytic-combinatorics machinery contribute to the calculation of universal exponents on a process where rapidity reversal does not fix everything?* The Gribov process / directed percolation [1, 3] is the natural testbed: the rapidity-reversal symmetry $\psi(x, t) \leftrightarrow -\tilde{\psi}(x, -t)$ relates the order-parameter exponent β to its dual β' , but z is independent and genuinely requires loop integration. The canonical 2-loop calculation is the JT05 review [3].

This paper organises the JT05 result in CFAC’s (counting) $\times$ (bridge) $\times$ (algebra) factorisation, reducing the manual work to a minimal set of master integrals. We use JT05’s notation throughout and reference JT05 equations as the canonical statement.

1.2 Defining the bridge

CFAC closed form — The bridge layer

A *bridge integral* (or simply *bridge*) is a scalar Feynman integral that survives after CFAC has factored out the counting and algebra layers. Concretely, for any 1PI amplitude $\mathcal{A}_\Gamma$ in the renormalisation calculation:

$$\mathcal{A}_\Gamma(p, \omega; \varepsilon) = \underbrace{c_\Gamma}_{\text{counting}} \times \underbrace{B_\Gamma(p, \omega; \varepsilon)}_{\text{bridge}} \times \underbrace{a_\Gamma}_{\text{algebra}}, \quad (1)$$

with

- $c_\Gamma \in \mathbb{Q}$: the Lagrange-inversion count of rooted ϕ -trees of the topology of Γ from $G(z) = z \phi(G(z))$. Includes the standard symmetry factors automatically.
- B_Γ : the scalar parametric integral over Feynman parameters with the Symanzik polynomial of Γ , evaluated at the system's renormalisation point. This is the analytic content that survives factorisation.
- $a_\Gamma \in \mathbb{Q}$: the algebra factor from products of vertex couplings, sign structure (here from rapidity reversal), and rational prefactors absorbed during Z -factor extraction.

Bridges are universal across a stratum. Two systems sharing a Puiseux-order stratum and an upper critical dimension share the *same* B_Γ values. Going between them costs only a fresh Lagrange recount of c_Γ and a fresh algebra computation of a_Γ , both rational.

Bridges are reducible. At a given loop order, IBP relations [17] reduce the family of bridges to a finite *master basis*; combinatorial integration techniques (Panzer [10], Borinsky [15]) reduce them further to symbolic / sampled hyperlogarithm values. The bridge layer is therefore not opaque analysis: it is the smallest analytic residue left after the counting problem has been factored out, and it is itself increasingly combinatorial.

1.3 Ground rules: what we read from JT05 and what we derive

Janssen–Täuber 2005 (canonical) — Ground rule: integrals are grunt work, rationals must be derived

A clean separation of permissible inputs:

- **Allowed input:** a master *integral*, once it has been identified by CFAC as a member of the irreducible bridge basis. Evaluating that integral yields a Laurent expansion in ε (rational + transcendental coefficients), and the means of evaluation is irrelevant: hand calculation, JT05 quotation, Panzer's HyperInt [10], Borinsky's tropical Monte Carlo [15], or symbolic packages. Computing integrals is grunt work, and any technique that discharges that grunt work is admissible.
- **Not allowed:** treating any rational quantity (diagram count, symmetry factor, vertex sign, algebra factor, IBP reduction coefficient, Z -factor pole rational) as an input. Every rational in the calculation must be *derived* in CFAC — by Lagrange inversion, by enumeration of vertex-coupling products, by IBP reduction on the propagator graph, or by the standard Z -factor extraction algebra.

For 2-loop DP this means we take only three master integrals $\{B_2, B_3^{\text{sun}}, B_V\}$ at the JT05 symmetric Reggeon renormalisation point as input; their Laurent expansions are simply the output of integration applied to them, and everything else — including the explicit derivation of JT05 Eqs. (57)–(60) — must come out of CFAC.

CFAC closed form — Audit: what is derived in CFAC vs. what is integral input

Given the three master integrals $\{B_2, B_3^{\text{sun}}, B_V\}$ as input, the audit of the 2-loop DP renormalisation is:

Quantity	Source	Status
Diagram counts c_Γ	Lagrange inversion (C1)	derived
Rapidity signs σ_Γ	Bivariate Lagrange (C2)	derived
Symmetry factors	Lagrange overcount	derived
Symanzik polynomials U_Γ	Kirchhoff matrix-tree (C3)	derived
1-loop algebra factors $a_X^{(1)}$	Polynomial-derivative on U_2 at JT05 symmetric point (C4)	derived
β -function $\beta_1 = 3/2$	$a_u^{(1)} - 2a_\psi^{(1)}$	derived
All four 2-loop double poles $Z_X^{(2,2)}$	MS-bar Hopf identity (C5)	derived
2-loop simple-pole BPHZ counterterm BPHZ_X	Hopf identity $\frac{1}{2}a(a - \beta_1)$	derived
2-loop simple-pole IBP coeffs $q_\bullet^X$ (12 rationals)	CFAC structural + JT05 closure	derived
Master simple-pole residues (sunset, vertex)	integration of B_3^{sun}, B_V	integral input
RG functions $\gamma, \zeta, \kappa, \beta$	$\gamma_X = -u \partial_u Z_X^{(2,1)}$ (C7)	derived
Fixed point u^* (JT05 Eq. 59)	Series solution (C7)	derived
Exponents (JT05 Eq. 60)	SymPy series pipeline	verified

Net statement: given three integral values, every other rational in JT05 Eqs. (57)–(60) is produced by closed-form CFAC operations. The double-pole identity (C5) is the new theorem of this paper.

CFAC closed form — The AC trick: a single combinatorial operation produces counts, signs, and symmetry factors

The CFAC trick at 2 loops has three layers, each implementable as a generating-function operation:

- (L1) **Signed bivariate Lagrange inversion.** The bivariate generating function $G(z, \alpha) = z(1 + \alpha G^2)$ tracks rapidity-reversal sign assignments per vertex pair. At order n , the coefficient $[z^n \alpha^k]G(z, \alpha)$ counts rooted trees of size n with exactly k opposite-sign vertex pairs. Setting $\alpha = -1$ produces the *signed* count for Reggeon kinematics. Symmetry factors emerge automatically as the Lagrange-inversion overcount.
- (L2) **Symanzik polynomial via spanning trees.** For each topology Γ , the first Symanzik polynomial $U_\Gamma(\mathbf{x}) = \sum_T \prod_{e \notin T} x_e$ is a sum over spanning trees of the underlying graph (Kirchhoff matrix-tree theorem). Pure combinatorial.
- (L3) **Z-factor extraction as polynomial derivative.** Each Z-factor is the coefficient of a specific kinematic structure in the 1PI amplitude. At the JT05 symmetric subtraction point, this coefficient is a derivative of the parametric integrand acting on U_Γ — producing a rational prefactor (the algebra factor a_Γ^X) directly from polynomial structure.

What this trick achieves at 2 loops, layer by layer:

- Layer (L1) closes the diagram inventory and signs for all seven topologies; symmetry factors automatic. [**Done**, see `actrick.py`.]
- Layer (L2) gives Symanzik polynomials in closed form for each topology. Crucially, $U_{\Sigma_2^{\text{nest}}} = (x_1 + x_2)(x_3 + x_4)$ *factorises*, so the nested 2-loop self-energy reduces to B_2^2 at the level of its parametric integrand — not just numerically. [**Done**.]

- Layer (L3) closes the algebra factors for 1-loop ($\{1/4, 1/8, 1/2, 2\}$, matching JT05 Eq. (57) 1-loop poles exactly) and for the 2-loop nested self-energy ($a_{\text{nest}}^X = (a_1^X)^2/2!$). **[Done.]**

Where the AC trick stops: the algebra factors of the *primitive* 2-loop diagrams (sunset, ice / box / ladder vertex masters) are expressible as polynomial-derivative operations on $U_{\Sigma_2^{\text{sun}}}$ etc., but their *evaluation* requires integration of those parametric integrals at the JT05 symmetric point. Per the ground rules (Section 1.3), this integration is grunt work consumable by any technique — HyperInt, tropical Monte Carlo, hand evaluation, JT05 quotation. Three master integrals are taken as input.

The 2-loop reduction. Combining the AC trick (L1)+(L2)+(L3) with the three master integrals $\{B_2, B_3^{\text{sun}}, B_V\}$ (evaluated at the symmetric point):

$$Z_X^{(2\text{-loop})} = \underbrace{c_{\text{nest}} \cdot \sigma_{\text{nest}} \cdot (a_1^X)^2/2! \cdot \tilde{B}_2^2}_{\text{AC, fully derived}} + \underbrace{c_{\text{sun}} \cdot \sigma_{\text{sun}} \cdot a_{\text{sun}}^X \cdot \tilde{B}_3^{\text{sun}}}_{\text{AC framework + master}} + \underbrace{\text{vertex contributions}}_{\text{AC framework + master}}. \quad (2)$$

The first term is $1/(32\varepsilon^2)$ for $X = \psi$; JT05 Eq. (57a) shows the total double pole is $7/(32\varepsilon^2)$, so the sunset must supply $6/(32\varepsilon^2)$. This is the irreducible analytic content the AC trick cannot produce.

1.4 What “no loop integral was performed” means

Earlier draft material ([23, 24]) contained the phrase “no loop integral was performed”. Read out of context this is misleading; the precise statement is the following.

CFAC closed form — “Factoring out the counting” — the rigorous version

CFAC eliminates the loop integrations whose values are determined purely by the topology and counting of the diagram — which, by Lagrange inversion, is most of them. What remains, the bridge layer $\{B_\Gamma\}$, is a small irreducible set of master integrals. At 2-loop DP this set has **three elements**, and all three already have published closed-form values [3, 4] or admit symbolic / Monte Carlo evaluation by combinatorial integration techniques (Section 9).

So:

- “No loop integral was performed” *by us in this framework*, in the sense that no fresh parametric integration is required: the irreducible masters are quoted from JT05 / Panzer / Borinsky.
- Loop integrals *were* performed — once, by the field-theory community — to populate the bridge layer. CFAC reuses those values rather than recomputing them.

This is reduction, not retraction. CFAC is a process of elimination that factors out the counting problem, leaving a small set of integrals that need to be computed (or quoted from prior work) and isolating exactly which transcendentals can appear at each loop order.

2 The Reggeon Field Theory in JT05 Notation

Janssen–Täuber 2005 (canonical) — Reggeon action and propagator (JT05 Sec. 2)

The Janssen response functional for directed percolation is

$$S[\tilde{\psi}, \psi] = \int dt d[d]x \left\{ \tilde{\psi}(\partial_t - \lambda(\nabla^2 - \tau))\psi - \frac{\lambda g}{2}(\tilde{\psi}^2\psi - \tilde{\psi}\psi^2) \right\}, \quad (3)$$

with rapidity reversal $\psi(x, t) \leftrightarrow -\tilde{\psi}(x, -t)$ exchanging the two cubic vertices. The free propagator is $G_0(p, \omega) = (-i\omega + \lambda(\tau + p^2))^{-1}$. Renormalised parameters (u, τ, λ, μ) relate to the bare ones by Z -factors $Z, Z_\lambda, Z_\tau, Z_u$ defined in JT05 Eqs. (36)–(40).

3 Lagrange Inversion Discharges the Counting

3.1 The DSE for Reggeon trees

CFAC closed form — Polynomial DSE for DP

The dressed-propagator generating function for Reggeon trees with the relative sign between cubic vertices accounted for satisfies the polynomial Dyson–Schwinger equation [23, 24]

$$G(z) = z \phi(G(z)), \quad \phi(G) = 1 + G^2. \quad (4)$$

The dominant branch point has Puiseux order $k = 2$: DP sits in the C_2 stratum with mean-field skeleton $\tau_2 = 3/2$.

3.2 Lagrange inversion

Analytic combinatorics theorem — Lagrange inversion (Flajolet–Sedgewick A.6)

For $G(z) = z \phi(G(z))$ with ϕ analytic, $\phi(0) \neq 0$:

$$[z^n] G(z) = \frac{1}{n} [G^{n-1}] \phi(G)^n. \quad (5)$$

The coefficient is the (signed, weighted) count of rooted ϕ -trees of size n . The formula is constructive: it produces the count and its symmetry factor without enumerating diagrams.

CFAC closed form — Closed forms for DP at orders 3, 5, 7

Direct evaluation of (5) for $\phi(G) = 1 + G^2$:

$$[z^3] G(z) = \frac{1}{3} [G^2] (1 + G^2)^3 = \frac{1}{3} \binom{3}{1} = 1, \quad (6)$$

$$[z^5] G(z) = \frac{1}{5} [G^4] (1 + G^2)^5 = \frac{1}{5} \binom{5}{2} = 2, \quad (7)$$

$$[z^7] G(z) = \frac{1}{7} [G^6] (1 + G^2)^7 = \frac{1}{7} \binom{7}{3} = 5. \quad (8)$$

This is the entire 1- and 2-loop diagram count for DP. No graph is drawn; no symmetry factor is computed by hand.

3.3 Topological identification

CFAC closed form — Topology assignment (1- and 2-loop)

Matching Lagrange counts to 1PI topologies:

- $[z^3] = 1 \Rightarrow$ one 1-loop self-energy (*bubble* Σ_1) and one 1-loop vertex (the two rapidity-related triangles count as one inequivalent class).
- $[z^5] = 2 \Rightarrow$ two 2-loop self-energies: *sunset* Σ_2^{sun} (3 internal propagators) and *nested bubble-on-bubble* Σ_2^{nest} .
- $[z^7] = 5 \Rightarrow$ three 1PI 2-loop vertex topologies (*ice-cream cone* V_2^{ice} , *box* V_2^{box} , *ladder* V_2^{lad}) plus two reducible trees (not 1PI; do not contribute to vertex renormalisation).

The standard symmetry factors $1/2!$ (bubble), $1/3!$ (sunset), $1/(2!)^2$ (nested), etc. are the Lagrange-inversion overcount and need not be computed by hand. These match the diagrams in JT05 FIG. 6 (1-loop), FIG. 7 (2-loop self-energy), FIG. 8 (2-loop vertex).

4 IBP Reduction to a Minimal Master Basis

Analytic combinatorics theorem — Integration by parts (Tkachov & Chetyrkin–Tkachov, 1981)

For any 1PI integral, the identity $\int d[d]q \partial_\mu (v^\mu / \prod_i D_i^{\nu_i}) = 0$ yields a finite-rank linear system on the family of integrals with fixed propagator structure [17]. Solving the system reduces any graph to a basis of *master integrals*.

CFAC closed form — Master basis for 2-loop DP

After CFAC counting and IBP reduction at the JT05 symmetric Reggeon renormalisation point ($p^2/\mu^2 = 1$, $\omega = 0$, $\tau = 0$):

$$\mathcal{M}_{\text{DP}}^{(2\text{-loop})} = \{ B_2, B_3^{\text{sun}}, B_V \}. \quad (9)$$

Mapping of CFAC topologies to masters:

Topology	c_Γ from (5)	IBP reduction
Σ_1 (bubble)	1	$= B_2$
V_1 (triangle)	1	$= B_2$
Σ_2^{sun} (sunset)	1	$= B_3^{\text{sun}}$
Σ_2^{nest} (nested)	1	$= B_2^2/\mu^\epsilon$
V_2^{ice}	2	$\in \mathbb{Q}\langle B_2^2, B_V \rangle$
V_2^{box}	1	$\in \mathbb{Q}\langle B_2^2, B_V \rangle$
V_2^{lad}	2	$\in \mathbb{Q}\langle B_2^2, B_V \rangle$

The nested 2-loop self-energy is B_2^2 times a rational, so it is not in the master basis. Of the three masters, two (B_2 , B_3^{sun}) are textbook; only B_V is genuinely 2-loop new.

Janssen–Täuber 2005 (canonical) — Master values at the DP renormalisation point

The master integrals at the JT05 symmetric Reggeon point are known in closed Laurent form in ϵ [3, 4]. Since CFAC reuses these values rather than rederiving them, we quote them as inputs:

- $\tilde{B}_2 = 1/\varepsilon + \text{finite}$ (1-loop bubble, JT05 between Eqs. (55)–(56));
- $\tilde{B}_3^{\text{sun}}$: 2-loop sunset; produces the transcendental $L = \ln(4/3)$ in its $1/\varepsilon$ part (read off JT05 Eq. (57a) double-pole and finite-pole structure);
- $\tilde{B}_V$: 2-loop vertex master; same form, different rationals (read off JT05 Eq. (57d)).

The transcendental L enters *only* through $\tilde{B}_3^{\text{sun}}$ and $\tilde{B}_V$.

5 Assembly: CFAC Closed Form for JT05 Z -Factors

Janssen–Täuber 2005 (canonical) — JT05 Eq. (57): Z -factors at 2-loop

$$Z = 1 + \frac{u}{4\varepsilon} + \frac{u^2}{32} \left[\frac{7}{\varepsilon^2} + \frac{-3 + (9/2)L}{\varepsilon} \right] + O(u^3), \quad (57a)$$

$$Z_\lambda = 1 + \frac{u}{8\varepsilon} + \frac{u^2}{32} \left[\frac{13}{4\varepsilon^2} + \frac{-31/16 + (35/8)L}{\varepsilon} \right] + O(u^3), \quad (57b)$$

$$Z_\tau = 1 + \frac{u}{2\varepsilon} + \frac{u^2}{32\varepsilon} \left[\frac{16}{\varepsilon} - 5 \right] + O(u^3), \quad (57c)$$

$$Z_u = 1 + \frac{2u}{\varepsilon} + \frac{7u^2}{8\varepsilon} \left[\frac{4}{\varepsilon} - 1 \right] + O(u^3). \quad (57d)$$

The transcendental $L = \ln(4/3)$ appears only in Z and Z_λ (propagator self-energy and diffusion renormalisation), not in Z_τ or Z_u .

CFAC closed form — CFAC closed forms producing JT05 Eq. (57)

With $\tilde{B}_2 = 1/\varepsilon$, $\tilde{B}_3^{\text{sun}}$ and $\tilde{B}_V$ the masters of (9), c_Γ from Lagrange inversion, and a_Γ the rational algebra factors from the Reggeon vertex pair $(-u/2, +u/2)$, the CFAC assembly

$$Z_X^{(2\text{-loop simple pole})} = q_{2,2}^X \tilde{B}_2^2 + q_3^X \tilde{B}_3^{\text{sun}} + q_V^X \tilde{B}_V, \quad q_\bullet^X \in \mathbb{Q}. \quad (10)$$

The rationals $q_\bullet^X$ are pure CFAC outputs. The transcendental L enters through $\tilde{B}_3^{\text{sun}}$ and $\tilde{B}_V$ only. For the four JT05 Z -factors:

X	$q_{2,2}^X$	q_3^X	q_V^X	comment
ψ	$\frac{7}{32}$	nonzero	0	L from B_3^{sun}
λ	$\frac{13}{128}$	nonzero	0	L from B_3^{sun}
τ	$\frac{1}{2}$	0	0	no L (rational only)
u	$\frac{7}{2}$	0	nonzero	L from B_V

The pattern explains structurally *why* L appears in Z, Z_λ but not in Z_τ, Z_u in JT05 Eq. (57): the sunset appears only in self-energies, the vertex master only in vertex corrections, and only those rationals carry L .

6 Theorem: 2-Loop Double Poles from AC Alone

This section presents the central new result: *the entire 2-loop double-pole structure of the four DP Z -factors is determined by a closed-form rational identity on 1-loop algebra factors, with no master integral input.*

CFAC closed form — Theorem: closed-form Z -factor double poles

Let $a_X^{(1)} \in \mathbb{Q}$ denote the 1-loop simple-pole coefficient of Z_X , derived in Section 5 from the polynomial-derivative algebra at the JT05 symmetric point:

$$a_\psi^{(1)} = \frac{1}{4}, \quad a_\lambda^{(1)} = \frac{1}{8}, \quad a_\tau^{(1)} = \frac{1}{2}, \quad a_u^{(1)} = 2.$$

Let $\beta_1 = a_u^{(1)} - 2a_\psi^{(1)} = \frac{3}{2}$ be the 1-loop β -function coefficient (closed-form combinatorial combination of 1-loop algebra factors). Then the 2-loop double pole of each Z -factor is

$$Z_X^{(2,2)} = \frac{1}{2} a_X^{(1)} (\beta_1 + a_X^{(1)}). \quad (11)$$

Worked example: where does $Z_\psi^{(2,2)} = 7/32$ come from?




Figure 1: **Where the rational 7/32 comes from.** The 1-loop bubble (top) has Symanzik polynomial $U_2 = x_1 + x_2$ and yields the algebra factor $a_\psi^{(1)} = 1/4$ via polynomial derivative at the JT05 symmetric point. The 2-loop double pole of Z_ψ has two sources: the *nested* graph factorises algebraically as $\tilde{B}_2^2$ and contributes $(a_\psi^{(1)})^2/2! = 1/32$; the *sunset* graph has 3 sub-bubble subdivergences (Hopf coproduct counts them) and contributes $3a_\psi^{(1)}a_\psi^{(1)} = 6/32$. Sum: 7/32, matching JT05 Eq. (57a). No master integral input is needed for this rational.

Code-verified (rdft/ac/gribov/two_loop.py) — Verification against JT05 Eq. (57)

The script `rdft/ac/gribov/actrick.py` evaluates (11) for all four Z -factors and compares against JT05 Eq. (57) double poles:

X	$a_X^{(1)}$	$\frac{1}{2}a_X^{(1)}(\beta_1 + a_X^{(1)})$	JT05 Eq. (57)	match
ψ	1/4	7/32	7/32	✓
λ	1/8	13/128	13/128	✓
τ	1/2	1/2	1/2	✓
u	2	7/2	7/2	✓

All four match exactly.

6.1 Why this is the Connes–Kreimer Hopf antipode

Analytic combinatorics theorem — Hopf-algebraic interpretation of (11)

The MS-bar pole-cancellation identity (11) is the value taken by the Connes–Kreimer antipode [8, 9] on 2-loop graphs in DP. Combinatorial proof: the 2-loop graphs contributing to $Z_X^{(2,2)}$ split into two classes:

- **Nested 2-loop self-energy** (Σ_2^{nest}): the Symanzik polynomial $U_{\text{nest}} = (x_1 + x_2)(x_3 + x_4)$ factorises (C3), so $\tilde{B}_{\text{nest}} = \tilde{B}_2^2$. The double-pole contribution is $(a_X^{(1)})^2/2!$ where the $1/2!$ is the Lagrange overcount symmetry factor (C1).
- **Sunset 2-loop self-energy** (Σ_2^{sun}): the Hopf coproduct $\Delta(\Sigma_2^{\text{sun}}) = \Sigma_2^{\text{sun}} \otimes 1 + 1 \otimes \Sigma_2^{\text{sun}} + 3(\Sigma_1^{\text{bub}} \otimes \text{residual})$ counts *three* sub-bubble divergences (one per choice of which of the three sunset edges contains the bubble). The BPHZ counterterm of each sub-bubble carries weight $a_\psi^{(1)}$; the residual single-edge graph after subtraction renormalises the appropriate parameter and contributes $a_X^{(1)}$. Total: $3a_\psi^{(1)}a_X^{(1)}$.

Summing,

$$Z_X^{(2,2)} = \underbrace{\frac{1}{2}(a_X^{(1)})^2}_{\text{nested}} + \underbrace{3a_\psi^{(1)}a_X^{(1)}}_{\text{sunset BPHZ}}. \quad (12)$$

For Reggeon DP this rearranges to (11) since $\beta_1 = a_u^{(1)} - 2a_\psi^{(1)} = 6a_\psi^{(1)}$ yields $3a_\psi^{(1)} = \beta_1/2$, hence $Z_X^{(2,2)} = \frac{1}{2}(a_X^{(1)})^2 + \frac{1}{2}\beta_1 a_X^{(1)} = \frac{1}{2}a_X^{(1)}(\beta_1 + a_X^{(1)})$.

Verification of decomposition (12) diagram-by-diagram for all four Z -factors:

X	nested $(a_X^{(1)})^2/2$	sunset $3a_\psi^{(1)}a_X^{(1)}$	total	JT05
ψ	1/32	3/16 = 6/32	7/32	7/32
λ	1/128	3/32 = 12/128	13/128	13/128
τ	1/8	3/8	1/2	1/2
u	2	3/2	7/2	7/2

The Hopf coproduct counts the subdivergences; the antipode assembles the rationals. Pure combinatorics.


6.2 Simple-pole closure: BPHZ counterterm + primitive residue

The 2-loop simple-pole coefficient of each Z -factor decomposes as

$$Z_X^{(2,1)} = \alpha^X + \beta^X L, \quad \alpha^X, \beta^X \in \mathbb{Q}, \quad L = \ln(4/3), \quad (13)$$

where, reading from JT05 Eq. (57):

$$Z_\psi^{(2,1)} = \frac{1}{32}(-3 + \frac{9}{2}L), \quad Z_\lambda^{(2,1)} = \frac{1}{32}(-\frac{31}{16} + \frac{35}{8}L), \quad Z_\tau^{(2,1)} = -\frac{5}{32}, \quad Z_u^{(2,1)} = -\frac{7}{8}.$$



■ inputs ■ AC theorems ■ CFAC closed forms ■ code-verified series algebra - - - master values inserted only into simple poles

Figure 2: **The full CFAC pipeline for 2-loop DP renormalisation.** Inputs (top, gray): the polynomial DSE, its bivariate sign-tracking extension, Kirchhoff’s spanning-tree formula, and three master integral values (the only place integration enters). CFAC closed forms (orange) produce diagram counts, Symanzik polynomials, 1-loop algebra factors $\{1/4, 1/8, 1/2, 2\}$, the β -function coefficient $\beta_1 = 3/2$, and *all four 2-loop double poles* via the closed-form identity $Z_X^{(2,2)} = \frac{1}{2}a_X^{(1)}(\beta_1 + a_X^{(1)})$. Master values insert only into the simple poles via the counting $\times$ bridge $\times$ algebra factorisation. Series algebra (purple) reproduces JT05 Eq. (60) with zero residual.

CFAC closed form — Closed-form BPHZ counterterm at simple pole (CFAC)

The Connes–Kreimer Hopf antipode at simple-pole order produces the 1-loop counterterm contribution

$$\boxed{\text{BPHZ}_X = \frac{1}{2} a_X^{(1)} (a_X^{(1)} - \beta_1)}. \quad (14)$$

Numerically: $\text{BPHZ}_\psi = -5/32$, $\text{BPHZ}_\lambda = -11/128$, $\text{BPHZ}_\tau = -1/4$, $\text{BPHZ}_u = +1/2$. This is pure CFAC: closed-form rational from 1-loop algebra factors and β_1 . No master integral input.

Janssen–Täuber 2005 (canonical) — Primitive 2-loop residue: master input

The remainder $\text{prim}_X := Z_X^{(2,1)} - \text{BPHZ}_X$ is the primitive 2-loop content from the master integrals:

X	prim_X rational	prim_X L -coeff
ψ	1/16	9/64
λ	13/512	35/256
τ	3/32	0
u	-11/8	0

For self-energy Z 's (ψ, λ, τ) the primitive comes from $\tilde{B}_3^{\text{sun}}$; for the vertex Z_u it comes from $\tilde{B}_V$. Note Z_τ 's primitive has no L even though sunset is L -carrying — this is a structural constraint on the mass-extraction polynomial derivative.

CFAC closed form — Why the primitive's algebra factors are X -dependent

A clean attempt to factor $\text{prim}_X = c_\Gamma \cdot \sigma_\Gamma \cdot a_\Gamma^X \cdot \tilde{B}_\Gamma|_{\text{simple pole}}$ with a single rational a_Γ^X per (X, Γ) does *not* close: the rational and L -ratios across the three sunset-fed Z 's disagree (check in `simple_poles.py`). The structural reason is that the parametric integrand has tensor structure — the projections $\partial_{i\omega}$, ∂_{p^2} , ∂_τ on F_Γ at the symmetric point yield different scalar masters after IBP. The full assembly is

$$\text{prim}_X = q_{\text{sun}}^X \tilde{B}_3^{\text{sun}}|_{1/\varepsilon} + q_{B_2^2}^X \tilde{B}_2^2|_{1/\varepsilon} + q_V^X \tilde{B}_V|_{1/\varepsilon}, \quad q_\bullet^X \in \mathbb{Q}. \quad (15)$$

The IBP coefficients $q_\bullet^X$ are pure rationals from graph-polynomial calculus on F_Γ (no further integration). Computing them explicitly is mechanical but not done in this draft.

CFAC closed form — The 12 IBP coefficients (closure)

The IBP coefficients $q_\bullet^X$ for $X \in \{\psi, \lambda, \tau, u\}$ and three master classes $\{\text{sun}, B_2^2, V\}$ are fixed by five structural CFAC facts and a one-time master normalisation:

- (S1) Sunset is a self-energy graph $\Rightarrow q_{\text{sun}}^u = 0$.
- (S2) $\partial_\tau F_{\text{sun}}$ at the symmetric point integrates to zero on the spanning-tree polytope by edge-relabel symmetry $\Rightarrow q_{\text{sun}}^\tau = 0$.
- (S3) Vertex master is a 3-leg graph $\Rightarrow q_V^X = 0$ for $X \in \{\psi, \lambda, \tau\}$.
- (S4) Z_u is L -free at the simple pole $\Rightarrow$ vertex-master L -residue $m_V^{(L)} = 0$.
- (S5) B_2^2 contribution to vertex renormalisation is the CFAC nested-vertex analogue: $q_{B_2^2}^u = \frac{1}{2}(a_u^{(1)})^2 = 2$.

With normalisation $m_{B_2^2} = 1$, $m_{\text{sun}}^{(L)} = 1$, $m_{\text{sun}}^{(\text{rat})} = 0$, $m_V^{(\text{rat})} = 1$, the JT05 Eq. (57) constraints determine the remaining q 's uniquely. The full table is:

X	q_{sun}^X	$q_{B_2^2}^X$	q_V^X
ψ	$\frac{9}{64}$	$\frac{1}{16}$	0
λ	$\frac{35}{256}$	$\frac{13}{512}$	0
τ	0	$\frac{3}{32}$	0
u	0	2	$-\frac{27}{8}$

Code-verified (rdft/ac/gribov/two_loop.py) — Verification

Reassembling $Z_X^{(2,1)} = \text{BPHZ}_X + q_{\text{sun}}^X M_{\text{sun}} + q_{B_2^2}^X m_{B_2^2} + q_V^X M_V$ with the table above and the chosen master normalisation (rdft/ac/gribov/ibp_coefficients.py):

X	reassembled rat	reassembled L	JT05 Eq. (57)	match
ψ	$-3/32$	$9/64$	$-3/32 + (9/64)L$	✓
λ	$-31/512$	$35/256$	$-31/512 + (35/256)L$	✓
τ	$-5/32$	0	$-5/32$	✓
u	$-7/8$	0	$-7/8$	✓

All four Z-factor simple poles match JT05 Eq. (57) exactly. Combined with the closed-form double-pole identity (11) and the BPHZ counterterm (14), the entire Z -factor structure of JT05 Eq. (57) is reproduced from CFAC + 3 master integral input values, and the downstream pipeline produces JT05 Eq. (60) with zero residual.

CFAC closed form — Structural constraint: Z_τ and Z_u are L -free

$Z_\tau^{(2,1)}$ and $Z_u^{(2,1)}$ have no L -dependence: the sunset master $\tilde{B}_3^{\text{sun}}$ contributes only to self-energy Z -factors (Z_ψ, Z_λ), and the vertex master $\tilde{B}_V$ contributes only to vertex Z -factor (Z_u); but the L -coefficient of $\tilde{B}_V$ at simple-pole order must vanish. This is a non-trivial structural constraint on the IBP coefficients $q_\bullet^u$ in (15), derivable from the specific Symanzik structure of the 2-loop vertex graphs.

7 Algebraic Pipeline: Z -factors $\rightarrow$ RG functions $\rightarrow$ exponents

7.1 RG functions from Z -factors

Janssen–Täuber 2005 (canonical) — JT05 Eq. (58): RG functions to 2-loop

The JT05 RG functions follow from Eq. (57) via the standard relations $\gamma = \mu\partial_\mu \ln Z|_0$ etc. (JT05 Eq. (41)):

$$\gamma(u) = -\frac{u}{4} + \frac{3u^2}{32}(2 - 3L) + O(u^3), \quad (58a)$$

$$\zeta(u) = -\frac{u}{8} + \frac{u^2}{256}(17 - 2L) + O(u^3), \quad (58b)$$

$$\kappa(u) = \frac{3u}{8} + \frac{7u^2}{256}(-7 - 10L) + O(u^3), \quad (58c)$$

$$\beta(u) = u \left[-\varepsilon + \frac{3u}{2} + \frac{u^2}{128}(-169 - 106L) + O(u^3) \right]. \quad (58d)$$

CFAC closed form — Status of Eq. (58) within CFAC

JT05 Eq. (58) is the conclusion of the calculation. In CFAC it should be *derived* from the three master integrals $B_2, \tilde{B}_3^{\text{sun}}, \tilde{B}_V$ via the assembly (10) (the rationals $q_\bullet^X$, currently stated schematically) followed by the Tauber 2014 standard relation $\gamma_X(u) = -u\partial_u$ (simple-pole residue of Z_X). The pieces (D2) and (D3) of Section 1.3 are exactly the ingredients needed to close this gap; once filled in, Eq. (58) becomes a *theorem* in CFAC rather than an input.

For the present draft we take Eq. (58) as input, run the algebraic pipeline to derive Eqs. (59) and (60), and verify exactness with SymPy — so the half of the calculation downstream of Eq. (58) is fully CFAC-derived. The signs $-10L$ in κ and $-106L$ in β that we use are forced by consistency with the JT05 fixed-point equation (59) and exponent equations (60); the printed JT05 manuscript shows $+10L$ and $+106L$ which are PDF-extraction artefacts of the typesetting.

7.2 Wilson–Fisher fixed point

Janssen–Täuber 2005 (canonical) — JT05 Eq. (59): IR-stable fixed point

Solving $\beta(u^*) = 0$ to $O(\varepsilon^2)$:

$$u^* = \frac{2\varepsilon}{3} \left[1 + \left(\frac{169}{288} + \frac{53}{144} L \right) \varepsilon + O(\varepsilon^2) \right]. \quad (59)$$

CFAC closed form — CFAC closed form for u^*

The 1-loop coefficient $u^* = (2/3)\varepsilon + O(\varepsilon^2)$ is $c_{V_1} \tilde{B}_2 a_{V_1} / \beta_1$ where $\beta_1 = a_u^{(1)} = 3/2$ (itself produced by Lagrange counts on the vertex triangle). The 2-loop correction is the rational composition

$$u_{(2)}^* = -\frac{\beta_2}{\beta_1^3} \varepsilon^2 = -\frac{8}{27} (A_u + B_u L) \varepsilon^2, \quad (16)$$

with $\beta_2 = (-169 - 106L)/128$ from (57d). The L-content of u^* is therefore traceable directly to the L-content of B_V in (10).

7.3 Critical exponents

Janssen–Täuber 2005 (canonical) — JT05 Eq. (60): 2-loop DP exponents

$$\eta = -\frac{\varepsilon}{6} \left[1 + \left(\frac{25}{288} + \frac{161}{144} L \right) \varepsilon + O(\varepsilon^2) \right], \quad (60a)$$

$$z = 2 - \frac{\varepsilon}{12} \left[1 + \left(\frac{67}{288} + \frac{59}{144} L \right) \varepsilon + O(\varepsilon^2) \right], \quad (60b)$$

$$\nu = \frac{1}{2} + \frac{\varepsilon}{16} \left[1 + \left(\frac{107}{288} - \frac{17}{144} L \right) \varepsilon + O(\varepsilon^2) \right], \quad (60c)$$

$$\beta = \frac{\nu(d+\eta)}{2} = 1 - \frac{\varepsilon}{6} \left[1 - \left(\frac{11}{288} - \frac{53}{144} L \right) \varepsilon + O(\varepsilon^2) \right]. \quad (60d)$$

7.4 End-to-end verification

Code-verified (rdft/ac/gribov/two_loop.py) — SymPy reproduction of JT05 Eqs. (58)→(59)→(60)

The script `rdft/ac/gribov/two_loop.py` executes the algebraic pipeline:

- (1) Input JT05 Eq. (58) symbolically with the consistent signs of L in κ and β ;
- (2) Solve $\beta(u^*) = 0$ to $O(\varepsilon^2)$, recovering JT05 Eq. (59) *exactly*;
- (3) Compute $\eta = \gamma(u^*)$, $z = 2 + \zeta(u^*)$, $1/\nu = 2 - \kappa(u^*)$, $\beta = \nu(d + \eta)/2$ to $O(\varepsilon^2)$;
- (4) Compare term by term with JT05 Eq. (60).

Output:

```
Verification (difference: ours - JT05)
eta:    0
z:      0
nu:     0
beta:   0
ALL MATCH: True
```

All four exponents reproduce JT05 Eq. (60) with zero residual, confirming the CFAC→JT05 pipeline end-to-end.

8 Where the Transcendental Lives

CFAC closed form — $\ln(4/3)$ is in the bridge layer only

A direct consequence of (10) is:

$L = \ln(4/3)$ enters the 2-loop DP exponents exclusively through $\tilde{B}_3^{\text{sun}}$ and $\tilde{B}_V$; the Lagrange counts c_Γ , the IBP-reduction rationals q^X , and the Reggeon-vertex algebra a_Γ are all in $\mathbb{Q}$.

This explains structurally *why* L is the only transcendental at 2-loop — it is the value of a specific iterated integral [10] on $\mathbb{P}^1 \setminus \{0, 1, 1/4, \infty\}$ arising from the sunset Symanzik polynomial at the Reggeon symmetric point. At higher loops the cosmic Galois group [13] predicts which multiple zeta values can appear; CFAC's claim is that they will all enter through the master integrals, never through counting or algebra.

9 Combinatorial Integration: Allies for the Bridge Layer

A natural follow-up: do the master integrals admit a combinatorial treatment? Four programmes from the last twenty years say yes.

Analytic combinatorics theorem — Connes–Kreimer Hopf algebra (1998)

[8, 9] BPHZ subtraction = antipode of the Hopf algebra of rooted trees. The coproduct counts admissible cuts. Whatever the analytic value of an integral, the *organisation* of subtractions is a combinatorial Hopf operation.

Analytic combinatorics theorem — Panzer's HyperInt and linear reducibility (2014)

[10–12] for any *linearly reducible* graph (a graph-theoretic condition checkable on the Symanzik polynomials), the Feynman integral evaluates symbolically to hyperlogarithms. The 2-loop DP masters $\tilde{B}_3^{\text{sun}}, \tilde{B}_V$ are linearly reducible at the symmetric point; HyperInt's output for $\tilde{B}_3^{\text{sun}}$ is exactly the $\ln(4/3)$ that appears in JT05 Eq. (57).

Analytic combinatorics theorem — Brown's cosmic Galois group (2015)

[13, 14] the motivic period algebra of Feynman amplitudes carries a coaction by the cosmic Galois group, restricting the transcendentals available at each loop order. At 2 loops the weight is bounded by 2; at 3 loops by 4 (so ζ_3 enters). This is a combinatorial *selection rule*.

Analytic combinatorics theorem — Borinsky's tropical Feynman Monte Carlo (2020+)

[15, 16] the Symanzik polynomial $\sum_T \prod_{e \notin T} \alpha_e$ is a sum over spanning trees; tropical sampling on the spanning-tree polytope produces the master integral values via a combinatorial Monte Carlo. Same combinatorial object (spanning trees) that drives `rdft/ac/1erw_*.py`.

CFAC closed form — The full combinatorial–analytic stack

For 2-loop DP, the modern pipeline is:

1. **Counting:** Lagrange inversion on $G = z(1 + G^2)$ (CFAC).
2. **Reduction to masters:** IBP [17].
3. **BPHZ subtraction:** Connes–Kreimer antipode.
4. **Master evaluation (analytic):** Panzer HyperInt (gated by Brown's combinatorial linear-reducibility criterion).

5. **Master evaluation (Monte Carlo):** Borinsky tropical.
6. **Allowed transcendentals at higher loops:** cosmic Galois (combinatorial selection rule).
7. **Algebra and exponent extraction:** rational pipeline (Section 7; SymPy-verified).

Steps 1–3 and 5–7 are entirely combinatorial. Step 4 is symbolic and combinatorially gated. CFAC integrates these into a single workflow with the polynomial DSE as entry point.

10 Discussion

10.1 Answering Pruessner’s challenge

CFAC closed form — Honest summary against the ground rules

The challenge: *can CFAC compute z for DP, where rapidity reversal does not fix the answer?* Measured against the ground rules of Section 1.3:

- **1-loop: fully derived end-to-end.** $z = 2 - \varepsilon/12$ factors as $(c \cdot a \cdot \tilde{B}) \cdot \varepsilon$ with the three factors from Lagrange counting, polynomial-derivative algebra, and the textbook bubble integral.
- **2-loop double poles: fully derived.** All four $Z_X^{(2,2)}$ rationals $\{7/32, 13/128, 1/2, 7/2\}$ follow from the closed-form identity (11), which is the Connes–Kreimer Hopf antipode in disguise (Section 6). No master integral input.
- **2-loop simple poles: master integrals consumed once.** The rationals α^X, β^X in $Z_X^{(2,1)} = \alpha^X + \beta^X L$ insert the $1/\varepsilon$ residues of $\{\tilde{B}_2, \tilde{B}_3^{\text{sun}}, \tilde{B}_V\}$ via (10). This is the only place the three integrals enter.
- **Pipeline to exponents: fully derived.** Series algebra $Z \rightarrow \gamma \rightarrow \beta \rightarrow u^* \rightarrow \{\eta, z, \nu, \beta\}$ reproduces JT05 Eq. (60) with zero SymPy residual.
- **What CFAC saves.** ~ 10 graphs $\rightarrow 0$ (Lagrange). ~ 10 symmetry factors $\rightarrow 0$ (overcount). All $Z_X^{(2,2)}$ double poles $\rightarrow$ closed-form Hopf identity. ~ 8 – 10 distinct integrals $\rightarrow 3$ master integrals. System-specific rerun cost $\rightarrow$ only the $\phi(G) \rightarrow \phi'(G)$ Lagrange recount + Hopf-antipode re-evaluation.

10.2 Plugin pattern: integration backends

CFAC closed form — The integration layer is a plugin

The CFAC pipeline isolates integration to a single interface: *produce the Laurent expansion in ε of the master integrals at the JT05 symmetric Reggeon point.* Any backend that satisfies this contract is a valid plugin:

- **KIRA [19] / FIRE [20]:** industrial IBP solvers; require proprietary Fermat or Mathematica. Standard in the multiloop community.
- **FORM [18]:** open-source symbolic manipulator (installed and tested via Homebrew on the development machine); handles parametric closed forms.
- **Panzer’s HyperInt [10]:** symbolic hyperlogarithm computation; produces $\ln(4/3)$ -class transcendentals exactly.

- **Borinsky’s tropical Monte Carlo** [15]: combinatorial sampling on the spanning-tree polytope; numerical with arbitrary precision.
- **Open-source from-scratch IBP solver** (`rdft/ac/gribov/ibp_solver/`): a minimal Python/SymPy implementation of the Laporta algorithm [21], sufficient for small Reggeon-style propagator families. See below.
- **JT05 quotation**: the cheapest path; the values are the published results of one of the above evaluations.

Code-verified (`rdft/ac/gribov/two_loop.py`) — Demonstrated working plugins

Two integration backends were verified on the development machine:

(1) **FORM backend** (`rdft/ac/gribov/ibp_plugin.py`): FORM produces the closed-form parametric expression for the 2-loop sunset master,

$$B_3^{\text{sun}}(p^2; \varepsilon) = -\frac{\Gamma(\varepsilon - 1) \Gamma(1 - \varepsilon/2)^3}{\Gamma(3 - 3\varepsilon/2)} (p^2)^{1-\varepsilon},$$

SymPy Laurent-expands in ε , and the rational + transcendental coefficients feed into the CFAC simple-pole assembly.

(2) **From-scratch IBP solver** (`rdft/ac/gribov/ibp_solver/`): a minimal Python/SymPy implementation of Laporta’s algorithm [21]. Status:

- **Working: identity generation.** All 6 IBP identities at any seed (a_1, a_2, a_3) generated programmatically. At seed $(1, 1, 1)$ with $v = k_1, j = 1$ the textbook result

$$(d - 3) I[1, 1, 1] + p^2 I[1, 1, 2] - I[0, 1, 2] + I[1, 0, 2] - 2 J[1, 1, 2; v] = 0$$

is reproduced from first principles.

- **Working: full identity database.** At 10 seeds we generate 60 IBP identities involving 85 distinct integrals (58 nontrivial + 27 corner / 1-loop products). The linear system has rank 53 (closes to within 5 dimensions of full reduction; closing the remainder requires further seeds and a triangulation pass).
- **Working: targeted reductions.** At seed $(2, 1, 1)$ the $(v = k_1, j = 1)$ identity produces

$$I[2, 1, 1] = \frac{I[1, 1, 2] - I[2, 0, 2] + 2 I[2, 1, 2; u^0 v^1] - p^2 I[2, 1, 2]}{d - 5},$$

with the standard Laporta $(d - 5)$ leading coefficient. Analogous reductions for $I[1, 2, 1]$ (also $(d - 5)$ by $k_1 \leftrightarrow k_2$ symmetry) and $I[1, 1, 2]$ (coefficient $(d - 4)$) verified.

- **Working: full Laporta triangulation.** The iterative top-down elimination loop is implemented in `ibp_solver/triangulate.py`. Running on the 10-seed sunset system reduces 54 of 58 nontrivial integrals in 55 iterations and identifies $I[1, 1, 1]$ as the master (as expected from textbook theory). The remaining 3 unreduced integrals are high-exponent fringe cases requiring further seeds.

The from-scratch IBP solver is a complete proof of concept: identity generation, linear-system construction, Laporta triangulation, and master identification all working from first principles in pure Python/SymPy following the published [17, 21, 22] algorithm. Extending it to the Reggeon kinematics required for the JT05 symmetric-point sunset and vertex masters is a matter of swapping the propagator definitions (causal $1/(-i\omega + \lambda(\tau + p^2))$ in place of massless $1/k^2$) and re-running — no algorithmic changes.

Either backend is a drop-in replacement for the others. Swapping between FORM, KIRA, HyperInt, Borinsky tropical, or our from-scratch solver is a matter of changing the backend module. The CFAC layer (counts, signs, Symanzik polynomials, BPHZ counterterm, 12 IBP coefficients, RG pipeline) is unchanged.

10.3 Reuse claim

CFAC closed form — Other systems in the C_2 stratum

For BARW with branching dominant ($\phi(G) = 1 + G^2$ as for DP, but with different vertex-sign assignments) or contact-process variants (same DSE), the bridge basis $\{B_2, B_3^{\text{sun}}, B_V\}$ is unchanged. The system-specific cost is recomputing the algebra factors a_Γ from the new vertex structure — a finite rational task. No fresh perturbative integration.

For conserved DP / Manna ($\phi(G) = 1 + G^3$, C_3 stratum, [24]), the bridge basis at the same $d_c = 4$ extends by one rank-3 master $B_3^{(3)}$; the Lagrange recount and IBP reduction are again finite rational tasks.

10.4 Scope

Janssen–Täuber 2005 (canonical) — Scope of the contribution

CFAC factors out the counting problem from the field-theoretic calculation. What remains — the bridge layer of three master integrals at 2-loop DP — is the irreducible analytic content. Those three masters are taken from JT05 / Täuber’s published work, and the same masters are reusable across every system in the C_2 stratum at $d_c = 4$. The transcendental $\ln(4/3)$ enters only through two of those three masters; combinatorics fixes everything else.

11 Reproducibility: the Code Stack

Code-verified (rdft/ac/gribov/two_loop.py) — End-to-end test suite: 8/8 passing

The directory `rdft/ac/gribov/` contains the complete working code stack for this paper. An end-to-end test harness (`run_all_tests.py`) exercises every component:

#	Test	Status
1	Lagrange + 1-loop algebra factors $\{\frac{1}{4}, \frac{1}{8}, \frac{1}{2}, 2\}$	✓
2	Double-pole closed form $Z_X^{(2,2)} = \frac{1}{2}a(\beta_1 + a)$	✓
3	Simple-pole BPHZ counterterm $\frac{1}{2}a(a - \beta_1)$	✓
4	12 IBP coefficients reproducing JT05 Eq. (57) simple poles	✓
5	Full pipeline $Z \rightarrow \beta \rightarrow u^* \rightarrow$ JT05 Eq. (60), zero residual	✓
6	From-scratch IBP solver: $(d - 3)$ identity for sunset	✓
7	Targeted reductions: $(d - 5)$ and $(d - 4)$ Laporta coefficients	✓
8	Full Laporta triangulation: $I[1, 1, 1]$ identified as master	✓
Summary		8/8 pass

Reproduce via `python3 run_all_tests.py`.

CFAC closed form — Code inventory

Paper and references:

- `gribov.tex` / `gribov.pdf` — this paper.

- `jt05.pdf` / `jt05.txt` — Janssen–Täuber 2005 (cond-mat/0409670), reference for masters.

CFAC pipeline (counting + algebra + Hopf + RG):

- `assembly.py` — Lagrange-inversion counts and 1-loop algebra factor derivation.
- `actrick.py` — AC trick at 2 loops: bivariate signed Lagrange and nested-graph derivation.
- `simple_poles.py` — BPHZ closed-form counterterm and primitive-residue extraction.
- `ibp_coefficients.py` — 12 IBP coefficients $q_\bullet^X$ from CFAC structural constraints.
- `two_loop.py` — algebraic pipeline $Z \rightarrow \beta \rightarrow u^*$ to JT05 Eq. (60).

Integration backend (plugin layer):

- `ibp_plugin.py` — FORM backend, computes the sunset Laurent expansion via Gamma-function closed form.
- `ibp_sunset.frm` — FORM script for the sunset.
- `ibp_solver/` — from-scratch open-source Python/SymPy Laporta reducer:
 - `sunset_identity.py` — one IBP identity by hand.
 - `sunset_full.py` — all 6 IBP identities per seed.
 - `reduce.py` — linear-system construction.
 - `triangulate.py` — Laporta top-down elimination (identifies $I[1, 1, 1]$ as the master).
 - `test_reduction.py` — $(d - 5)$, $(d - 4)$ coefficient verifications.

Tests:

- `run_all_tests.py` — 8 end-to-end tests, all passing on the development machine (macOS 26 / Python 3.13 / SymPy 1.13 / FORM 5.0.0 / tectonic for the paper build).

Janssen–Täuber 2005 (canonical) — Dependencies and licensing

The CFAC pipeline (tests 1–5) has no external dependencies beyond the Python/SymPy standard scientific stack. The integration backend (tests 6–8) uses:

- **FORM** [18] (GPL, installed via Homebrew on the development machine).
- **SymPy** (BSD, pip-installable).

No proprietary software is required for the entire pipeline to run. KIRA [19] and FIRE [20] would slot into the integration backend with the same interface; their availability depends on the group’s setup (Mathematica licence and/or Fermat).

12 Open Items

1. **(D2) IBP reduction tables.** Work out the explicit rational-coefficient linear combinations expressing each of V_2^{ice} , V_2^{box} , V_2^{lad} , Σ_2^{nest} in the master basis $\{B_2^2, B_V, B_3^{\text{sun}}\}$. Finite-rank linear algebra; can be automated with FIRE / KIRA / LiteRed. This closes the gap between Section 5’s schematic $q_\bullet^X$ table and an explicit derivation of JT05 Eq. (57).
2. **(D3) Algebra factors a_Γ .** Enumerate the rational a_Γ for each of the seven 1- and 2-loop

topologies from the Reggeon vertex pair $(-u/2, +u/2)$ together with the Z -factor extraction conventions ($\partial_{i\omega}$, ∂_{p^2} , etc.). Pure bookkeeping; can be done by hand or by a small SymPy routine.

3. **Independent evaluation of the masters.** Run Panzer’s HyperInt on B_2 , B_3^{sun} , B_V at the JT05 symmetric point. Cross-check against the JT05 quoted values, removing the JT05 numerical-input dependency entirely.
4. **Reuse on Manna C_3 .** Apply the same pipeline with $\phi(G) = 1 + G^3$ and the rank-3 bridge basis at $d_c = 4$; cross-check against the C_3 paper.
5. **Pedagogical exposition.** Distill Sections 3 to 5 into a 6–8 page note for circulation to Pruessner and group — the *simple statement* requested in correspondence.
6. **Higher loops.** At 3 loops the master basis grows; cosmic-Galois selection rules constrain the transcendentals available; Panzer’s HyperInt evaluates linearly-reducible cases symbolically. CFAC’s counting layer extends uniformly.

References

- [1] H. K. Janssen, “On the nonequilibrium phase transition in reaction-diffusion systems with an absorbing stationary state,” *Z. Phys. B* **42**, 151 (1981).
- [2] J. B. Bronzan and J. W. Dash, “Higher-order ε -terms in the renormalization-group approach to Reggeon field theory,” *Phys. Rev. D* **10**, 4208 (1974).
- [3] H. K. Janssen and U. C. Täuber, “The field theory approach to percolation processes,” *Annals of Physics* **315**, 147 (2005); arXiv:cond-mat/0409670.
- [4] U. C. Täuber, *Critical Dynamics: A Field Theory Approach to Equilibrium and Non-Equilibrium Scaling Behavior*, Cambridge University Press (2014).
- [5] G. Pruessner and R. Garcia-Millan, “Field theories of active particle systems and their entropy production,” *Rep. Prog. Phys.* (2025); arXiv:2211.11906.
- [6] R. Garcia-Millan, J. Pausch, B. Walter, G. Pruessner, “Field-theoretic approach to the universality of branching processes,” *Phys. Rev. E* **98**, 062107 (2018); arXiv:1808.08418.
- [7] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press (2009).
- [8] A. Connes and D. Kreimer, “Hopf algebras, renormalization and noncommutative geometry,” *Commun. Math. Phys.* **199**, 203 (1998); hep-th/9808042.
- [9] D. Kreimer, “On the Hopf algebra structure of perturbative quantum field theories,” *Adv. Theor. Math. Phys.* **2**, 303 (1998); q-alg/9707029.
- [10] E. Panzer, “Algorithms for the symbolic integration of hyperlogarithms with applications to Feynman integrals,” *Comput. Phys. Commun.* **188**, 148 (2015); arXiv:1403.3385.
- [11] E. Panzer, “Feynman integrals and hyperlogarithms,” PhD thesis, Humboldt-Universität zu Berlin (2015); arXiv:1506.07243.
- [12] F. C. S. Brown, “On the periods of some Feynman integrals,” arXiv:0910.0114 (2008).
- [13] F. Brown, “Feynman amplitudes, coaction principle, and cosmic Galois group,” *Commun. Number Theory Phys.* **11**, 453 (2017); arXiv:1512.06409.

- [14] F. Brown, “Notes on motivic periods,” arXiv:1512.06410 (2015).
- [15] M. Borinsky, “Tropical Monte Carlo quadrature for Feynman integrals,” *Ann. Inst. H. Poincaré D* **10**, 635 (2023); arXiv:2008.12310.
- [16] M. Borinsky and H. J. Munch, “Tropical Feynman integration in the Minkowski regime,” arXiv:2302.08955 (2023).
- [17] V. A. Smirnov, *Analytic Tools for Feynman Integrals*, Springer Tracts in Modern Physics **250** (2012).
- [18] J. A. M. Vermaseren, “New features of FORM,” arXiv:math-ph/0010025 (2000). <https://www.nikhef.nl/~form/>.
- [19] P. Maierhöfer, J. Usovitsch, and P. Uwer, “Kira — A Feynman integral reduction program,” *Comput. Phys. Commun.* **230**, 99 (2018); arXiv:1705.05610.
- [20] A. V. Smirnov, “FIRE6: Feynman Integral REduction with modular arithmetic,” *Comput. Phys. Commun.* **247**, 106877 (2020); arXiv:1901.07808.
- [21] S. Laporta, “High-precision calculation of multi-loop Feynman integrals by difference equations,” *Int. J. Mod. Phys. A* **15**, 5087 (2000); arXiv:hep-ph/0102033.
- [22] K. G. Chetyrkin and F. V. Tkachov, “Integration by parts: The algorithm to calculate β -functions in 4 loops,” *Nucl. Phys. B* **192**, 159 (1981).
- [23] S. Amarteifio, “Reaction-diffusion field theory via analytic combinatorics: from stoichiometry to critical exponents,” Percolation Labs (2026). [[paper/cfac/cfac_theorem.tex](#)]
- [24] S. Amarteifio, “CFAC Theorem II: admissible, multivariate and log-corrected extensions; signed-projector C_3 for conserved DP,” Percolation Labs (2026). [[paper/cfac/cfac_theorem_II.tex](#)]