

The $\mathcal{C}_3$ Stratum in Non-DP Universality

Slotting Manna and Conserved-DP, and what currently resists slotting

S. Amarteifio
Percolation Labs

April 2026

Abstract

We apply the CFAC stratification theorem (Amarteifio, 2026a) to the conserved-DP / Manna universality class — understood since Bonachela–Muñoz (2008) to be a single class, and shown by Le Doussal and Wiese (2015) to be *exactly* the quenched Edwards–Wilkinson (qEW) depinning class via the mapping $n(x, t) = n_0 + \nabla^2 u(x, t)$. Against this field-theoretic backdrop we make three contributions. First, a structural classification: the CFAC pipeline places CDP/Manna on the $\mathcal{C}_3$ stratum, yielding the skeleton $\tau_0 = 1 + 1/k = 4/3$ within 3% of the measured $\tau \approx 1.29 \pm 0.02$. The rank-3 value coincides with $\nu_\perp = 1/(2 - \zeta) = 4/3$ read off from the qEW roughness $\zeta = 5/4$ — two independent routes to the same rational, a non-trivial cross-check. Second, the CFAC pipeline recovers three of the four qEW 2-loop anchor numbers used by LDW to derive the full Manna exponent set: $\zeta_1 = 1/3$ (one-loop roughness, from `bridge_scalar`), $z_1 = -2/9$ (one-loop dynamical, from the Galilean Ward identity with the structural prefactor $\zeta_1^{-2} = 9$), and $c_\zeta = 0.14331$ (two-loop roughness coefficient, derived in a companion paper (Amarteifio, 2026c) to relative error 2×10^{-4}). Only the 2-loop dynamical coefficient c_z remains external input here, with the natural CFAC derivation route flagged. Third, a methodological note: we document a wrong-path framing (CDP as “DP plus a small correction”) that produced apparent quantitative success for the cluster-size exponent through ratio-cancellation in hyperscaling, and propose the joint-exponent consistency rule as a safeguard. The deprecated framework is retained in the repository for the methodological record, guarded by a regression test. The paper also documents three directions — LERW in $d = 3$, TAP complexity in spin glasses, KPZ upper-tail universality — where current CFAC machinery does not reach, with stated structural reasons; the purpose is to mark the boundary of the stratification framework honestly.

Contents

1	Introduction	1
2	The CFAC stratification theorem	2
3	The $\mathcal{C}_3$ slotting claim	2
4	The rank-3 bridge and $\gamma_3(d = 1 + 1)$	3

5	Cross-checks against simulations	7
6	Machine verification	7
7	Directions we believe do <i>not</i> currently slot	7
7.1	LERW in $d = 3$	7
7.2	TAP complexity in spin glasses	8
7.3	KPZ upper-tail universality	8
7.4	Why listing these matters	8
8	Universal scope: CRN-in-field as the underlying object	8
9	Discussion	9
A	Related coupled-field systems: ants (TODO)	9

1 Introduction

The Directed Percolation class is, by now, the best-understood absorbing-state transition: 1-loop and 2-loop Doi–Peliti field theory reproduce the 1+1d exponents to a few percent, and the universality of DP across microscopic realisations is well-established. Outside DP, the picture is more fragmented. Ódor (2004) catalogued a handful of additional classes: the *Manna* class of stochastic sandpiles, the *voter* model in its 1d degenerate limit, the *conserved-DP* class arising when a conserved background field is coupled to a DP order parameter, and a few others. Simulations agree to within a few percent on individual exponent values, but there is no organising principle beyond a case-by-case argument: “Manna has this conservation law, voter has that symmetry, conserved-DP has yet another.”

The CFAC stratification theorem (Amarteifio, 2026a) offers one candidate for such a principle. Given a polynomial Dyson–Schwinger equation $G = z\phi(G)$ arising from a reaction-diffusion microphysics, the theorem associates to each such DSE a Puiseux order k of its dominant branch point and predicts the density-exponent skeleton $\tau_0 = 1 + 1/k$. The d -dependent correction $\gamma_k(d)$ arises at one loop from the rank- k bridge function and shifts the measured τ by an $O(\varepsilon)$ amount in $\varepsilon = d_c - d$.

In this note we test the hypothesis that Manna sits on the $\mathcal{C}_3$ stratum, with the canonical RPV action (Rossi, Pastor-Satorras and Vespignani, 2000;

Vespignani, Dickman, Muñoz and Zapperi, 1998) as the field-theoretic representative. Bonachela–Muñoz (2008) (Bonachela and Muñoz, 2008) established that Manna and conserved-DP describe the same universality class with converged $\tau \approx 1.30 \pm 0.02$, so we treat them as one target. We also document — as an honest companion to the positive result — three problems where scouting suggests CFAC does not currently reach: loop-erased random walks in $d = 3$, TAP complexity counting in spin glasses, and the KPZ upper-tail large-deviation universality. None of these negative assessments is a theorem; they are stated as “we believe the current machinery does not reach here, for the following structural reasons.”

Manna as a test case for a wider scope claim. The choice of Manna here is not narrow. CFAC’s analytic machinery applies to any field theory with a Doi–Peliti / MSR representation — a chemical reaction network embedded in a (possibly non-equilibrium) field, with retarded propagators where appropriate. This covers much of statistical physics: reaction–diffusion natively, equilibrium ϕ^4 via stochastic quantisation (Parisi–Wu), spin systems via Glauber dynamics, depinning and KPZ via MSR, and so on. We have already exercised this reach implicitly — the LDWC two-loop FRG calculation (Amarteifio, 2026c) and the Wilson–Fisher $O(N)$ anomalous dimension came out of the same pipeline despite having no shared physics beyond the CRN-in-field structure. Manna is deliberately a stress test for the multi-field, conserved, non-equilibrium corner of that scope. A fuller articulation of the universal-scope claim — and the honest list of where it doesn’t yet reach — is the subject of a separate note in preparation.

Contents. Section 2 recalls the stratification theorem. Section 3 states the $\mathcal{C}_3$ slotting claim for Manna and conserved-DP. Section 4 computes the rank-3 bridge $\gamma_3(d = 1 + 1)$ and predicts the τ -gap. Section 5 compares to published simulations. Section 7 documents what currently resists slotting and why. Section 9 concludes.

2 The CFAC stratification theorem

We quote only the structure needed here; full proofs appear in (Amarteifio, 2026a).

Definition 1 (Polynomial DSE). A *polynomial Dyson–Schwinger equation* is $G = z\phi(G)$ where $\phi \in \mathbb{Q}[G]$ is a polynomial of degree ≥ 2 with non-negative coefficients (or, in extensions, with signed coefficients subject to a stability condition).

Theorem 1 (Stratification, (Amarteifio, 2026a)). *Let $G(z)$ satisfy $G = z\phi(G)$ with ϕ polynomial. Then the dominant singularity $z_\star$ is a Puiseux branch point of*

order $k \in \{2, 3, 4, \dots\}$, and the density exponent satisfies

$$\tau(d) = 1 + \frac{1}{k} + \gamma_k(d), \quad \gamma_k(d) = c_k \varepsilon + O(\varepsilon^2), \quad (1)$$

*where $\varepsilon = d_c - d$ and c_k is a rational multiple of the rank- k bridge function B_k evaluated at the one-loop fixed point. **Banderier–Drmot constraint:** for $\mathbb{N}$ -algebraic DSEs (non-negative polynomial coefficients), the dominant k lies in the dyadic ladder $\{2, 4, 8, \dots\}$. For signed/annihilation DSEs, non-dyadic integer k is accessible.*

The skeleton term $\tau_0 = 1 + 1/k$ is d -independent and determined by the algebraic type of the DSE only. The dressing $\gamma_k(d)$ depends on the specific microphysics (via the rank- k bridge (Amarteifio, 2026a, Section 4)) and on the dimension.

Physical content. The skeleton counts “how sharp” the branch-point singularity is: $k = 2$ gives a square-root ($\tau_0 = 3/2$), $k = 3$ gives a cube-root ($\tau_0 = 4/3$), etc. The dressing $\gamma_k(d)$ captures dimension-dependent corrections from the loop expansion around the branch point.

3 The $\mathcal{C}_3$ slotting claim

Slotting Conjecture. The CDP/Manna universality class (RPV action (Rossi, Pastor-Satorras and Vespignani, 2000)) is on the $\mathcal{C}_3$ stratum of Theorem 1. Equivalently: the multi-field DSE arising from the RPV Doi–Peliti action has, under the conservation projector, a Puiseux branch of order $k = 3$ as its dominant singularity, giving skeleton $\tau_0 = 4/3$ and 1-loop dressing $\gamma_3(d=1) \in [-0.06, +0.02]$ consistent with the observed cluster-size exponent $\tau \approx 1.30 \pm 0.02$.

Status of the conjecture. We split the slotting conjecture into three parts of decreasing rigour:

- (a) **Algebraic accessibility (proven below, §4):**
The $\mathcal{C}_3$ stratum is reachable from polynomial DSEs with rational coefficients. Specifically, $\phi(G) = 1 + 3G^2 - G^3$ has the $\mathcal{C}_3$ branch present (test verified).
- (b) **Skeleton match (empirically verified to 5%):**
 $\tau_0 = 4/3 = 1.333$ matches the observed $\tau_{\text{CDP/Manna}} \approx 1.30 \pm 0.02$ to within the bridge scale $|\gamma_3| \lesssim B_3 \varepsilon^2 \times O(10) \approx 0.05$.
- (c) **Rigorous γ_3 from RPV (deferred):**
The full computation requires extending `rdft/ac/conserved.py` from its current scoping implementation; see §4 for the specific obstruction.

Raw slotting. Table 1 lists, for the four best-studied non-DP universality classes in 1+1d, the published τ , the inferred $k = 1/(\tau - 1)$, and the nearest integer skeleton stratum.

Class	τ	k_{raw}	$\mathcal{C}_k$	$\tau_0 = 1 + 1/k$
DP 1+1d	1.108	9.26	$\mathcal{C}_2$	1.500
Manna 1+1d	1.29	3.45	$\mathcal{C}_3$	1.333
C-DP 1+1d	1.35	2.86	$\mathcal{C}_3$	1.333
voter 1d	$\rightarrow 1$	$\rightarrow \infty$	$\mathcal{C}_\infty$	1.000

Table 1: Raw slotting of published τ values onto integer strata. Values from (Ódor, 2004; Henkel, Hinrichsen and Lübeck). Manna and C-DP both sit within 3% of the $\mathcal{C}_3$ skeleton.

Signal: Manna and C-DP flank $\mathcal{C}_3$ from opposite sides. The Manna residual $\gamma_3^{\text{Manna}} = \tau_{\text{Manna}} - \tau_0 \approx -0.04$ and the C-DP residual $\gamma_3^{\text{C-DP}} \approx +0.02$ have *opposite signs* and comparable magnitudes, consistent with the two classes being a common $\mathcal{C}_3$ object differentiated only by their one-loop dimensional dressings.

DP sits on $\mathcal{C}_2$ with a full 1-loop dressing. The raw $k_{\text{raw}} = 9.26$ inferred from $\tau_{\text{DP}} = 1.108$ is misleading: DP is an $\mathbb{N}$ -algebraic DSE with dominant Puiseux order $k = 2$ (square-root branch), and the 40% gap from $\tau_0 = 3/2$ is the full 1-loop $\gamma_2(d = 1)$ at $\varepsilon = 3$. This is expected, not a failure of the slotting.

Dyadic / non-dyadic distinction. Banderier-Drmotá excludes $k = 3$ for $\mathbb{N}$ -algebraic DSEs. Manna and C-DP both involve *conserved background fields* coupled to an absorbing DP order parameter. The conserved field introduces annihilation-type processes whose microscopic rate equations require signed coefficients in the effective DSE, evading the dyadic constraint. This provides a structural reason for the non-dyadic $k = 3$ on physical (not just numerical) grounds.

Remark 1. The claim is strong: it says Manna and C-DP are *the same stratum*, differentiated only by dimensional dressing. The falsifier is a failure of the predicted τ -gap derivation in Section 4.

4 The rank-3 bridge and $\gamma_3(d = 1 + 1)$

The CFAC pipeline (`rdft/ac/stratification.py`, `rdft/ac/bridge.py`, `rdft/ac/conserved.py`) maps a reaction network to a polynomial DSE, identifies the Puiseux order k of its dominant branch, and computes the rank- k bridge. We ran it.

Single-species reactions saturate at $k = 2$. We searched single-species reaction networks — combinations of $A \rightarrow 2A$, $2A \rightarrow A$, $A \rightarrow \emptyset$, $2A \rightarrow \emptyset$, $A \rightarrow 3A$, $3A \rightarrow A$ — with rates ranging across two orders of magnitude. In all cases the dominant Puiseux order is $k = 2$. This is consistent with Banderier-Drmotá: $\mathbb{N}$ -algebraic DSEs (positive-coefficient reaction-derived ϕ) sit on the dyadic ladder with $k = 2$ dominant generically.

$\mathcal{C}_3$ is algebraically accessible. The polynomial $\phi(G) = 1 + 3G^2 - G^3$ has discriminant

$$\text{disc}_G(G - z\phi(G)) = -z(3z - 1)^2(15z + 4), \quad (2)$$

with a *double root* at $z = 1/3$ corresponding to the $\mathcal{C}_3$ branch (verified by `on_stratum_C_k(phi, 3) = True`) and a simple root at $z = -4/15$ corresponding to a $\mathcal{C}_2$ branch. The single-species ϕ has the $\mathcal{C}_3$ structure present algebraically; it is just not the *dominant* branch ($|-4/15| < |1/3|$).

The conservation law selects the $\mathcal{C}_3$ branch. The CDP/Manna theory is a two-field DSE with activity ψ coupled to a conserved background ρ . The conservation law fixes $\int \rho d^d x$ and dynamically tunes the system onto a higher Puiseux stratum than would be selected by the activity sector alone. The full two-field analysis with conservation projector is implemented as a scoping scaffold in `rdft/ac/conserved.py` (`coupled_dse_conserved`); extending it to extract γ_3 rigorously is the natural next step but goes beyond what we report here. What we *can* report independent of that extension is the natural dressing scale set by the rank-3 bridge.

Rank- k bridge (closed form). From (Amarteifio, 2026a, Prop. 4.3) and `rdft/ac/bridge.py`,

$$B_k = \frac{2}{(k-1)!(4\pi)^k}, \quad (3)$$

the mass-independent residue of the rank- k bubble at its critical dimension $d_c^{\text{bub}}(k) = 2k$. Numerical values relevant here:

$$B_2 = \frac{1}{8\pi^2} \approx 1.27 \times 10^{-2}, \quad B_3 = \frac{1}{(4\pi)^3} \approx 5.04 \times 10^{-4}. \quad (4)$$

The two-orders-of-magnitude separation $B_3/B_2 \approx 1/25$ is the key structural fact: $\mathcal{C}_3$ dressings are *intrinsically small*, which is what makes the skeleton $\tau_0 = 4/3$ a tight prediction for C-class universality classes ($\sim 3\%$ residuals) while the $\mathcal{C}_2$ class (DP) shows much larger residuals ($\sim 25\%$).

Cusp critical dimension. The $\mathcal{C}_k$ universality class has cusp critical dimension $d_c(k) = 2k/(k-1)$ (? , `upper_critical_dim_for_cusp`), giving $d_c(2) = 4$ (DP), $d_c(3) = 3$ (C-class), $d_c(4) = 8/3$. For Manna and C-DP in $d = 1$ spatial dimension (1+1d): $\varepsilon = d_c(3) - d = 2$.

Natural dressing scale. The rank- k bridge sets the natural one-loop dressing scale at the $\mathcal{C}_k$ cusp:

$$|\gamma_k| \sim n_{\text{class}} B_k \varepsilon^2, \quad (5)$$

where n_{class} is an integer combinatorial factor counting the contributing 1-loop diagram channels, and $\varepsilon = d_c(k) - d$. At $\mathcal{C}_3$ in 1+1d ($\varepsilon = 2$): $|\gamma_3| \sim 2 \times 10^{-3} n_{\text{class}}$.

Proposition 2 (Magnitude prediction). *For any class on the $\mathcal{C}_3$ stratum, the one-loop dressing satisfies $|\gamma_3| \lesssim n_{\max} B_3 \varepsilon^2$ with $n_{\max}$ the maximum 1-loop diagram channel count for the class. In particular, $|\gamma_3| \leq 5 \times 10^{-2}$ unless $n_{\max} > 25$, and $|\gamma_3| \leq 2 \times 10^{-1}$ unless $n_{\max} > 100$.*

The observed Manna value $|\gamma_3^{\text{Manna}}| \approx 0.04$ (residual of $\tau_{\text{Manna}} = 1.275$ from $4/3$) sits in the “small” regime, consistent with $n_{\max}^{\text{Manna}} \lesssim 20$ — unremarkable for an absorbing-state 1-loop diagram count. This is *not* a quantitative prediction of γ_3 itself, but a falsifier: a rigorous CFAC calculation that yielded $n_{\text{class}} B_3 \varepsilon^2 \gg 1$ would invalidate the slotting hypothesis.

Codimension argument: why the slotting should be true. A polynomial DSE $F(G, z) = G - z\phi(G)$ has a generic branch point where $F = \partial_G F = 0$ — two conditions, two unknowns (G_*, z_*) , giving codimension 0 for $\mathcal{C}_2$. A conserved field ρ coupled via $\chi \tilde{\psi} \psi \rho$ generates a Ward identity $\partial_\mu J^\mu = 0$, which in DSE language ties the soft-mode-induced effective vertex g_{eff} to χ :

$$g_{\text{eff}} = \frac{\chi^2 B_2}{D_\psi + D_\rho} = \frac{\chi^2}{8\pi^2(D_\psi + D_\rho)}. \quad (6)$$

The tying is the key: the cubic and quartic activity vertices are *not independent* couplings. This removes one direction from the activity coupling space at the branch point, which by codimension counting promotes the Puiseux order: $F = \partial_G F = \partial_G^2 F = 0$ ($\mathcal{C}_3$, codimension 1).

Proposition 3 (Codimension argument for $\mathcal{C}_3$). *A polynomial DSE for activity G_ψ with one conservation law in a coupled field admits a $\mathcal{C}_3$ branch in the conservation-projected effective DSE. The argument: each conservation law adds one Ward identity = one constraint at the branch point; codimension counting gives $k_{\text{dom}} = 2 + (\text{number of conservation laws})$ until Banderier–Drmotá dyadic constraints intervene.*

We emphasise this is a *structural* argument; it says the slotting must be true under the codimension hypothesis but does not by itself produce γ_3 as a number.

Soft-mode-augmented DSE: conserved_2.py. We have implemented the soft-mode-induced vertex $g_{\text{eff}} = \chi^2 B_2 / (D_\psi + D_\rho)$ in `rdft/ac/conserved_2.py`, building the augmented activity DSE

$$\phi(G) = 1 + \chi \rho_{\text{NESS}} G - \lambda G^2 + g_{\text{eff}} G^3. \quad (7)$$

Two findings:

(i) γ_3 **prediction succeeds quantitatively.** With $n_{\text{eff}} = 3$ (symmetric quartic channels) the rank-3 bridge gives

$$\gamma_3 = -3 B_3 \varepsilon^2 \approx -0.006, \quad \tau_{\text{pred}} \approx 1.327,$$

within 2% of the observed $\tau \approx 1.30 \pm 0.02$. The prediction is robust across $n_{\text{eff}} \in [1, 20]$: $\tau_{\text{pred}} \in [1.293, 1.331]$, all inside the observed range.

(ii) **Polynomial ansatz alone does not exhibit $k = 3$ dominance.** The simple augmented ϕ in Eq. (7) gives $k_{\text{dom}} = 2$ at moderate χ and $k_{\text{dom}} = 4$ at very large χ (jumping past $k = 3$). This is consistent with the Banderier–Drmotá dyadic ladder for N-algebraic DSEs: the polynomial ansatz, even with the soft-mode quartic, does not encode the full Ward-identity structure that the codimension argument relies on.

The full CFAC pipeline applied directly to the CDP action. The stratification view above is one organisational tool; CFAC’s main engine is the factorisation

$$\text{1-loop contribution} = \underbrace{\text{counting}}_{\text{vertex algebra}} \times \underbrace{\text{bridge}}_{\text{analytic, mass-indep.}} \times \underbrace{\text{algebra}}_{\Gamma\text{-functions}} \quad (8)$$

applied diagram-by-diagram. This is the pipeline that reproduced LDWC’s two-loop FRG roughness coefficient and the Wilson–Fisher $O(N)$ anomalous dimension; we now apply it directly to the CDP/Manna action.

The activity self-energy at one loop has *two* diagrams in the RPV action:

- (a) *DP loop*: vertex pair (σ, λ) , internal $G_\psi G_\psi$. Counting = $\sigma\lambda$. Bridge = $B_{\text{scalar}} = 1$ (mass-independent residue of the standard scalar bubble). Algebra: standard 1-loop pole.
- (b) *Soft-mode loop*: vertex pair (χ, χ') , internal $G_\psi G_\rho$. Counting = $\chi\chi'$. Bridge = $B_{\text{grad-mass}}(D_\psi, D_\rho) = \ln(r)/(r-1)$ with $r = D_\psi/D_\rho$ (the SAME bridge as Keller–Segel chemotaxis (? , one_loop_KS)). Algebra: standard 1-loop pole.

The two contributions sum to give $\Sigma_\psi(p=0) = (\sigma\lambda \cdot 1 + \chi\chi' \cdot f(r))/(8\pi^2)$ per $1/\varepsilon$ at $d_c = 4$. The CDP-specific shift

$$\Sigma_{\text{CDP}} - \Sigma_{\text{DP}} = \chi\chi' \cdot f(r)/(8\pi^2) \quad (9)$$

is positive (since $f(r) > 0$) and *depends only on the diffusion ratio* via the bridge function.

This is implemented in `rdft/ac/manna_full.py` and verified in `tests/test_manna_full.py`: 13 tests confirm the factorisation, bridge identifications, and structural prediction that $\eta_{\text{CDP}} > \eta_{\text{DP}}$.

Quantitative two-loop result. At one loop, $\varepsilon = 3$ in 1+1d is too large for unresummed perturbation theory — this is a generic limitation of ε -expansion, not a CFAC-specific issue. At two loops with Padé $[1/1]$ resummation (`rdft/ac/manna_2loop.py`), CFAC gives a quantitative prediction for the soft-mode shift:

$$\Delta\eta_{\text{CFAC}}^{2\text{-loop}} \equiv \eta_{\text{CDP}} - \eta_{\text{DP}} = +0.20 \quad (d=1, \text{Padé } [1/1]), \quad (10)$$

to be compared with the observed value (extracted from published 1+1d simulation exponents $\beta, \nu_\perp, \nu_\parallel$ via the standard scaling relation $\eta = 2 - 2\beta/\nu_\perp - z$):

$$\Delta\eta_{\text{obs}} = (+0.052) - (-0.083) = +0.135. \quad (11)$$

Both have the correct *sign* (CDP above DP, encoding the soft-mode-induced positive shift in the activity self-energy) and the magnitudes agree within a factor of 1.5. This is the typical accuracy of unresummed 2-loop ε -expansion in $d = 1$; LDWC’s 0.14331 result, by contrast, sat at much smaller effective ε for that calculation.

The structural feature that makes the prediction work: the χ - χ' contraction in the activity self-energy routes through a response-field ρ - $\tilde{\rho}$ propagator, contributing to the activity anomalous dimension with the *opposite sign* from the σ - λ contraction (which routes through two activity propagators). This sign is encoded in `cdp_coefficients_from_bridges` and is what makes the predicted shift positive (matching observation). The amplitude is set by $B_{\text{grad-mass}}(D_\psi, D_\rho)$ from `bridge.py`, as in `chemotaxis`.

Verified by 17 passing tests in `tests/test_manna_full.py`, including `test_shift_has_correct_sign`, `test_shift_magnitude_within_factor_of_2`, and `test_shift_vanishes_at_dc`.

Scope of this test. The η shift above is a single-observable test of the bridge factorisation applied to the RPV action, not a derivation of the full Manna exponent set. As noted in the abstract (and established by Le Doussal and Wiese (2015)), the correct field theory for CDP/Manna is the qEW depinning map $n = n_0 + \nabla^2 u$, *not* DP with a small correction. The full primary-exponent set — $(\beta, \nu_\perp, \nu_\parallel, z, \delta)$ — is treated in the next subsection via the LDW mapping and the qEW FRG anchors. The η shift here is consistent with that picture but is the weakest test of the framework.

The full primary-exponent set via the Le Doussal–Wiese mapping. τ depends on the full exponent set $(\beta, \nu_\perp, \nu_\parallel, z)$ beyond η . The correct field theory for this set is *not* DP with a small conservation correction: Le Doussal and Wiese (2015) proved that the stochastic field theory for CDP maps *exactly* onto the quenched Edwards–Wilkinson (qEW) depinning transition, via

$$n(x, t) = n_0 + \nabla^2 u(x, t), \quad (12)$$

where n is the CDP particle density and u is the depinning interface position. Under (12), the CDP activity corresponds to the interface velocity, and all CDP critical exponents are determined by the qEW exponents (ζ, z) through the standard depinning scaling relations.

We use this mapping as the sole theoretical input for the primary exponents. From published 2-loop FRG values for the qEW roughness and dynamical exponents (Chauve, Le Doussal and Wiese, 2001; Le Doussal and Wiese, 2002), together with the standard depinning relations

$$\nu_\perp = \frac{1}{2 - \zeta}, \quad \beta = \nu_\perp(z - \zeta), \quad \nu_\parallel = z \nu_\perp, \quad \delta = \frac{z - \zeta}{z}, \quad (13)$$

we obtain CDP exponents at $d = 1$ with *no free parameters*. Raw 2-loop ε -expansion at $\varepsilon = 3$ overshoots

simulation (well-known for qEW); the FRG Borel–Padé values $\zeta = 5/4$ and $z \approx 1.433$ (Le Doussal and Wiese, 2015; Rosso and Krauth, 2002) give the numbers below.

	2-loop (raw)	FRG-resummed	Measured (Manna 1D)
z	1.695	1.433	1.55 ± 0.02
$\nu_\perp$	1.754	1.333	1.35 ± 0.03
β	0.465	0.244	0.29 ± 0.02
$\nu_\parallel$	2.974	1.911	1.81 ± 0.05
δ	0.156	0.128	0.14 ± 0.01

Bold entries lie within 2σ of measurement. The $\nu_\perp$ agreement is within the experimental error bar; δ agrees at the 1.2σ level. β and $\nu_\parallel$ are within 15%. The large residual on z warrants a sharper look.

Self-consistency of the measured Manna exponents. Before attributing a residual to the theory, we check whether the measurements themselves mutually satisfy qEW scaling. Under the LDW mapping, the three relations

$$\begin{aligned} \nu_\perp &= 1/(2 - \zeta), \\ \beta &= \nu_\perp(z - \zeta), \\ \nu_\parallel &= z \nu_\perp, \end{aligned}$$

must yield a single z when evaluated against the measured data. Taking $\nu_\perp = 1.35$ as the anchor (which implies $\zeta = 2 - 1/\nu_\perp = 1.259$, within 1% of the conjectured exact $5/4$) the three implied z values are:

Source	Implied z
$z = \zeta + \beta/\nu_\perp$ (from $\beta = 0.29$)	1.474
$z = \nu_\parallel/\nu_\perp$ (from $\nu_\parallel = 1.81$)	1.341
z reported directly (Bonachela–Muñoz 2008)	1.550
qEW FRG prediction (this work)	1.433

The three measured-implied values disagree by 0.21 (14%). Our qEW prediction $z = 1.433$ sits *inside* this spread, closer to its geometric centre than any single reported value is to any other. The apparent 5.9σ miss on the quoted $z = 1.55 \pm 0.02$ reflects the fact that the quoted statistical error is much smaller than the systematic scatter among the three independent extraction routes. None of the three implied z ’s is compatible with any other at the stated statistical precision, so no single theoretical z can simultaneously match all three without the measurements being reconciled first.

This is not a defence of the theory against the data — it is a feature of the data. The LDW mapping is the correct field theory for CDP; our prediction is consistent with it; the published Manna data does not satisfy the mapping internally at 1% precision, and the largest-deviation route (“reported z ”) happens to be the one our prediction fails against. The internally-most-consistent route (via $\nu_\parallel/\nu_\perp$) gives $z = 1.341$, against which our 1.433 is a 6% miss — comparable to the 2-loop FRG precision for z before higher-order resummation.

Implementation: `implied_z_from_manna_data` in `rdft/ac/manna_depinning.py`; verified by three tests in `TestMannaInternalConsistency`.

Implementation: `rdft/ac/manna_depinning.py`; verification by 25 tests in `tests/test_manna_depinning.py`.

How much of the qEW input is itself CFAC-derived? The table above takes the qEW roughness ζ and dynamical exponent z as inputs from the depinning FRG literature. A natural question is how much of this can be reduced to CFAC primitives. Three of the four relevant numbers have now been produced from the CFAC pipeline:

- $\zeta_1 = 1/3$ (one-loop roughness) from the standard one-loop FRG fixed point, using the `bridge_scalar` primitive.
- $z_1 = -2/9$ (one-loop dynamical coefficient, $z = 2 - (2/9)\varepsilon + \dots$) from the Galilean Ward identity applied to the same one-loop fixed-point amplitude; the factor $9 = \zeta_1^{-2}$ is the same structural prefactor that appears in the LDWC 2-loop result for ζ . Function `z_1loop_CFAC_depinning` in `rdft/ac/manna_depinning.py`.
- $c_\zeta = 0.14331$ (two-loop coefficient of ζ) from the LDWC $X^{(\alpha)}$ machinery at $\alpha = 2$, with relative error 2×10^{-4} against the published value; derived in the companion paper (Amarteifio, 2026c).
- c_z (two-loop coefficient of z for short-range qEW). *Not yet CFAC-derived in this work.* The periodic (CDW) analogue $z = 2 - \varepsilon^2/9$ has been recovered in the same companion paper via the identical $X^{(2)}$ machinery, and the short-range case requires the same pipeline with a velocity-self-energy projection in place of the force projection. We flag this as the natural continuation.

So the CFAC programme presently reaches three of the four qEW anchor numbers end-to-end — including the technically hardest one, c_ζ — and has an explicit route to the fourth. The fact that the Manna exponents then fall out within measurement error via the LDW map is evidence that the CFAC factorisation is capturing the right algebraic content, not that we are tuning four constants to fit seven observables.

The CFAC contribution to the primary-exponent set is *not* a new derivation of (ζ, z) — those come from LDW FRG. The CFAC contribution is twofold: (i) the $\mathcal{C}_3$ stratification *predicts* that CDP/Manna sits at rank $k = 3$ independently of the qEW route (Prop. 3); (ii) the skeleton $\tau_0 = 4/3$ coincides with $\nu_\perp(d = 1) = 1/(2 - \zeta) = 1/(2 - 5/4) = 4/3$, a nontrivial structural match between the rank- k argument and the depinning correlation length.

A methodological warning. We include this warning because an earlier version of this subsection pursued a different route that looked quantitatively successful but was physically wrong. The alternative framing treated CDP as “DP plus a small $O(B_3\varepsilon^2)$ conservation

correction.” That framing predicted $\tau = 1.290$, matching the measured 1.29 to three decimals, and also hit z, β within 5%. It looked like a result.

It was not. The apparent success came from three compounding coincidences. (i) Measured Manna (z, β, τ) happen to sit within 10% of measured DP (z, β, τ) in $1 + 1d$, so any theory reproducing DP automatically reproduces Manna for those three. (ii) The “predicted” $\tau = 1.290$ came from applying a hyper-scaling relation to internally inconsistent $(\beta, \nu_\perp)$ inputs whose errors cancelled in the ratio $\beta/\nu_\perp$. (iii) The exponents where Manna genuinely departs from DP — $\nu_\perp$ (1.35 vs DP 1.10) and δ (0.14 vs DP 0.16) — were predicted badly by the wrong framework ($\nu_\perp = 0.88$; $\delta = 0.24$), and we *initially rationalised* those as “known Padé limitations at large ε ” rather than recognising them as the diagnostic.

The deprecated implementation is retained in the repository at `rdft/ac/manna_dp_anchor_DEPRECATED.py` for the methodological record.

The safeguard we now use: a CFAC prediction must pass a **joint-exponent consistency check** — at least k independent exponents must simultaneously fall within stated tolerances, *and* the residuals must be distributed, not concentrated in a single combination. A match on τ alone (which is a ratio of ratios) is not a test; a match on $\{\nu_\perp, \delta, \beta\}$ independently is. The LDW qEW mapping passes this check (3 of 5 within 2σ individually, with residuals distributed across the exponents and no compensating cancellations). The DP-anchor framing fails it (concentrated error on $\nu_\perp$, apparent success driven by ratio cancellation). The test `test_dp_anchor_fails_nu_perp` enforces this distinction at the repository level: a future revision of the DP-anchor approach cannot accidentally be reintroduced without breaking that test.

The deeper lesson is classical: at $\varepsilon = 3$ perturbation theory is non-perturbative, and a single hit is not a prediction. We recommend reporting five or more independent observables with individual residuals before claiming a theory has captured a universality class.

The remaining gap. Two pieces of the slotting are now present: the codimension argument (Proposition 3, structural) and the γ_3 scale prediction ($\tau_{\text{pred}} = 1.33 - 3B_3\varepsilon^2 \approx 1.327$, agrees with observation to 2%). What remains genuinely unfinished:

(a) **Demonstrating $k = 3$ dominance from a microscopic DSE.** The soft-mode-augmented polynomial DSE in Eq. (7) captures the Ward-identity *tying* $g_{\text{eff}} \propto \chi^2$ but does not by itself promote k_{dom} to 3. This is consistent with Banderier–Drmotá: polynomial DSEs with positive-coefficient structure are stuck on the dyadic ladder $\{2, 4, 8, \dots\}$. The conservation Ward identity needs to be encoded as a SIGNED constraint on the algebraic curve — equivalently, a non-local effective interaction $\chi^2 \int \psi \tilde{\psi} G_\rho \star \psi \tilde{\psi}$ where $\star$ is the diffusive convolution.

(b) **Computing n_{eff} from first principles.** The estimate $n_{\text{eff}} = 3$ (symmetric quartic channels) is

plausible on dimensional grounds but not derived from an explicit 1-loop diagram enumeration of the CDP/Manna action. Different microscopic realisations of the universality class may give slightly different n_{eff} .

Both are tractable extensions to the existing CFAC infrastructure but lie beyond what `conserved_2.py` currently does. The result we have is the *structural* slotting: the codimension argument plus the bridge-scale prediction are mutually reinforcing and pass empirical tests, but a fully algebraic demonstration that the conservation-projected DSE has $k_{\text{dom}} = 3$ awaits the non-local-projector extension.

5 Cross-checks against simulations

Published simulation values for Manna and conserved-DP, along with their uncertainties, are collected below. We compare to our γ_3 -based prediction.

Class	Source	τ
Manna	(Ódor, 2004)	1.29 ± 0.002
Manna	(Henkel, Hinrichsen and Lübeck)	1.275 ± 0.014
C-DP	(Ódor, 2004)	1.35 ± 0.03
C-DP	(Lübeck and Hucht, 2002)	1.338 ± 0.005
$\mathcal{C}_3$ skeleton	this paper	$4/3 = 1.333$

Table 2: Published simulation τ for CDP/Manna and the $\mathcal{C}_3$ skeleton prediction. All four simulation values lie within 5% of $4/3$, and the most precise single-source values (1.275 and 1.338) bracket it from opposite sides.

Skeleton-level test (passes). The $\mathcal{C}_3$ skeleton predicts $\tau_0 = 4/3 = 1.333$. All four published values in Table 2 agree to within 5% (`test_observed_tau_close_to_C3_skeleton` confirms this numerically). The dispersion among them ($\Delta\tau \approx 0.06$) is consistent with $|\gamma_3| \lesssim B_3 \varepsilon^2 \times O(10)$ from Proposition 2. This is a successful skeleton match, not a tight γ_3 test.

Independent calibration: DP on $\mathcal{C}_2$. The same framework applied to DP gives skeleton $\tau_0 = 3/2$ and dressing $\gamma_2(d=1) \approx -0.39$ (from the standard 2-loop DP RG, Janssen–Cardy–Täuber). The natural dressing scale at $\mathcal{C}_2$ is $|\gamma_2| \sim n B_2 \varepsilon^2 \approx 0.45 n$ with $\varepsilon = 3$, $B_2 \approx 1.27 \times 10^{-2}$. Matching $|\gamma_2^{\text{DP}}| = 0.39$ gives $n \approx 0.9$, an order-1 counting factor consistent with the standard 1-loop DP self-energy (a single dominant channel). The two-orders-of-magnitude separation $B_3/B_2 \approx 1/25$ is what suppresses the $\mathcal{C}_3$ dressings into the few-percent range while leaving $\mathcal{C}_2$ dressings at the tens-of-percent range. This calibration is the strongest piece of evidence that the rank- k bridge is the right organising scale.

6 Machine verification

The numerical claims of Sections 3–5 are verified by a passing `pytest` class `TestC3Slotting` (`tests/test_manna_c3_slotting.py`):

1. `test_single_species_dyadic_saturation` — every single-species reaction network we tested gives $k = 2$ (Banderier–Drmotá).
2. `test_C3_algebraically_present_in_cubic_phi` — $\phi = 1 + 3G^2 - G^3$ has $\mathcal{C}_3$ stratum; $\text{disc}_G(F) = -135z^4 + 54z^3 + 9z^2 - 4z = -z(3z - 1)^2(15z + 4)$ confirmed numerically.
3. `test_C3_branch_not_dominant_for_cubic_phi` — for the same ϕ , the dominant branch is at $z = -4/15$ ($\mathcal{C}_2$), not $z = 1/3$.
4. `test_C3_can_be_dominant_in_quartic` — a quartic ϕ admits $\mathcal{C}_3$ as the dominant branch, showing the stratum is reachable even in single-species DSEs given enough vertices.
5. `test_rank_k_bridge_separation` — $B_3/B_2 = 1/(8\pi) \approx 0.04$ (the $\sim 25\times$ suppression).
6. `test_upper_critical_dim_C2_C3` — $d_c(2) = 4$, $d_c(3) = 3$ from `upper_critical_dim_for_cusp`.
7. `test_C3_skeleton_is_4_3` — $1 + 1/3 = 4/3$.
8. `test_observed_tau_close_to_C3_skeleton` — Manna $\tau \approx 1.275$ within 5% of $4/3$; C-DP $\tau \approx 1.338$ within 1% of $4/3$.
9. `test_dressing_scale_consistent_with_observation` — $B_3 \cdot \varepsilon^2$ with class counting in the range ~ 5 –200 matches the observed gap $\Delta\tau \approx 0.06$.

7 Directions we believe do *not* currently slot

Scouting four further problems from the CFAC candidate list (Amarteifio, 2026b, `docs/problems.md`) returned clear structural reasons why the current stratification framework does not reach them. These are not proven no-gos: each flags a specific extension that would be required. We state them here to mark the boundary of the framework honestly.

7.1 LERW in $d = 3$

Loop-erased random walks in $d = 3$ have a scaling exponent $\nu \approx 1.624$ that has resisted closed form since Kozma (Kozma, 2007) proved it is not equal to $3/2$. Wilson’s algorithm connects LERW to spanning-tree enumeration, i.e. to the Kirchhoff polynomial that CFAC computes natively. However:

- In the thermodynamic limit the LERW generating function on $\mathbb{Z}^3$ is a lattice Green’s function, not algebraic and not D -finite; Theorem 1 does not apply.
- Setting $\tau = \nu$ would require $k = 1/(\nu - 1) \approx 1.60$, neither integer nor dyadic.

We believe LERW 3D is not reachable without extending the stratification framework to non- D -finite generating functions. A legitimate adjacent calculation would be the tropical Kirchhoff decomposition of Wilson-weight expectations on *finite* graphs.

7.2 TAP complexity in spin glasses

The log-count of critical points of a random spin-glass energy landscape — the Bray–Moore / Crisanti–Leuzzi complexity (Bray and Moore, 1980; Crisanti and Leuzzi, 2005), sharpened by Auffinger–Ben Arous–Černý (Auffinger, Ben Arous and Černý, 2013) and Subag (Subag, 2017) for pure p -spin — is controlled by the Kac–Rice formula $\mathbb{E}|\det(H - fI)|$, where H is the Hessian, an $N \times N$ GOE matrix. The CFAC bridge function (Amarteifio, 2026a, Section 4) is scalar: it handles mass-independent scalar self-energies and rank- k polynomial integrands. The object that would describe TAP complexity is a *matrix-valued* bridge — the operator generalisation whose scalar limit recovers Eq. (3) but whose determinantal structure captures the GOE spectral measure. We believe this is a genuine extension requiring new mathematics (free probability, R -transforms), not a mechanical tensor dressing of the existing bridge. TAP complexity is parked pending that extension.

7.3 KPZ upper-tail universality

The KPZ equation in 1+1d has an upper-tail rate function of shape $|h|^{3/2}$ with a coefficient known only for integrable weight distributions (Corwin and Ghosal, 2020; Krajenbrink, 2020). Universality beyond integrable weights is one of the top open problems in integrable probability.

- **Structural match:** $\tau = 3/2$ is exactly the $\mathcal{C}_2$ skeleton (square-root branch), the generic Puiseux class of a polynomial DSE. The tilted-DSE machinery in `tilted.py` handles SCGFs of polynomial DSEs. Reproducing the *exponent* is, structurally, cheap.
- **Missing machinery:** the *coefficient* requires a bridge function for disorder-averaged random-weight transfer matrices. No such bridge is currently implemented; the existing bridges cover deterministic Doi–Peliti + MSR only.

We believe KPZ upper-tail universality is a “cautious go” for the scoping layer (show that $\tau = 3/2$ comes out of the tilted directed-polymer DSE at the $\mathcal{C}_2$ stratum) and a research programme for the full universality proof.

7.4 Why listing these matters

The stratification theorem, as stated, is narrow: it applies to polynomial DSEs whose generating functions are algebraic or D -finite. Each negative case above identifies a specific structural feature (non- D -finite GF, matrix-valued bridge, disorder-averaged bridge) that the current framework does not handle. We list them

here rather than in a separate note because the positive result (Manna/C-DP on $\mathcal{C}_3$) is only interpretable against the clear boundary of what the framework currently covers.

8 Universal scope: CRN-in-field as the underlying object

We have framed the Manna calculation in narrow terms (one universality class, one stratum), but the machinery used here is not Manna-specific. This section makes the wider scope claim explicit so that the Manna result can be read as one specific test of it.

The scope claim. *Any statistical-physics system whose dynamics admits a Doi–Peliti / MSR action representation — a chemical reaction network embedded in a (possibly non-equilibrium) field, with retarded propagators where appropriate — falls within the CFAC analytic framework.* The two-step procedure is: (i) write the CRN-in-field representation of the system; (ii) run the CFAC pipeline (filter diagrams by causal contraction and bridge factorisation, compute Symanzik polynomials, apply BPHZ multiplicities $K_G|_{\alpha=1}$, sector-decompose, and extract the residual list of free integrals).

Demonstrated reach. The CFAC pipeline as it stands has been exercised across a heterogeneous list of systems:

- **LDWC depinning:** 2-loop FRG for an elastic manifold, multi-field MSR action, doubled-propagator hat diagram. (Amarteifio, 2026c)
- **LDWC long-range elasticity:** same pipeline, fractional propagator exponents.
- **LDWC periodic CDW:** same pipeline, $N_{\text{res}} = 0$.
- **Wilson–Fisher $O(N)$ η at 2-loop:** equilibrium ϕ^4 , no FRG. Reproduces $\eta = (N+2)/[2(N+8)^2]\varepsilon^2$.
- **Directed percolation 2-loop:** pure Doi–Peliti. (Amarteifio, 2026d)
- **Manna / C-DP:** this paper (multi-field, conserved, non-equilibrium).

None of these are in the same physics class. What they share is the CRN-in-field structure. This is evidence for the scope claim, not proof.

Routes from common stat-physics to CRN-in-field.

Reaction–diffusion: Direct (Doi–Peliti).

Equilibrium ϕ^4 / $O(N)$: Stochastic quantisation (Parisi–Wu) gives a Langevin equation; its MSR action is a CRN-in-field. The field-theoretic anomalous dimension η falls out of the same pipeline as Manna’s τ .

Spin systems: Glauber dynamics is literally a CRN of spin flips; mapping to a continuous order parameter is standard.

KPZ / surface growth: MSR action for the height field; the noise vertex defines a CRN.

Hydrodynamic fluctuations: MSR for Navier–Stokes; the pressure constraint is a Lagrange-multiplier field; the noise is a CRN.

Lattice gauge (stochastic quantisation): Langevin in field space, MSR action with gauge-fixing constraints.

The technical step in each case is well-known; CFAC inherits the machinery once the action is in standard form.

Where the scope claim genuinely doesn’t reach (yet). The boundary cases of Section 7 sharpen this:

- **Matrix-valued fields** (TAP complexity, RMT, replica $n \rightarrow 0$): the CRN is matrix-valued, and the bridge function in CFAC is currently scalar. A matrix-valued bridge (free probability / R-transforms) is a research extension.
- **Non- D -finite generating functions** (LERW in $d = 3$, lattice Green’s functions): the stratification theorem requires algebraic or D -finite GFs.
- **Topologically constrained systems** (BRST-quantised gauge theory, field theories with global anomalies): the constraint algebra is not currently handled by the pipeline.

Manna’s role in the wider programme. Manna is deliberately a stress test for the multi-field, conserved, non-equilibrium corner of the scope. If CFAC handles Manna successfully (the slotting holds, the rank-3 bridge gives the right order of magnitude, and the conservation projector in the multi-field DSE selects $\mathcal{C}_3$ as predicted), it is direct evidence that the wider scope claim is plausible: *a representative system from a distinct stat-phys corner falls in line with the same machinery*. A separate note in preparation will state the scope claim formally and tabulate the demonstrated cases as a reference artefact.

9 Discussion

Summary of the solid contributions. Three things are established by this paper and the companion (Amarteifio, 2026c):

1. **Stratum classification.** The CFAC codimension argument (Prop. 3) places CDP/Manna on the $\mathcal{C}_3$ Puiseux stratum, giving a skeleton $\tau_0 = 4/3$ within 3% of the measured $\tau \approx 1.29$. This is an organising principle for non-DP universality that does not require knowledge of the underlying field theory.
2. **Cross-check via two independent routes.** The rank-3 skeleton value $4/3 = 1 + 1/k$ equals $\nu_\perp = 1/(2 - \zeta) = 1/(2 - 5/4) = 4/3$ read from the qEW mapping with $\zeta = 5/4$. Two independent derivations — combinatorial stratification and depinning field theory — land on the same rational.

3. **Three of four qEW anchor numbers CFAC-derived.** The LDW mapping (Le Doussal and Wiese, 2015) converts qEW (ζ, z) into the full Manna exponent set. Three of the four 2-loop anchor numbers are now reached end-to-end from CFAC primitives: $\zeta_1 = 1/3$, $z_1 = -2/9$, and $c_\zeta = 0.14331$ (the last to relative error 2×10^{-4} , in the companion paper). Only the 2-loop dynamical coefficient c_z remains external input, with the derivation route flagged.

What is *not* claimed.

- We do not claim a first-principles derivation of the Manna exponents from within CFAC. The qEW mapping is the proper field theory (Le Doussal and Wiese, 2015); CFAC reaches the anchor numbers that feed it.
- We do not claim the $\mathcal{C}_3$ stratum prediction is proven from a microscopic DSE. The conservation Ward identity encoding as a signed projector on the algebraic curve is the open gap flagged in Section 4 and remains so.
- We do not claim a better numerical match to the Manna measurements than the measurements internally support. The published 1+1d Manna exponents imply three different values of z depending on which scaling route is used (spread 14%); our prediction sits inside that spread.

Methodological contribution. The paper documents a wrong-path framing (CDP as DP-plus-small correction) that produced apparent quantitative success through ratio cancellation in hyperscaling. The deprecated implementation is retained in the repository with a regression test that would flag any accidental reintroduction. The joint-exponent consistency rule proposed in Section 4 is the explicit safeguard.

Next steps. (i) Port the LDWC $X^{(\alpha)}$ machinery to compute the short-range qEW dynamical coefficient c_z at 2-loop, closing the last external input. (ii) Encode the conservation Ward identity as a signed projector on the algebraic curve to derive $\mathcal{C}_3$ stratum dominance microscopically. (iii) Test the slotting on a further non-DP class not used in the hypothesis formation (parity-conserving branching annihilation random walks, voter class at $k \rightarrow \infty$ with log corrections).

A Related coupled-field systems: ants (TODO)

A natural sibling of CDP/Manna in the CFAC programme is the chemotactic ant colony, treated in (Amarteifio, 2026e) as the canonical coupled-field system: a Doi–Peliti particle sector (ants in two states $S \rightleftharpoons R$) coupled to an MSR continuous-field sector (pheromone density). The ant construction shares the multi-field, non-equilibrium structure that made CDP/Manna the test case in this paper.

The two systems sit on *different* Puiseux strata, however:

- CDP/Manna: $\mathcal{C}_3$ (skeleton $\tau_0 = 4/3$, this paper).
- Ants: $\mathcal{C}_2$ saddle-node, with order parameter exponent $\beta = 1/2$ verified to 2.7% by simulation ((Amarteifio, 2026e, §4)). Conservation enters differently — the ant “conservation” is the total particle number, which fixes $\langle S+R \rangle$ but does not generate the same Ward identity that promotes CDP from $\mathcal{C}_2$ to $\mathcal{C}_3$.

The two cases together would be a stronger test of the codimension argument (Proposition 3) than either alone:

- **TODO 1.** Apply the `conserved_2.py` (and eventual `conserved_3.py`) machinery to the ant DSE and verify that it returns $\mathcal{C}_2$ dominant, with $\beta = 1/2$ falling out of the rank-2 bridge as in standard DP.
- **TODO 2.** Identify the precise reason the ant conservation does *not* promote to $\mathcal{C}_3$: is it that the ant’s $S + R$ conservation is an exact lattice constraint rather than a Ward identity at the field-theoretic level? Or that the conservation acts on the activity sector itself rather than coupling activity to a separate conserved background?
- **TODO 3.** Construct a modified ant model that adds a *second* conservation law (e.g., a conserved environmental field) and check whether the codimension argument predicts $\mathcal{C}_4$ promotion, observing what numerical β emerges.

These cross-tests would either strengthen the slotting framework (by showing it correctly predicts *which* coupled-field systems sit on which strata) or expose its failure mode (a coupled-field system with conservation that nonetheless does not slot as predicted). Both outcomes are informative. We defer this work to a follow-up note.

References

- Amarteifio, S., 2026. *CFAC theorem: stratification of Dyson–Schwinger equations by Puiseux order*. Percolation Labs preprint.
- Amarteifio, S., 2026. Candidate problems for the CFAC framework. `rdft/docs/problems.md`.
- Auffinger, A., Ben Arous, G., and Černý, J., 2013. Random matrices and complexity of spin glasses. *Comm. Pure Appl. Math.* **66**, 165.
- Bonachela, J. A. and Muñoz, M. A., 2008. Confirming and extending the hypothesis of universality in sandpiles. *Phys. Rev. E* **78**, 041102.
- Bray, A. J. and Moore, M. A., 1980. Metastable states in spin glasses. *J. Phys. C* **13**, L469.
- Rossi, M., Pastor-Satorras, R. and Vespignani, A., 2000. Universality class of absorbing phase transitions with a conserved field. *Phys. Rev. Lett.* **85**, 1803.
- Vespignani, A., Dickman, R., Muñoz, M. A. and Zapperi, S., 1998. Driving, conservation, and absorbing states in sandpiles. *Phys. Rev. Lett.* **81**, 5676.
- Le Doussal, P. and Wiese, K. J., 2015. Exact mapping of the stochastic field theory for Manna sandpiles to interfaces in random media. *Phys. Rev. Lett.* **114**, 110601. arXiv:1410.1930.
- Le Doussal, P. and Wiese, K. J., 2016. Directed percolation with a conserved field and the depinning transition. *Phys. Rev. E* **94**, 042138. arXiv:1604.05556.
- Chauve, P., Le Doussal, P. and Wiese, K. J., 2001. Renormalization of pinned elastic systems: how does it work beyond one loop? *Phys. Rev. Lett.* **86**, 1785. arXiv:cond-mat/0006197.
- Le Doussal, P. and Wiese, K. J., 2002. 2-loop functional renormalization group theory of the depinning transition. arXiv:cond-mat/0205108.
- Rosso, A. and Krauth, W., 2002. Roughness at the depinning threshold for a long-range elastic string. *Phys. Rev. E* **65**, 025101(R).
- Amarteifio, S., 2026. Reading the LDWC two-loop roughness coefficient through the CFAC lens. Percolation Labs preprint (`paper/cfac/ldw_kirchhoff_recovery.tex`).
- Amarteifio, S., 2026. Multi-Loop Feynman Integrals as Counting Problems. Percolation Labs preprint (`paper/two_loop_worked/main.tex`).
- Amarteifio, S., 2026. Ant Colonies as Canonical Coupled-Field Systems: A CFAC Recipe Book. Percolation Labs preprint (`paper/cfac/ac_ants.tex`).
- Corwin, I. and Ghosal, P., 2020. Lower tail of the KPZ equation. *Duke Math. J.* **169**, 1329.
- Crisanti, A. and Leuzzi, L., 2005. Spherical $2+p$ spin-glass model. *Phys. Rev. B* **73**, 014412.
- Henkel, M., Hinrichsen, H. and Lübeck, S., 2008. *Non-Equilibrium Phase Transitions, Vol. 1: Absorbing Phase Transitions*. Springer.
- Krajenbrink, A., 2020. From Painlevé transcendents to the KPZ equation. Ph.D. thesis, Ecole Normale Supérieure, Paris.
- Kozma, G., 2007. The scaling limit of loop-erased random walk in three dimensions. *Acta Math.* **199**, 29.
- Lübeck, S. and Hucht, A., 2002. Absorbing state phase transition in conserved DP. *J. Phys. A* **35**, 4853.
- Ódor, G., 2004. Universality classes in non-equilibrium lattice systems. *Rev. Mod. Phys.* **76**, 663.
- Subag, E., 2017. The complexity of spherical p -spin models. *Ann. Probab.* **45**, 3385.