

Semester Final Review U1-U7

U1 (pg. 14: 1-5, pg. 20: 1-5)

pg. 14

1. a) The main difference between physical properties and chemical properties is that physical properties can be observed without altering the identity of the substance, whereas chemical properties require the identity to be changed.
b) An example of a physical property could be color. Although change in color indicates a chemical change, the color of a substance can be observed without chemically changing the identity of the substance. On the other hand, reactivity is a chemical property because it requires a change in identity.
2. a) Physical, doesn't change identity
b) Physical, doesn't change identity
c) Changes identity when it burns.
3. Indefinite volume and indefinite shape: gas, definite volume but indefinite shape: liquid, definite volume and definite shape: solid.
4. Mixtures are blends of different types of matter. Pure substances must have a fixed composition.
5. Same, sucrose is always the same molecule.

pg. 20

1. Oxygen, Sulfur, Copper, Silver
2. Fe, N, Ca, Hg
3. Elements in a group are most likely to undergo the same kinds of reactions because they share properties that relate to their valence shell electrons.
4. Metals are lustrous, malleable, ductile, conductive, dense, and have high melting and boiling points (usually solids). Nonmetals are brittle, dull, less reactive, and are usually gaseous and have low melting points. Metalloids are lustrous, brittle, with moderate reactivity and melting/boiling points. Metalloids are usually solids.
5. It would usually be more unreactive because it wouldn't be in a bond. You can determine whether or not it's a metal from its position on the periodic table.

U2 (pg. 29: 2-4, pg. 40: 3-6, pg. 55: 1-9)

2. a. Qualitative
b. Qualitative
c. Quantitative
3. Hypotheses are testable statements, and theories are generalizations. Theories usually aren't created before hypotheses.
4. Models help explain hypotheses and theories.

3. a. $10.5\text{g} = 0.0105\text{kg}$
b. $1.57\text{km} = 1570\text{m}$
c. $3.54\mu\text{g} = 0.00000354\text{g}$
d. $3.5\text{mol} = 3500000\mu\text{mol}$
e. $1.2\text{L} = 1200\text{mL}$
f. $358\text{cm}^3 = 0.000358\text{m}^3$
g. $548.6\text{mL} = 548.6\text{cm}^3$
4. a. $1000000\text{cm}^3 / 1\text{m}^3$
b. $2.54\text{cm} / 1\text{in.}$
c. $0.000001\text{g} / 1\mu\text{g}$
d. $1000000\text{m} / 1\text{Mm}$
5. a. $(d=m/v) \rightarrow 1.71\text{g/mL}$
b. $7.75\text{g} = 1.71\text{g/mL} * v, v = 4.53\text{mL}$
6. No, the student is not performing this calculation. Their conversion factor checks out, but mg should be on the top in order to convert from g to mg.

1. Student A's results are more accurate because they are more evenly distributed around 8.94g/cm^3 . Student B's results are more precise because they are closer together.
2. a. 4
b. 2
c. 6
d. 1
e. 4
3. 2.7

4. a. $52.13\text{g} + 1.7502\text{g} = 53.88\text{g}$
 b. $12\text{m} * 6.41\text{m} = 77\text{m}$
 c. $16.25\text{g} / 5.1442\text{mL} = 3.159 \text{ g/mL}$
5. a. $0.0154\text{g} + 0.286\text{g} = 0.301\text{g} = 3.01 * 10^{-1}\text{g}$
 b. $7\,023\,000\,000\text{g} - 66\,200\,000\text{g} = 6\,956\,800\,000\text{g} = 6.9568 * 10^{-9}\text{g}$
 c. $8.99 * 10^{-4}\text{m} * 3.57 * 10^4\text{m} = 32.1\text{m} = 3.21 * 10^1\text{m}$
 d. $2.17 * 10^{-3}\text{g} / (5.002 * 10^4\text{mL}) = 0.4338 * 10^{-7}\text{g/mL} = 4.338 * 10^{-8}\text{mL}$
6. a. $5.6 * 10^5$
 b. $3.34 * 10^4$
 c. $4.120 * 10^{-4}$
7. a. $215.6\text{g} - 110.4\text{g} = 105.2\text{g}$
 b. $105.2\text{g} / 114\text{mL} = 0.923\text{g/mL}$
8. $(m=dv) \rightarrow 19.3\text{g/cm}^3 * 5.0 * 10^{-3}\text{cm}^3 = 0.096\text{g}$
9. Direct variation can be represented by the equation $y=kx$. Inverse variation can be represented by the equation $k=xy$.

U3 (pg. 67: 1-2, pg. 72: 1-3, pg. 82: 1-3, pg. 83: 1-6)

pg. 67

1. i. All matter is composed of extremely small particles called atoms.
 ii. Atoms of an element are identical, and atoms of different elements must differ.
 iii. Atoms cannot be divided, created, or destroyed.
 iv. Atoms combine in simple whole-number ratios.
 v. In chemical reactions, atoms are combined, subdivided, and rearranged.
2. iii explains the law of conservation of mass. ii and iv explain the law of definite proportions. iv and v explain the law of multiple proportions.

pg. 72

1. a. Smallest particle of an element that has the element's properties
 b. Negatively charged subatomic particle in atoms
 c. Center of an atom, has protons and neutrons
 d. Positively charged subatomic particle in atoms
 e. Subatomic particle in an atom without a charge
2. a. Thomson discovered the electron through the cathode ray experiment and came up with the plum pudding model.

- b. Millikan discovered the neutron through his oil drop experiment.
 - c. Rutherford discovered the proton through his gold foil experiment.
3. The protons and neutrons have a mass of 1 and make up the nucleus, the center of the atom. The electrons have an approximate mass of 0 and orbit the nucleus. Protons have a positive 1 charge, electrons have a negative 1 charge, and neutrons don't have a charge.

pg. 82

1. $1.50 \times 10^{12} \text{ atoms Pb} \times \text{mol} / (6.022 \times 10^{23} \text{ atoms}) = 2.49 \times 10^{-12} \text{ mol Pb}$
2. $2500 \text{ atoms Sn} \times \text{mol} / (6.022 \times 10^{23} \text{ atoms}) = 4.2 \times 10^{-23} \text{ mol Sn}$
3. $2.75 \text{ mol Al} \times (6.022 \times 10^{23} \text{ atoms}) / \text{mol} = 1.66 \times 10^{24} \text{ atoms Al}$

pg. 83

1. a. Atomic number - number of protons
 b. Mass number - number of protons and neutrons
 c. Relative atomic mass - atomic mass relative to 1/12 the mass of carbon-12
 d. Average atomic mass - average atomic mass in nature
 e. Mole - as many particles as there are in 12g of carbon-12, aka avogadro's number of particles
 f. Avogadro's number - 6.022×10^{23} - number of particles in one mole of a substance
 g. Molar mass - mass of a mole of a substance
 h. An atom with a different number of neutrons
2. a. Sodium-23 - p+: 11, n0: 12, e-: 11
 b. Calcium-40 - p+: 20, n0: 20, e-: 20
 c. $^{64}_{22}\text{Cu}$ - p+: 22, n0: 42, e-: 22
 d. $^{108}_{47}\text{Ag}$ - p+: 47, n0: 61, e-: 47
3. a. $^{28}_{14}\text{Si}$, silicon-28
 b. $^{56}_{26}\text{Fe}$, iron-56
4. 39.10u, 39.10g
5. a. $2.00 \text{ mol N} \times 14.0067 \text{ g N} / \text{mol N} = 28.0 \text{ g N}$
 b. $3.01 \times 10^{23} \text{ atoms Cl} \times 17 \text{ g Cl} / (6.022 \times 10^{23} \text{ atoms Cl}) = 8.50 \text{ g Cl}$
6. a. $12.15 \text{ g Mg} \times \text{mol Mg} / 24.305 \text{ g Mg} = 0.4999 \text{ mol Mg}$
 b. $1.50 \times 10^{23} \text{ atoms F} \times \text{mol} / (6.022 \times 10^{23} \text{ atoms}) = 0.249 \text{ mol F}$

U4 (pg. 97: 1-5, pg. 104: 1-3, pg. 116: 1-5)

pg. 97

1. A major shortcoming of Rutherford's model was that it didn't explain the distribution of the electrons.
2. $c = \lambda \nu$. c : speed of light, λ : wavelength, ν : frequency.
3. a. Electromagnetic radiation is composed of waves of energy.
 - b. Wavelength is the minimum length of the wave pattern that repeats itself.
 - c. Frequency is wavelengths per second.
 - d. A quantum of energy is the minimum amount of energy gained or lost by an atom.
 - e. A particle of EMR that carries a quantum of energy.
4. Light acts both as a wave and as a particle. In the double slit experiment, light passes straight through one slit like particles would, but when passing through two slits, the particles interfere with each other and act like waves.
5. Bohr's model of the hydrogen atom has a nucleus in the center, and energy levels on the outside. When a photon is gained, the electron moves further away from the nucleus, and when the photon is lost, light is emitted and the electron falls back down.

pg. 104

1. a. Levels on the Bohr model
 - b. Quantum numbers are properties of atoms that relate to their orbitals.
2. a. The four quantum numbers are
 - n : Principal quantum number
 - ℓ : Angular momentum quantum number
 - m : Magnetic quantum number
 - m_s : Spin quantum number
3. The principal quantum number indicates the main energy level occupied by the outermost electron.
 - The angular momentum quantum number indicates the type of orbital.
 - The magnetic quantum number indicates the orientation of the orbital.
 - The spin quantum number ($+1/2$ or $-1/2$) indicates the spin of the electron.

pg. 116

1. a. An atom's electron configuration is how the electrons are organized.
 - b. Aufbau principle, Pauli exclusion principle, Hund's rule.
2. Orbital notation, noble-gas notation, and electron configuration notation.
3. An octet of electrons is 8 electrons. Noble gases contain octets of electrons.
4. a. $1s^2 2s^2 2p^2$
 [He] $2s^2 2p^2$

1s: 1↓, 2s: 1↓, 2p: 1 1

b. 1s² 2s² 2p⁶

[He] 2s² 2p⁶

1s: 1↓, 2s: 1↓, 2p: 1↓ 1↓ 1↓

c. 1s² 2s² 2p⁶ 3s² 3s⁴

[Ne] 3s² 3s⁴

1s: 1↓, 2s: 1↓, 2p: 1↓ 1↓ 1↓, 3s: 1↓, 3p: 1↓ 1 1

5. a. Phosphorus

b. Potassium

c. Silicon

d. Selenium

U5 (pg. 129: 1-2, pg. 141: 1-5, pg. 156: 1-3)

pg. 129

1. a. Stanislao Cannizzaro

b. Dmitri Mendeleev

c. Henry Moseley

2. Physical and chemical properties are periodic **functions** of their atomic numbers.

pg. 141

1. s, p, d, and f blocks

2. a. Alkali metals

b. Alkaline earth metals

c. Transition metals

d. Halogens

e. Noble gases

3. The last group can determine the group number.

4. 3d¹⁰ 4s² 4p³

5. Period 4

D-block

Group 9

pg. 156

1. a. Atomic radius decreases across a period and increases down a group. It decreases across a period because it gains protons while the electrons are on the same energy level, pulling the energy level closer. It increases down a group because the increase in the number of energy levels also increases the atomic radius (electrons repel).
- b. First ionization energy increases across a period and decreases down a group. Across a period, the strength of the pull on the farthest electron increases because the electrons are closer to the nucleus. Down a group, it decreases because the electrons are farther away from the nucleus, so they are pulled less.
- c. Electron affinity increases across a period and decreases down a group. Across a period, there is a greater change in energy when electrons are added because more electrons are being added to the same energy level (or new energy level). Down a group, it decreases because the outermost energy level is farther away from the nucleus, so it will be more insignificant.
- d. Ionic radius decreases across a period and increases down a group. It decreases across a period because it gains protons while the electrons are on the same energy level, pulling the energy levels closer. It increases across a group due to the **shielding effect**.
- e. Electronegativity increases across a period, decreases across a group, and is 0 for the noble gases. It increases across a period because as atoms are closer to getting a full octet, they can gain an electron more easily. Noble gases don't have any electronegativity because they already have full octets. Electronegativity decreases across a group because, as the radius increases, it becomes more difficult for the protons in the nucleus to pull the electrons.
2. a. They are more inconsistent.
- b. The reason they are more inconsistent is because they don't all form octets.
3. Group 1: 1, Group 2: 2, Groups 13-18: 8-10

U6 (pg. 167: 1-5, pg. 179: 1-4, pg. 197: 1-6)

pg. 167

1. Covalent bonding is caused by the nuclei of multiple atoms pulling the same electron. Ionic bonding is caused by the difference in ionic charge after the electrons are taken to form an octet.
2. If the electronegativity difference is between 0 and 0.3, it is a nonpolar covalent bond. If it is between 0.3 and 1.7, it is a polar covalent bond. If the difference is greater than 1.7, then it is likely an ionic bond.
3. a. $4.0 - 1.0 = 3.0$, ionic
- b. $2.5 - 1.9 = 0.6$, polar covalent
- c. $2.8 - 2.5 = 0.3$, nonpolar covalent





4. Br & I, Cu & S, Li & F.
5. a. Cu & Cl would have the greater ionic character because the electronegativity difference is greater.
- b. Chlorine has the greater negative charge in Cu & Cl because it is better at pulling electron(s) from copper.


pg. 179

1. a. Bond length is the average distance between two bonded atoms.
- b. Bond energy is the energy required to break a bond.
2. Chemical compounds will generally try to form an octet of electrons in its highest occupied energy level

3. a. 1
- b. 2
- c. 3

4.





pg. 197


1. VSEPR theory

2.


a) Linear



b) Tetrahedral



c) Trigonal planar



3. Lone pairs and bonds on the central atom.

4. We didn't do hybridization.

5. Hydrogen bond forces are the strongest intermolecular forces and contribute to the high boiling point of water.

6. Boiling points can be used to test the strength of intermolecular forces. When something is boiling, the pulling forces between the molecules have been overcome.

U7 (pg. 219: 2-4, pg. 223: 1, pg. 232: 1-6, pg. 237: 1-4)

pg. 219

2. a. AlBr_3
 b. Na_2O
 c. MgI_2
 d. PbO
 e. SnI_2
 f. Fe_2S_3
 g. $\text{Cu}(\text{NO}_3)_2$
 h. $(\text{NH}_4)_2\text{SO}_4$
3. a. Sodium iodide
 b. Magnesium sulfide
 c. Calcium oxide
 d. Potassium sulfide
 e. Copper(I) bromide
 f. Iron(II) chloride
4. a. NaOH
 b. $\text{Pb}(\text{NO}_3)_2$
 c. FeSO_4
 d. P_2O_3
 e. CSe_2
 f. $\text{HC}_2\text{H}_3\text{O}_2$
 g. HClO_3
 h. H_2SO_3

pg. 223

1. We did not do oxidation numbers.

pg. 232

1. formula mass: $2 \cdot (14.007\text{u N}) + 8 \cdot (1.008\text{u H}) + (12.011\text{u C}) + 3 \cdot (15.999\text{u O}) = 96.086\text{u}$ $(\text{NH}_4)_2\text{CO}_3$
 molar mass: 96.086g $(\text{NH}_4)_2\text{CO}_3$
2. 2 mol N
 8 mol H
 1 mol C
 3 mol O
3. formula mass: $2 \cdot (55.845\text{u Fe}) + 3 \cdot (32.06\text{u S}) + 12 \cdot (15.999\text{u O}) = 399.86\text{u}$ $\text{Fe}_2(\text{SO}_4)_3$
 $3.25\text{mol Fe}_2(\text{SO}_4)_3 \cdot 399.86\text{u Fe}_2(\text{SO}_4)_3 = 1299.54\text{g Fe}_2(\text{SO}_4)_3$

4. formula mass: $9 \times (12.011 \text{ u C}) + 8 \times (1.008 \text{ u H}) + 4 \times (15.999 \text{ u O}) = 180.159 \text{ u C}_9\text{H}_8\text{O}_4$
 $100.0 \text{ mg C}_9\text{H}_8\text{O}_4 \times 1 \text{ g} / 1000 \text{ mg} / (180.159 \text{ g/mol C}_9\text{H}_8\text{O}_4) \times 6.022 \times 10^{23} \text{ molecules/mol} = 3.343 \times 10^{20} \text{ molecules aspirin}$
5. formula mass: $2 \times (14.007 \text{ u N}) + 8 \times (1.008 \text{ u H}) + (12.011 \text{ u C}) + 3 \times (15.999 \text{ u O}) = 96.086 \text{ u (NH}_4)_2\text{CO}_3$
 $2 \times (14.007 \text{ u N}) / 96.086 \text{ u (NH}_4)_2\text{CO}_3 = 29.155\% \text{ N}$
 $8 \times (1.008 \text{ u H}) / 96.086 \text{ u (NH}_4)_2\text{CO}_3 = 8.392\% \text{ H}$
 $(12.011 \text{ u C}) / 96.086 \text{ u (NH}_4)_2\text{CO}_3 = 12.500\% \text{ C}$
 $3 \times (15.999 \text{ u O}) / 96.086 \text{ u (NH}_4)_2\text{CO}_3 = 49.952\% \text{ O}$
6. CuSO_4 formula mass: $(63.55 \text{ u Cu}) + (32.06 \text{ u S}) + 4 \times (16.00 \text{ u O}) = 159.61 \text{ u CuSO}_4$
 H_2O formula mass: $2 \times (1.01 \text{ u H}) + (16.00 \text{ u O}) = 18.02 \text{ u H}_2\text{O}$
 $n = (58.55 / 18.02) / (103.75 \text{ u CuSO}_4 / 159.60 \text{ u CuSO}_4) \approx 5$
 $5.0 \text{ moles H}_2\text{O}$

pg. 237

1. $36.48\% \text{ Na} \times \text{mol} / 22.99 \text{ g} = 0.01587 \text{ mol Na}$
 $25.41\% \text{ S} \times \text{mol} / 32.06 \text{ g} = 0.007926 \text{ mol S}$
 $38.11\% \text{ O} \times \text{mol} / 16.00 \text{ g} = 0.02382 \text{ mol O}$
 Na_2SO_3
2. $53.70\% \text{ Fe} \times \text{mol} / 55.845 \text{ g} = 0.009616 \text{ mol Fe}$
 $46.30\% \text{ S} \times \text{mol} / 32.06 \text{ g} = 0.01444 \text{ mol S}$
 $n \approx 1.5$
 Fe_2S_3
3. $1.04 \text{ g K} \times (\text{mol K} / 39.098 \text{ g K}) = 0.0266 \text{ mol K}$
 $0.70 \text{ g Cr} \times (\text{mol Cr} / 51.996 \text{ g Cr}) = 0.013 \text{ mol Cr}$
 $0.86 \text{ g O} \times (\text{mol O} / 16.00 \text{ g O}) = 0.054 \text{ mol O}$
 K_2CrO_4
4. $4.04 \text{ g N} \times (\text{mol N} / 14.01 \text{ g N}) = 0.288 \text{ mol N}$
 $11.46 \text{ g O} \times (\text{mol O} / 16.00 \text{ g O}) = 0.7162 \text{ mol O}$
 N_2O_5
empirical formula mass: $2 \times 14.01 \text{ g N} + 5 \times 16.00 \text{ g O} = 108.02 \text{ g N}_2\text{O}_5$
 $n = 108.0 \text{ g} / 108.02 \text{ g} = 1$
 N_2O_5