




Ex. 3. - res:

$$\begin{aligned} \text{The } Q_2 &= (a_1, a_2) \\ (a_1, a_2) &\subseteq A^2 \Rightarrow \\ \Rightarrow R_X &= \mathcal{V}(Q) = Q_2 \\ Q_2 &= Q_1 \end{aligned}$$



$$\begin{aligned} Q_2 &= Q_1 \\ (X, Y) &\in A^2 \\ (X, Y) &\in R_X \\ (X, Y) &\in R_X \end{aligned}$$





$\partial U \partial^2$

$$\begin{aligned} & \partial \partial \cdot \\ & (X \mid z) \in A \\ & (X \mid z) \in \partial \partial \cdot \\ & (X \mid z) \in \partial \partial \cdot \end{aligned}$$

$$\Rightarrow \partial \partial \cdot \text{ have } \cdot$$

$$\Rightarrow \partial \partial \cdot = \partial \partial \cdot$$

$$\partial \partial \cdot A \rightarrow A \quad \partial \partial \cdot A \rightarrow A$$

$$\Rightarrow \mathcal{A} \neq \mathcal{B} \times \mathcal{C} \quad \mathcal{A} \neq \mathcal{B} \times \mathcal{C}$$

$$= \mathcal{A} \rightarrow \mathcal{B} \times \mathcal{C}.$$