


Ex. 3. - RES:

Let $Q_1 = (q_1), (q_2)$

$(q_2) \cup A^2 \Rightarrow$


$\Rightarrow R_X = Q_1(Q) = Q_2$



$Q_2 = Q_0 Q$

$(X^2) \in A^2$

$(X^2) \in A^2$



$\partial U \partial^2$

$$\begin{aligned}
 & \partial \partial^2 \cdot \\
 & (X \cap \partial^2) \cap \partial^2 \in A \\
 & (X \cap \partial^2) \cap \partial^2 \in \partial U \partial^2 \\
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2 \\
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2
 \end{aligned}$$

$$\begin{aligned}
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2 \\
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2 \\
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2
 \end{aligned}$$

$$\begin{aligned}
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2 \\
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2 \\
 & \Rightarrow \partial U \partial^2 \cap \partial^2 \in \partial U \partial^2
 \end{aligned}$$

$$f(x) = y$$

$$\Delta \rightarrow R(x, y)$$

$$= H(R(x, y))$$

$$H(R(x, y)) = R$$

$$H(y) = \Delta$$

$$H(x) = \Delta$$

$$(x, y) = (y, x) \in R \Rightarrow$$

$$\Rightarrow \chi(R(x, y)) = 0 \in \mathbb{Z}_2$$

$$\chi(x) = y \rightarrow R(x, y) =$$

$$\Rightarrow \chi(x) = y \rightarrow R(x, y) = 0 = 0$$

$$\wedge \neq 0, \text{ deci.}$$

$$\Rightarrow \| \phi \|_A =$$

