

Practicum 1

Peter Claussen

Ben Derenge

Stephanie Liebl

9/16/2021

Overview

We have been given an Excel data file with data collected for a study from a call center. These data include health metrics from **employees** gathered over an eight month period.

Variables that are of particular interest have been highlighted in the original data. These include variables relating to weight change and demographics. Specifically, we will consider in this preliminary proposal the variables

Highlighted in yellow

- `shift`
- `Total_Met_Min`

Highlighted in orange

- `gender`
- `Age`
- `height`
- `weightgain`
- `lbs_gained`
- `BMI`
- `Vig.ex.Time`
- `Mod.ex.time`
- `Walk.ex.Time`

We will use the convention that text rendered in **sans serif** font denote variable or column names found in the original data set, or derived variables calculated from columns in the original data, while *italics* will denote real-world processes or phenomena of interest. Thus, **weightgain** denotes the data column in the original data file, while *weight gain* denotes some measure of the change in weight by individuals over the study period.

. We have been tasked to provide an analysis to address two specific aims:

- **(SA1)** Does *total metabolic minutes* have an effect on *weight gain*?
- **(SA2)** Does *shift* have an effect on *weight gain*?

We will refer to as **SA1** and **SA2** in further discussion. We address *total metabolic minutes*, *weight gain* and *shift* in the following sections.

Total MET minutes

Total MET Minutes is a composite measure obtained from survey responses (International Physical Activity Questionnaire (IPAQ) short form) to estimate overall physical activity. Physical activity is divided into categories described as *vigorous*, *moderate* and *walking*.

Total MET Minutes can be calculated from 3 data columns. We are given the formula

$$\text{Total_met_min} = 8 \cdot \text{Vig_ex_time} + 4 \cdot \text{Mod_ex_time} + 3.3 \cdot \text{Walk_ex_time}$$

to calculate *Total MET Minutes* from the data.

`Total_met_min` contains many missing values. We will create a new data column, `CalcTMM` that is calculated from existing data columns as given in the formula above. To visualize the number of missing values, we plot missing values as 0 in the following figure.




Figure 1: Total MET Minutes calculated from data, plotted against the data column title `Total_Met_Min`

We can see that the original data column `Total_met_min` has a large number of missing values. If we use this data column as given, we will reduce the number of observations in the analysis. This graph also confirms that the formula given above was used to compute `Total_met_min` from the data.

Shift

Shift takes values of the form 7am, 8am, ..., 2pm, other. There are some columns that have missing values for *shift*. We propose that *shift* be modeled as an ordinal data type, with missing values grouped with *other* and *other* takes an ordinal value greater than 2pm.

```
metrics.dat$shift[metrics.dat$shift==''] <- 'other'
shift.levels <- c(paste(c(7:11), 'am', sep=''), paste(c(12, 1:2), 'pm', sep=''), 'other')
metrics.dat$shift <- factor(metrics.dat$shift, shift.levels)
summary(metrics.dat$shift)
```

##	7am	8am	9am	10am	11am	12pm	1pm	2pm	other
##	31	115	56	50	44	14	8	15	19

Weight Gain

While the data include a variable (data column) named **weightgain**, we are asked to consider other response variables, including change in weight (**lbs_gained**) and change in BMI. Thus, our first task is to determine the appropriate response variable. The choice of response variable will dictate both choice of statistical method (i.e. logistic regression vs linear regression) and methods for data cleaning.

weightgain (Binomial Response)




Figure 2: **weightgain** versus Total MET Minutes (calculated) and Shift

At first glance, this appears to be a logistic regression - the response variable (**weightgain**) should be binary (Yes/No), ignoring a small number of missing values represented as light gray blocks in the plot. However, we are asked to consider alternate response variables (BMI, pounds gained) as markers for the general response "weight gain", as opposed to the specific data variable 'weightgain'


Pounds gained (Continuous Response)

The **pounds_gained** column contains only non-negative integer values and missing values. This implies every known observation has gained weight. Checking corresponding values in the **weightgain** column confirms all the NA in **pounds_gained** are "No" in **weightgain**. This will affect the interpretation of SA.1 and SA.2. To use **pounds_gained** as the response variable, we would be ignoring all observations who lost weight.

Plots of Pounds_Gained


120 missing rows and 1 row zero need to be excluded for the following plots.

Calculated TMM vs Pounds Gained



The bar plot above suggest shift does have potential to predict pounds_gained. The column 'pounds_gained' seems to decrease on average as shifts move later into the day, with an exception for the 2pm shift. If we don't classify Missing shifts into the "other" category, the missing group has the highest gained weight of all the shifts. As the plot shows some relation around adjacent shifts, we may rank the shifts as opposed to treating shifts like independent factors.

Targeting Binary outcome with weightgain




weightgain


If desired, `weightgain` can be predicted from `Total_met_minutes`, as most of the “No”s from `weightgain` have NAs or zeros from `total_met_minutes`. This might change if we predict the missing TMM values using the equation from earlier.

Change in BMI


BMI at the end of the study period has been recorded, while initial BMI has not. We can, however, calculate an initial BMI from other data columns. We calculated the initial BMI using the following calculation:
`initial_BMI=bweight/(height^2) * 703`



In the scatter plot visualizing (`initial_BMI`) versus final BMI (`'BMI'`), it appears that there is a strong linear relationship between the variables. The scatter plot also suggests that (`metrics.dat$BMI`) is frequently greater than (`metrics.dat$initial_BMI`).




To look at the difference between the initial BMI and the final BMI for call center employees, we created a new column (`BMI_diff`) by subtracting the initial BMI from the final BMI. Thus, in this column, positive values represent increases in BMI over the eight month period.




The BMI calculation is dependent on whether weight is a measure of pounds or kilograms. To ensure that the BMI values in the given dataset are correct, we manually calculated the BMI using the following formula: $\text{bweight}/(\text{height}^2)*703$. We then plotted our calculated BMI against the given BMI with a $y=x$ line to verify the values are all the same.

```
# Were the BMI calculations correct?
# BMI = (weight/height^2)*703
CalcBMI <- ''
CalcBMI <- ((metrics.dat$bweight)/(metrics.dat$height)^2)*703

plot(CalcBMI, metrics.dat$BMI)
abline(0,1,col='red')
```



If BMI is chosen to be the response variable, two relationships to be examined are `shift` and `BMI_diff` as well as `CalcTMM` and `BMI_diff`.



```
## Warning in `[<-.factor`(`*tmp*`, metrics.dat$weightgain == "", value =
## structure(c(2L, : invalid factor level, NA generated
## [1] "Female" "Male" "Missing"
## [1] No Yes <NA>
## Levels: No Yes
```

Points for consideration

Selection of response variable.

We have proposed three response variables of interest as surrogates for weight gain, specifically `weightgain`, `pounds_gained`, and `initial_BMI`.

Pounds Gained

The skew of `pounds_gained` should be addressed. If this is our response variable, we would consider either data transformation, or a generalized linear model of the poisson family.

Using `Pounds_gained` would result in 121 fewer responses.

The box-whisker plot `pounds_gained` vs `shift` shows a curvilinear response. We should be able to create a straightforward model predicting `weight_gained` from `shift`. The relationship between `weight_gained` and `Calculated Total Met Minutes` would require further exploratory analysis, as any relationship is not apparent from the scatterplot.

BMI (Difference and Initial)

Similar to the variable `lbs_gained`, the box-whisker plot of `shift` vs `BMI_diff` shows a relationship between the variables. However, this relationship is not as skewed nor as linear as that of `shift` and `lbs_gained`. The scatter plot of `CalcTMM` vs `BMI_diff` also looks similar to `CalcTMM` vs `lbs_gained`, however the relationship between `CalcTMM` vs `BMI_diff` appears less linear.

Given our findings, our preference is to use the initial BMI as a predictor variable rather than response for either `lbs_gained` or `weightgain`. We would like feedback from the client as to which is preferred moving forward.


Additional predictor variables.

SA1 and **SA2** include only one predictor variable each (`Total_Met_Min` and `shift`, respectively). This implies a simple regression of one-way ANOVA analysis. However, the data includes other demographic variables that may be confounded with the primary (“yellow”) predictors

- Does the client wish us to perform multiple regression and variable selection with the “orange” highlighted variables?
- Does the client wish us to perform multiple regression and variable selection with any additional variables in the original data file?

We include, for further discussion, a pairs plot of the variables described in this document:

```
## Warning in `[<-.factor`(`*tmp*`, thisvar, value = 0): invalid factor level, NA
## generated
```



This provides a visualization of the scope of the variable selection problem, if restricted to the highlight variables and additional variables described in this document. We also provide as an addendum a summary of the original data and calculated columns, as used in this document.

##	shift	Total_Met_Min	gender	Age	height
##	8am :115	Min. : 0.0	Female :248	Min. : 0.00	Min. : 0.00
##	9am : 56	1st Qu.: 0.0	Male : 99	1st Qu.:25.00	1st Qu.:63.00
##	10am : 50	Median : 357.8	Missing: 5	Median :30.00	Median :66.00
##	11am : 44	Mean : 797.4		Mean :30.88	Mean :62.67
##	7am : 31	3rd Qu.:1315.5		3rd Qu.:39.25	3rd Qu.:69.00
##	other : 19	Max. :5542.0		Max. :64.00	Max. :82.00
##	(Other): 37				
##	weightgain	lbs_gained	BMI	Vig.ex.Time	Mod.ex.time
##	No :111	Min. : 0.00	Min. : 0.00	Min. : 0.00	Min. : 0.00
##	Yes :237	1st Qu.: 0.00	1st Qu.: 0.00	1st Qu.: 0.00	1st Qu.: 0.00

```
## NA's: 4 Median : 7.75 Median :24.12 Median : 27.00 Median : 30.00
## Mean :11.05 Mean :20.02 Mean : 76.16 Mean : 73.64
## 3rd Qu.:16.25 3rd Qu.:29.46 3rd Qu.:120.00 3rd Qu.: 90.00
## Max. :70.00 Max. :48.25 Max. :840.00 Max. :1680.00
##
## Walk.ex.Time initial_BMI BMI_diff CalcTMM
## Min. : 0.00 Min. : 0.00 Min. : 0.000 Min. : 0
## 1st Qu.: 9.75 1st Qu.: 0.00 1st Qu.: 0.000 1st Qu.: 263
## Median : 60.00 Median : 0.00 Median : 0.000 Median : 822
## Mean : 122.92 Mean :12.36 Mean : 1.311 Mean : 1302
## 3rd Qu.: 136.25 3rd Qu.:24.66 3rd Qu.: 2.107 3rd Qu.: 1746
## Max. :1680.00 Max. :44.93 Max. :12.014 Max. :12852
##
```

Finally, a brief note about typesetting. This document was produced in RMarkdown. The original .Rmd with R code and additional details of our preliminary analysis is available upon request.

Analysis

(SA1) Does *total metabolic minutes* have an effect on *weight gain*?

```
metrics.dat$WG <- FALSE
metrics.dat$WG[metrics.dat$weightgain=='Yes'] <- TRUE
SA1.model1 <- glm(WG ~ Total_Met_Min, data=metrics.dat, family = binomial)
summary(SA1.model1)
```

```
##
## Call:
## glm(formula = WG ~ Total_Met_Min, family = binomial, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -1.6816  -1.4654   0.8735   0.9115   0.9145
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  6.556e-01  1.419e-01  4.620 3.83e-06 ***
## Total_Met_Min 8.653e-05  1.113e-04  0.777   0.437
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##      Null deviance: 444.80  on 351  degrees of freedom
## Residual deviance: 444.18  on 350  degrees of freedom
## AIC: 448.18
##
## Number of Fisher Scoring iterations: 4
```

```
SA1.model2 <- glm(WG ~ gender + Age + height + initial_BMI + shift + Total_Met_Min, data=metrics.dat, family = binomial)
```

```
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
```

```
summary(SA1.model2)
```

```
##
## Call:
## glm(formula = WG ~ gender + Age + height + initial_BMI + shift +
##      Total_Met_Min, family = binomial, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -1.51652  -0.65045   0.00000   0.00007   1.89746
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   0.5782398  0.7299439   0.792  0.42826
## genderMale    -1.1273546  0.4278148  -2.635  0.00841 **
## genderMissing -1.4887416  1.3192900  -1.128  0.25913
## Age           -0.0099970  0.0124288  -0.804  0.42120
## height        -0.0097402  0.0084465  -1.153  0.24884
## initial_BMI    1.0378329 53.6633437   0.019  0.98457
## shift8am       0.2432249  0.5848390   0.416  0.67750
## shift9am       0.0896331  0.6419207   0.140  0.88895
## shift10am      0.1754489  0.6896795   0.254  0.79919
## shift11am     -0.1743358  0.6829998  -0.255  0.79853
## shift12pm     -0.6585172  1.2429023  -0.530  0.59624
## shift1pm      -0.6599904  1.2696066  -0.520  0.60318
## shift2pm       1.3148289  0.9624159   1.366  0.17188
## shifttother    0.2566556  0.9969791   0.257  0.79684
## Total_Met_Min -0.0001585  0.0001730  -0.916  0.35941
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##      Null deviance: 444.80  on 351  degrees of freedom
## Residual deviance: 222.29  on 337  degrees of freedom
## AIC: 252.29
##
## Number of Fisher Scoring iterations: 19
```

```
SA1.model3 <- glm(WG ~ gender + Age + height + initial_BMI + shift + Vig.ex.Time + Mod.ex.time + Walk.
```

```
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
```

```
summary(SA1.model3)
```

```
##
## Call:
## glm(formula = WG ~ gender + Age + height + initial_BMI + shift +
##      Vig.ex.Time + Mod.ex.time + Walk.ex.Time, family = binomial,
##      data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -1.61831  -0.64716   0.00000   0.00007   1.97001
##
```

```

## Coefficients:
##               Estimate Std. Error z value Pr(>|z|)
## (Intercept)   0.7399115  0.7474675   0.990  0.32223
## genderMale    -1.1322070  0.4350468  -2.602  0.00925 **
## genderMissing -1.3287561  1.3424536  -0.990  0.32227
## Age           -0.0125228  0.0126939  -0.987  0.32388
## height        -0.0082960  0.0085646  -0.969  0.33273
## initial_BMI    1.0388815 52.7797311   0.020  0.98430
## shift8am       0.2209635  0.5913108   0.374  0.70864
## shift9am       0.0165692  0.6444526   0.026  0.97949
## shift10am      0.2603850  0.7060986   0.369  0.71230
## shift11am     -0.2877916  0.6977417  -0.412  0.68000
## shift12pm     -0.7274323  1.2609414  -0.577  0.56401
## shift1pm      -0.8133505  1.2766194  -0.637  0.52405
## shift2pm       1.3022479  0.9987646   1.304  0.19228
## shifttother    0.2162509  1.0008318   0.216  0.82893
## Vig.ex.Time   -0.0029012  0.0019167  -1.514  0.13012
## Mod.ex.time   -0.0014295  0.0020003  -0.715  0.47484
## Walk.ex.Time   0.0003392  0.0008589   0.395  0.69289
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##      Null deviance: 444.80  on 351  degrees of freedom
## Residual deviance: 218.72  on 335  degrees of freedom
## AIC: 252.72
##
## Number of Fisher Scoring iterations: 19

```

```

library(MASS)
best.model <- stepAIC(SA1.model3, direction = "both")

```

```

## Start:  AIC=252.72
## WG ~ gender + Age + height + initial_BMI + shift + Vig.ex.Time +
##      Mod.ex.time + Walk.ex.Time

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

##
##      Df Deviance    AIC
## - shift      8  223.13 241.13
## - Walk.ex.Time 1  218.87 250.87
## - Mod.ex.time  1  219.26 251.26
## - height      1  219.66 251.66

```

```

## - Age          1    219.70 251.70
## <none>          1    218.72 252.72
## - Vig.ex.Time  1    221.25 253.25
## - gender        2    227.26 257.26
## - initial_BMI   1    429.73 461.73

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
##
## Step:  AIC=241.13
## WG ~ gender + Age + height + initial_BMI + Vig.ex.Time + Mod.ex.time +
##       Walk.ex.Time

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

##           Df Deviance    AIC
## - Walk.ex.Time  1    223.24 239.24
## - Mod.ex.time   1    224.13 240.13
## - height        1    224.21 240.21
## - Age           1    224.33 240.33
## - Vig.ex.Time   1    224.76 240.76
## <none>          1    223.13 241.13
## - gender        2    230.92 244.92
## + shift         8    218.72 252.72
## - initial_BMI   1    437.35 453.35

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
##
## Step:  AIC=239.24
## WG ~ gender + Age + height + initial_BMI + Vig.ex.Time + Mod.ex.time

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

```

```

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

##           Df Deviance    AIC
## - Mod.ex.time 1    224.13 238.13
## - height       1    224.39 238.39
## - Age          1    224.39 238.39
## - Vig.ex.Time  1    224.85 238.85
## <none>         223.24 239.24
## + Walk.ex.Time 1    223.13 241.13
## - gender       2    230.95 242.95
## + shift        8    218.87 250.87
## - initial_BMI  1    437.41 451.41

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

##
## Step:  AIC=238.13
## WG ~ gender + Age + height + initial_BMI + Vig.ex.Time

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

##           Df Deviance    AIC
## - Age        1    225.08 237.08
## - height     1    225.54 237.54
## <none>       224.13 238.13
## - Vig.ex.Time 1    227.00 239.00
## + Mod.ex.time 1    223.24 239.24
## + Walk.ex.Time 1    224.13 240.13
## - gender     2    231.65 241.65
## + shift      8    219.29 249.29
## - initial_BMI 1    437.51 449.51

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

##
## Step:  AIC=237.08
## WG ~ gender + height + initial_BMI + Vig.ex.Time

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

```

```
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
```

```
##           Df Deviance    AIC
## <none>           225.08 237.08
## - Vig.ex.Time    1    227.83 237.83
## - height         1    228.09 238.09
## + Age            1    224.13 238.13
## + Mod.ex.time    1    224.39 238.39
## + Walk.ex.Time   1    225.08 239.08
## - gender         2    232.14 240.14
## + shift          8    220.06 248.06
## - initial_BMI    1    437.54 447.54
```

```
summary(best.model)
```

```
##
## Call:
## glm(formula = WG ~ gender + height + initial_BMI + Vig.ex.Time,
##      family = binomial, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -1.46600  -0.69238   0.00000   0.00008   1.84845
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   0.656867   0.492878   1.333   0.1826
## genderMale    -0.966429   0.404680  -2.388   0.0169 *
## genderMissing -1.231707   1.256998  -0.980   0.3271
## height        -0.013160   0.007628  -1.725   0.0845 .
## initial_BMI    1.031397  53.955748   0.019   0.9847
## Vig.ex.Time   -0.002617   0.001710  -1.530   0.1260
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##      Null deviance: 444.80  on 351  degrees of freedom
## Residual deviance: 225.08  on 346  degrees of freedom
## AIC: 237.08
##
## Number of Fisher Scoring iterations: 19
```



```
metrics.dat$LBS <- round(metrics.dat$pounds_gained)
SA1.model1 <- glm(LBS ~ Total_Met_Min, data=metrics.dat, family = poisson)
summary(SA1.model1)
```

```
##
## Call:
## glm(formula = LBS ~ Total_Met_Min, family = poisson, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -5.205  -4.618  -1.088   1.371  12.032
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  2.367e+00  2.031e-02 116.552 < 2e-16 ***
## Total_Met_Min 4.318e-05  1.471e-05   2.934  0.00334 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 5090.1  on 351  degrees of freedom
## Residual deviance: 5081.7  on 350  degrees of freedom
## AIC: 6111.3
##
## Number of Fisher Scoring iterations: 5
```

```
SA1.model2 <- glm(LBS ~ gender + Age + height+ initial_BMI + shift + Total_Met_Min, data=metrics.dat, family = poisson)
summary(SA1.model2)
```

```
##
## Call:
## glm(formula = LBS ~ gender + Age + height + initial_BMI + shift +
##      Total_Met_Min, family = poisson, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -4.769  -3.175  -1.672   1.033  10.920
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  2.215e+00  9.792e-02 22.624 < 2e-16 ***
## genderMale   -1.061e-01  3.754e-02 -2.827  0.00470 **
## genderMissing 4.747e-02  1.506e-01  0.315  0.75264
## Age         -6.690e-03  1.386e-03 -4.825 1.40e-06 ***
## height      -2.445e-03  1.345e-03 -1.818  0.06910 .
## initial_BMI  4.263e-02  1.346e-03 31.672 < 2e-16 ***
## shift8am     8.326e-02  6.017e-02  1.384  0.16645
## shift9am    -1.629e-01  6.964e-02 -2.339  0.01936 *
## shift10am   -3.238e-01  7.077e-02 -4.575 4.76e-06 ***
## shift11am   -4.464e-01  7.893e-02 -5.656 1.55e-08 ***
## shift12pm   -7.710e-01  1.161e-01 -6.640 3.14e-11 ***
## shift1pm    -9.990e-01  1.775e-01 -5.630 1.80e-08 ***
## shift2pm     2.307e-01  8.480e-02  2.721  0.00651 **
```

```

## shiftother      8.895e-02  8.324e-02   1.069  0.28525
## Total_Met_Min  1.209e-06  1.571e-05   0.077  0.93863
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 5090.1  on 351  degrees of freedom
## Residual deviance: 3698.8  on 337  degrees of freedom
## AIC: 4754.4
##
## Number of Fisher Scoring iterations: 6

SA1.model3 <- glm(LBS ~ gender + Age + height+ initial_BMI + shift + Vig.ex.Time + Mod.ex.time + Walk.
summary(SA1.model3)

##
## Call:
## glm(formula = LBS ~ gender + Age + height + initial_BMI + shift +
##      Vig.ex.Time + Mod.ex.time + Walk.ex.Time, family = poisson,
##      data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -4.9014  -3.1527  -1.6651   0.9505  10.8537
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   2.186e+00  9.861e-02  22.168 < 2e-16 ***
## genderMale    -8.833e-02  3.762e-02  -2.348 0.018890 *
## genderMissing  2.027e-01  1.536e-01   1.320 0.186795
## Age          -7.011e-03  1.390e-03  -5.045 4.55e-07 ***
## height       -1.980e-03  1.361e-03  -1.454 0.145924
## initial_BMI    4.282e-02  1.352e-03  31.673 < 2e-16 ***
## shift8am       9.156e-02  6.039e-02   1.516 0.129524
## shift9am      -2.026e-01  7.001e-02  -2.894 0.003809 **
## shift10am     -3.064e-01  7.103e-02  -4.314 1.60e-05 ***
## shift11am     -4.253e-01  7.931e-02  -5.363 8.20e-08 ***
## shift12pm     -7.959e-01  1.163e-01  -6.845 7.64e-12 ***
## shift1pm      -9.922e-01  1.775e-01  -5.590 2.27e-08 ***
## shift2pm       2.381e-01  8.540e-02   2.788 0.005306 **
## shiftother     9.449e-02  8.335e-02   1.134 0.256945
## Vig.ex.Time   -6.168e-04  1.643e-04  -3.754 0.000174 ***
## Mod.ex.time    6.107e-05  1.090e-04   0.560 0.575149
## Walk.ex.Time   3.179e-04  7.458e-05   4.263 2.02e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 5090.1  on 351  degrees of freedom
## Residual deviance: 3673.8  on 335  degrees of freedom
## AIC: 4733.5
##
## Number of Fisher Scoring iterations: 6

```

```
best.model <- stepAIC(SA1.model3, direction = "both")
```

```
## Start: AIC=4733.47
## LBS ~ gender + Age + height + initial_BMI + shift + Vig.ex.Time +
##      Mod.ex.time + Walk.ex.Time
##
##           Df Deviance    AIC
## - Mod.ex.time  1   3674.1 4731.8
## <none>                3673.8 4733.5
## - height       1   3675.9 4733.5
## - gender       2   3681.2 4736.8
## - Vig.ex.Time  1   3688.3 4746.0
## - Walk.ex.Time 1   3690.7 4748.3
## - Age         1   3699.4 4757.0
## - shift       8   3923.4 4967.0
## - initial_BMI 1   4770.8 5828.5
##
```


```
## Step: AIC=4731.78
## LBS ~ gender + Age + height + initial_BMI + shift + Vig.ex.Time +
##      Walk.ex.Time
##
##           Df Deviance    AIC
## <none>                3674.1 4731.8
## - height       1   3676.2 4731.8
## + Mod.ex.time  1   3673.8 4733.5
## - gender       2   3681.4 4735.0
## - Vig.ex.Time  1   3688.5 4744.1
## - Walk.ex.Time 1   3691.5 4747.1
## - Age         1   3699.8 4755.4
## - shift       8   3926.0 4967.6
## - initial_BMI 1   4771.0 5826.6
```

```
summary(best.model)
```

```
##
## Call:
## glm(formula = LBS ~ gender + Age + height + initial_BMI + shift +
##      Vig.ex.Time + Walk.ex.Time, family = poisson, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -4.9179  -3.1527  -1.6689   0.9571  10.8533
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  2.191e+00  9.817e-02  22.317 < 2e-16 ***
## genderMale   -8.830e-02  3.762e-02  -2.347 0.018913 *
## genderMissing 1.971e-01  1.532e-01   1.286 0.198348
## Age        -7.022e-03  1.389e-03  -5.054 4.33e-07 ***
## height     -1.966e-03  1.360e-03  -1.446 0.148273
## initial_BMI  4.278e-02  1.349e-03  31.703 < 2e-16 ***
## shift8am     8.850e-02  6.014e-02   1.472 0.141151
## shift9am    -2.049e-01  6.989e-02  -2.931 0.003375 **
## shift10am   -3.095e-01  7.082e-02  -4.370 1.24e-05 ***
```

```
## shift11am      -4.290e-01  7.902e-02  -5.429  5.66e-08 ***
## shift12pm      -7.961e-01  1.163e-01  -6.847  7.55e-12 ***
## shift1pm       -9.940e-01  1.775e-01  -5.601  2.13e-08 ***
## shift2pm        2.430e-01  8.489e-02   2.863  0.004201 **
## shifttother     9.492e-02  8.334e-02   1.139  0.254694
## Vig.ex.Time    -5.953e-04  1.598e-04  -3.724  0.000196 ***
## Walk.ex.Time    3.213e-04  7.425e-05   4.327  1.51e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
## Null deviance: 5090.1 on 351 degrees of freedom
## Residual deviance: 3674.1 on 336 degrees of freedom
## AIC: 4731.8
##
## Number of Fisher Scoring iterations: 6
```

```
par(mfrow=c(2,2))
plot(best.model)
```



```
SA1.model1 <- glm(pounds_gained ~ Total_Met_Min, data=metrics.dat, family = gaussian)
summary(SA1.model1)
```

```
##
## Call:
## glm(formula = pounds_gained ~ Total_Met_Min, family = gaussian,
## data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
```

```

## -13.398 -10.652 -3.498 4.922 59.348
##
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept)  1.065e+01  8.660e-01  12.300  <2e-16 ***
## Total_Met_Min 4.955e-04  6.562e-04   0.755    0.451
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for gaussian family taken to be 167.6111)
##
##      Null deviance: 58759  on 351  degrees of freedom
## Residual deviance: 58664  on 350  degrees of freedom
## AIC: 2805.7
##
## Number of Fisher Scoring iterations: 2
SA1.model2 <- glm(pounds_gained ~ gender + Age + height+ initial_BMI + shift + Total_Met_Min, data=metrics)
summary(SA1.model2)

##
## Call:
## glm(formula = pounds_gained ~ gender + Age + height + initial_BMI +
##      shift + Total_Met_Min, family = gaussian, data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -15.296   -7.001   -3.223    3.177   52.215
##
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept)  9.364e+00  3.330e+00   2.812  0.00521 **
## genderMale   -1.109e+00  1.396e+00  -0.795  0.42746
## genderMissing 7.381e-01  5.547e+00   0.133  0.89422
## Age         -6.461e-02  5.050e-02  -1.279  0.20163
## height      -1.099e-02  4.271e-02  -0.257  0.79709
## initial_BMI  4.628e-01  4.779e-02   9.684 < 2e-16 ***
## shift8am     1.145e+00  2.304e+00   0.497  0.61957
## shift9am    -1.248e+00  2.567e+00  -0.486  0.62722
## shift10am   -3.194e+00  2.618e+00  -1.220  0.22336
## shift11am   -4.066e+00  2.668e+00  -1.524  0.12846
## shift12pm   -6.743e+00  3.666e+00  -1.839  0.06675 .
## shift1pm    -7.696e+00  4.509e+00  -1.707  0.08875 .
## shift2pm     4.273e+00  3.626e+00   1.179  0.23935
## shifttother  1.401e+00  3.456e+00   0.405  0.68552
## Total_Met_Min -9.806e-05  5.906e-04  -0.166  0.86823
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for gaussian family taken to be 128.9481)
##
##      Null deviance: 58759  on 351  degrees of freedom
## Residual deviance: 43455  on 337  degrees of freedom
## AIC: 2726.1
##

```

```
## Number of Fisher Scoring iterations: 2
SA1.model3 <- glm(pounds_gained ~ gender + Age + height+ initial_BMI + shift + Vig.ex.Time + Mod.ex.time,
summary(SA1.model3)

##
## Call:
## glm(formula = pounds_gained ~ gender + Age + height + initial_BMI +
##      shift + Vig.ex.Time + Mod.ex.time + Walk.ex.Time, family = gaussian,
##      data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -16.745   -7.087   -3.171    3.088   51.973
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   9.117e+00  3.330e+00   2.738  0.00652 **
## genderMale    -9.337e-01  1.398e+00  -0.668  0.50474
## genderMissing  1.794e+00  5.605e+00   0.320  0.74912
## Age          -6.580e-02  5.066e-02  -1.299  0.19493
## height       -8.471e-03  4.283e-02  -0.198  0.84335
## initial_BMI   4.625e-01  4.760e-02   9.718 < 2e-16 ***
## shift8am      1.222e+00  2.312e+00   0.528  0.59751
## shift9am     -1.586e+00  2.572e+00  -0.617  0.53789
## shift10am    -3.062e+00  2.634e+00  -1.162  0.24590
## shift11am    -3.907e+00  2.682e+00  -1.457  0.14619
## shift12pm    -6.930e+00  3.669e+00  -1.889  0.05981 .
## shift1pm     -7.806e+00  4.511e+00  -1.731  0.08445 .
## shift2pm      4.525e+00  3.638e+00   1.244  0.21434
## shiftother    1.463e+00  3.459e+00   0.423  0.67269
## Vig.ex.Time  -6.023e-03  5.916e-03  -1.018  0.30937
## Mod.ex.time  -1.681e-05  4.751e-03  -0.004  0.99718
## Walk.ex.Time  3.512e-03  3.029e-03   1.159  0.24711
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for gaussian family taken to be 128.9494)
##
##      Null deviance: 58759  on 351  degrees of freedom
## Residual deviance: 43198  on 335  degrees of freedom
## AIC: 2728
##
## Number of Fisher Scoring iterations: 2
best.model <- stepAIC(SA1.model3, direction = "both")

## Start:  AIC=2728.02
## pounds_gained ~ gender + Age + height + initial_BMI + shift +
##      Vig.ex.Time + Mod.ex.time + Walk.ex.Time
##
##              Df Deviance    AIC
## - gender      2    43270 2724.6
## - Mod.ex.time  1    43198 2726.0
## - height      1    43203 2726.1
```

```

## - Vig.ex.Time 1 43332 2727.1
## - Walk.ex.Time 1 43371 2727.4
## - Age 1 43416 2727.8
## <none> 43198 2728.0
## - shift 8 45802 2732.6
## - initial_BMI 1 55376 2813.4
##
## Step: AIC=2724.61
## pounds_gained ~ Age + height + initial_BMI + shift + Vig.ex.Time +
## Mod.ex.time + Walk.ex.Time
##
## Df Deviance AIC
## - Mod.ex.time 1 43270 2722.6
## - height 1 43289 2722.8
## - Vig.ex.Time 1 43412 2723.8
## - Walk.ex.Time 1 43449 2724.1
## - Age 1 43491 2724.4
## <none> 43270 2724.6
## + gender 2 43198 2728.0
## - shift 8 45884 2729.2
## - initial_BMI 1 55406 2809.6
##
## Step: AIC=2722.61
## pounds_gained ~ Age + height + initial_BMI + shift + Vig.ex.Time +
## Walk.ex.Time
##
## Df Deviance AIC
## - height 1 43289 2720.8
## - Vig.ex.Time 1 43427 2721.9
## - Walk.ex.Time 1 43453 2722.1
## - Age 1 43491 2722.4
## <none> 43270 2722.6
## + Mod.ex.time 1 43270 2724.6
## + gender 2 43198 2726.0
## - shift 8 45891 2727.3
## - initial_BMI 1 55408 2807.7
##
## Step: AIC=2720.76
## pounds_gained ~ Age + initial_BMI + shift + Vig.ex.Time + Walk.ex.Time
##
## Df Deviance AIC
## - Vig.ex.Time 1 43451 2720.1
## - Walk.ex.Time 1 43473 2720.3
## <none> 43289 2720.8
## - Age 1 43597 2721.3
## + height 1 43270 2722.6
## + Mod.ex.time 1 43289 2722.8
## + gender 2 43203 2724.1
## - shift 8 45914 2725.5
## - initial_BMI 1 55686 2807.4
##
## Step: AIC=2720.08
## pounds_gained ~ Age + initial_BMI + shift + Walk.ex.Time
##

```

```

##           Df Deviance    AIC
## - Walk.ex.Time 1    43570 2719.1
## <none>           43451 2720.1
## - Age           1    43732 2720.3
## + Vig.ex.Time  1    43289 2720.8
## + height        1    43427 2721.9
## + Mod.ex.time   1    43434 2721.9
## + gender        2    43353 2723.3
## - shift         8    46008 2724.2
## - initial_BMI   1    55858 2806.5
##
## Step: AIC=2719.05
## pounds_gained ~ Age + initial_BMI + shift
##
##           Df Deviance    AIC
## <none>           43570 2719.1
## - Age           1    43864 2719.4
## + Walk.ex.Time  1    43451 2720.1
## + Vig.ex.Time   1    43473 2720.3
## + height        1    43546 2720.8
## + Mod.ex.time   1    43568 2721.0
## + gender        2    43467 2722.2
## - shift         8    46125 2723.1
## - initial_BMI   1    55930 2804.9

summary(best.model)


##
## Call:
## glm(formula = pounds_gained ~ Age + initial_BMI + shift, family = gaussian,
##      data = metrics.dat)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -15.616   -7.044   -3.328    3.251   52.616
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  8.40582    2.51758   3.339 0.000934 ***
## Age         -0.07103    0.04686  -1.516 0.130484
## initial_BMI  0.45721    0.04649   9.835 < 2e-16 ***
## shift8am     1.26638    2.28850   0.553 0.580376
## shift9am    -0.96581    2.53060  -0.382 0.702957
## shift10am   -2.94019    2.59101  -1.135 0.257270
## shift11am   -4.01664    2.65485  -1.513 0.131219
## shift12pm   -6.70662    3.64844  -1.838 0.066900 .
## shift1pm    -7.55090    4.48336  -1.684 0.093057 .
## shift2pm     4.18539    3.60450   1.161 0.246391
## shifttother  1.93054    3.30282   0.585 0.559261
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for gaussian family taken to be 127.7726)
##
##      Null deviance: 58759  on 351  degrees of freedom

```




```
## Residual deviance: 43570  on 341  degrees of freedom
## AIC: 2719
##
## Number of Fisher Scoring iterations: 2
```

```
par(mfrow=c(2,2))
plot(best.model)
```



```
par(mfrow=c(1,1))
hist(metrics.dat$pounds_gained)
```

Histogram of metrics.dat\$pounds_gained



Zero-Inflated Poisson

```
library(pscl)
```

```
## Classes and Methods for R developed in the  
## Political Science Computational Laboratory  
## Department of Political Science  
## Stanford University  
## Simon Jackman  
## hurdle and zeroinfl functions by Achim Zeileis
```

```
summary(m1 <- zeroinfl(LBS ~ Total_Met_Min, data = metrics.dat))
```

```
##  
## Call:  
## zeroinfl(formula = LBS ~ Total_Met_Min, data = metrics.dat)  
##  
## Pearson residuals:  
##      Min      1Q  Median      3Q      Max  
## -1.6521 -1.2215 -0.4045  0.5716  6.8119  
##  
## Count model coefficients (poisson with log link):  
##              Estimate Std. Error z value Pr(>|z|)  
## (Intercept)  2.820e+00  2.005e-02 140.617  <2e-16 ***  
## Total_Met_Min 4.099e-06  7.573e-06   0.541    0.588  
##  
## Zero-inflation model coefficients (binomial with logit link):  
##              Estimate Std. Error z value Pr(>|z|)
```

```

## (Intercept)   -0.5512065   0.1443698  -3.818 0.000135 ***
## Total_Met_Min -0.0001232   0.0001327  -0.928 0.353371
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Number of iterations in BFGS optimization: 1
## Log-likelihood: -1646 on 4 Df
summary(m2 <- zeroinfl(LBS ~ gender + Age + height+ initial_BMI + shift + Total_Met_Min, data = metrics

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
##
## Call:
## zeroinfl(formula = LBS ~ gender + Age + height + initial_BMI + shift +
##   Total_Met_Min, data = metrics.dat)
##
## Pearson residuals:
##      Min       1Q   Median       3Q      Max
## -4.6135 -0.9533 -0.4833  0.4673 11.1797
##
## Count model coefficients (poisson with log link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   2.818e+00  1.028e-01  27.417 < 2e-16 ***
## genderMale     5.622e-02  3.798e-02   1.480  0.13877
## genderMissing  5.336e-01  1.999e-01   2.669  0.00760 **
## Age            -3.180e-03  1.412e-03  -2.252  0.02430 *
## height         2.848e-03  1.348e-03   2.113  0.03461 *
## initial_BMI    5.156e-03  1.504e-03   3.427  0.00061 ***
## shift8am       -3.424e-02  6.183e-02  -0.554  0.57973
## shift9am       -1.796e-01  7.095e-02  -2.532  0.01135 *
## shift10am      -3.562e-01  7.123e-02  -5.000  5.73e-07 ***
## shift11am      -5.370e-01  7.940e-02  -6.763  1.35e-11 ***
## shift12pm      -6.707e-01  1.176e-01  -5.703  1.18e-08 ***
## shift1pm       -1.064e+00  1.779e-01  -5.982  2.21e-09 ***
## shift2pm        6.419e-02  8.494e-02   0.756  0.44980
## shifttother    -7.220e-02  8.571e-02  -0.842  0.39959
## Total_Met_Min  -2.115e-05  1.143e-05  -1.850  0.06424 .
##
## Zero-inflation model coefficients (binomial with logit link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  -1.082e-01  7.766e-01  -0.139   0.8892
## genderMale    1.132e+00  4.513e-01   2.508   0.0121 *
## genderMissing 1.542e+00  1.348e+00   1.144   0.2525
## Age           1.110e-02  1.280e-02   0.867   0.3857
## height        1.186e-02  8.548e-03   1.387   0.1654
## initial_BMI   -1.048e+00  8.819e+01  -0.012   0.9905
## shift8am      -7.619e-01  6.444e-01  -1.182   0.2371
## shift9am      -5.511e-01  7.007e-01  -0.786   0.4316
## shift10am     -5.332e-01  7.465e-01  -0.714   0.4750
## shift11am     -2.023e-01  7.442e-01  -0.272   0.7858
## shift12pm      7.471e-02  1.272e+00   0.059   0.9532
## shift1pm       5.738e-02  1.297e+00   0.044   0.9647
## shift2pm      -1.859e+00  9.972e-01  -1.865   0.0622 .
## shifttother   -7.946e-01  1.036e+00  -0.767   0.4429

```

```

## Total_Met_Min 9.749e-05 2.014e-04 0.484 0.6284
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Number of iterations in BFGS optimization: 1
## Log-likelihood: -1416 on 30 Df

summary(m3 <- zeroinfl(LBS ~ gender + Age + height+ initial_BMI + shift + Vig.ex.Time + Mod.ex.time + W

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
##
## Call:
## zeroinfl(formula = LBS ~ gender + Age + height + initial_BMI + shift +
## Vig.ex.Time + Mod.ex.time + Walk.ex.Time, data = metrics.dat)
##
## Pearson residuals:
##      Min      1Q  Median      3Q      Max
## -4.4256 -0.9573 -0.4910  0.3451 11.1840
##
## Count model coefficients (poisson with log link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  2.767e+00  1.002e-01  27.609 < 2e-16 ***
## genderMale   6.827e-02  3.788e-02   1.802 0.071497 .
## genderMissing 6.030e-01  1.616e-01   3.731 0.000190 ***
## Age         -3.062e-03  1.397e-03  -2.192 0.028355 *
## height       3.137e-03  1.355e-03   2.315 0.020587 *
## initial_BMI  5.156e-03  1.493e-03   3.454 0.000553 ***
## shift8am     -2.914e-02  6.072e-02  -0.480 0.631235
## shift9am     -2.125e-01  7.086e-02  -2.999 0.002707 **
## shift10am    -3.510e-01  7.105e-02  -4.940 7.79e-07 ***
## shift11am    -5.257e-01  7.928e-02  -6.631 3.34e-11 ***
## shift12pm    -7.020e-01  1.169e-01  -6.005 1.92e-09 ***
## shift1pm     -1.051e+00  1.777e-01  -5.919 3.25e-09 ***
## shift2pm     6.546e-02  8.553e-02   0.765 0.444040
## shifttother  -7.768e-02  8.574e-02  -0.906 0.364918
## Vig.ex.Time  -3.656e-04  1.702e-04  -2.148 0.031692 *
## Mod.ex.time   3.819e-05  9.368e-05   0.408 0.683542
## Walk.ex.Time  2.608e-04  6.702e-05   3.891 9.97e-05 ***
##
## Zero-inflation model coefficients (binomial with logit link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept) -0.2128543  0.7876161  -0.270  0.7870
## genderMale   1.1402087  0.4565210   2.498  0.0125 *
## genderMissing 1.4103404  1.3642942   1.034  0.3013
## Age         0.0130986  0.0130264   1.006  0.3146
## height      0.0106396  0.0086723   1.227  0.2199
## initial_BMI -1.0476335 87.0640084  -0.012  0.9904
## shift8am     -0.7409939  0.6472858  -1.145  0.2523
## shift9am     -0.4694167  0.7007571  -0.670  0.5029
## shift10am    -0.5717069  0.7605367  -0.752  0.4522
## shift11am    -0.1113415  0.7555825  -0.147  0.8828
## shift12pm     0.1928469  1.3016796   0.148  0.8822
## shift1pm     0.2235354  1.3032794   0.172  0.8638
## shift2pm    -1.8717091  1.0290655  -1.819  0.0689 .

```

```

## shiftother      -0.7791681  1.0369783  -0.751   0.4524
## Vig.ex.Time     0.0022485  0.0018770   1.198   0.2309
## Mod.ex.time     0.0011189  0.0019658   0.569   0.5692
## Walk.ex.Time    -0.0006442  0.0008913  -0.723   0.4698
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Number of iterations in BFGS optimization: 1
## Log-likelihood: -1409 on 34 Df

```