

Abstract

IceCube has measured high energy astrophysical neutrinos for the first time providing a powerful new probe of the universe, but many questions still remain. I will explore one strange quirk in the data. Despite generally large astrophysical uncertainties, I will show that this tension cannot be resolved with standard physics. The simplest consistent explanation is that some neutrinos are decaying. Finally, I will wrap up with predictions and a path forward.

Finding the Unexpected in IceCube

Peter B. Denton

N-Talk

September 21, 2018

PRL 121, 121802 (2018)

with I. Tamborra



The Niels Bohr
International Academy

VILLUM FONDEN



Finding the Unexpected in IceCube

Peter B. Denton

Neutrino-Talk

September 21, 2018

PRL 121, 121802 (2018)

with I. Tamborra



The Niels Bohr
International Academy

VILLUM FONDEN



New Probe of the Universe

IceCube has measured







- ▶ the first high energy (10 TeV - few PeV) astrophysical neutrinos
- ▶ their spectrum
- ▶ their flavor at the Earth
- ▶ oscillation parameters

Still working on

- ▶ determining their sources*
 - ▶ the data is quite isotropic
- ▶ particle physics
- ▶ ...



*the recent blazar event contributes $\sim 1\%$ to the total flux

IceCube Measures Tracks and Cascades




ν_μ CC

ν_e, ν_τ CC (and all NC)



Tension

$$\Phi(E) = \Phi_0 E^{-\gamma}$$



$$\Delta\gamma = 0.54$$

“The p-value for obtaining the combined fit result and the result reported here from an unbroken powerlaw flux is 3.3σ , and is therefore in significant **tension**.”

IC [1607.08006](#)

“This [cascade] fit [is] in **tension** with previous results based on through-going muons”

IC [1808.07629](#)

Conventional Wisdom

- ▶ High energy neutrinos are produced from full π decay
- ▶ Flavor ratio at source of 1:2:0 converts to 1:1:1* at Earth
- ▶ All neutrinos have the same energy[†]

*the fact that this ratio is 1:1:1 is coincidental not fundamental

[†]also a coincidence; kinematic corrections are small

Conventional Wisdom

- ▶ High energy neutrinos are produced from full π decay
- ▶ Flavor ratio at source of 1:2:0 converts to 1:1:1* at Earth
- ▶ All neutrinos have the same energy[†]

Some of these *must* be incorrect.

*the fact that this ratio is 1:1:1 is coincidental not fundamental


[†]also a coincidence; kinematic corrections are small

Need a phenomenon that non-trivially depends on
energy and **flavor** at the same time

Muon Cooling

$$\pi \rightarrow \nu_\mu + \mu$$

$$\mu \rightarrow \nu_\mu + \nu_e + e$$




- ▶ E.g. synchrotron
- ▶ More ν_μ at high energy
- ▶ E_b determined by B field

Muon Cooling

$$\pi \rightarrow \nu_\mu + \mu$$

$$\mu \rightarrow \nu_\mu + \nu_e + e$$



- ▶ E.g. synchrotron
- ▶ More ν_μ at high energy
- ▶ E_b determined by B field
- ▶ This doesn't work at all!
- ▶ Oscillations kill this
 - ▶ $\mu - \tau$ symmetry
- ▶ $\max \Delta\gamma \simeq 0.2$

Other Options

Neutron decay: $n \rightarrow p + e + \bar{\nu}_e$

- ▶ Produces extra ν_e 's
- ▶ Produced with pions in $p\gamma$ interactions
- ▶ Also come from photodissociation of heavy ions

Other Options

Neutron decay: $n \rightarrow p + e + \bar{\nu}_e$

- ▶ Produces extra ν_e 's
- ▶ Produced with pions in $p\gamma$ interactions
- ▶ Also come from photodissociation of heavy ions

But

- ▶ Neutrino energies are ~ 2 -3 orders of magnitude less for $p\gamma$
- ▶ Neutrino flux from heavy ions is also suppressed

D. Biehl, et al. [1705.08909](#)

X. Rodrigues, et al. [1711.02091](#)

New Physics!

We need a stronger effect, so we look to new physics.

- ▶ NSI with ultra-light mediators ($m \ll 1$ eV)

A. Joshipura, S. Mohanty [hep-ph/0310210](#)

M. Bustamante, S. Agarwalla [1808.02042](#)

- ▶ Pseudo-dirac neutrinos

L. Wolfenstein [Nucl. Phys. B186, 147 \(1981\)](#)

S. Pakvasa, A. Joshipura, S. Mohanty [1209.5630](#)

- ▶ Electrophilic dark matter decay
- ▶ Neutrino decay

New Physics!

We need a stronger effect, so we look to new physics.

- ▶ NSI with ultra-light mediators ($m \ll 1$ eV)

A. Joshipura, S. Mohanty [hep-ph/0310210](#)

M. Bustamante, S. Agarwalla [1808.02042](#)


- ▶ Pseudo-dirac neutrinos

L. Wolfenstein [Nucl. Phys. B186, 147 \(1981\)](#)

S. Pakvasa, A. Joshipura, S. Mohanty [1209.5630](#)

- ▶ Electrophilic dark matter decay

- ▶ Neutrino decay



New Physics!

We need a stronger effect, so we look to new physics.

- ▶ NSI with ultra-light mediators ($m \ll 1$ eV) weak

A. Joshipura, S. Mohanty [hep-ph/0310210](#)

M. Bustamante, S. Agarwalla [1808.02042](#)


- ▶ Pseudo-dirac neutrinos weak

L. Wolfenstein [Nucl. Phys. B186, 147 \(1981\)](#)

S. Pakvasa, A. Joshipura, S. Mohanty [1209.5630](#)

- ▶ Electrophilic dark matter decay strong but CMB

- ▶ Neutrino decay
strong, 3.4σ



Some Neutrinos Decay



Model recipe:

1. ν -decay depletes ν 's at low energy
2. Want fewer ν_μ at low energy
3. Let ν_2 and ν_3 decay
4. Keep ν_1 stable



*NO preferred at $\sim 3\sigma$

Some Neutrinos Decay



Mr. Stark,
I don't feel so good...

Model recipe:

1. ν -decay depletes ν 's at low energy
2. Want fewer ν_μ at low energy
3. Let ν_2 and ν_3 decay
4. Keep ν_1 stable



*NO preferred at $\sim 3\sigma$

Decay Process

- ▶ Neutrinos couple to a light/massless scalar ϕ : **Majoran**
- ▶ Secondaries?

Decay Process

- ▶ Neutrinos couple to a light/massless scalar ϕ : **Majoran**
- ▶ Secondaries?

Visible

- ▶ Lighter states regenerated
- ▶ Energetics depend strongly on absolute mass scale

Decay Process

- ▶ Neutrinos couple to a light/massless scalar ϕ : **Majoran**
- ▶ Secondaries?

Visible

- ▶ Lighter states regenerated
- ▶ Energetics depend strongly on absolute mass scale

Invisible

- ▶ Right handed neutrinos
- ▶ ν_L with much lower energies
- ▶ Unparticles, ...

Decay Process

- ▶ Neutrinos couple to a light/massless scalar ϕ : **Majoran**
- ▶ Secondaries?

Visible

- ▶ Lighter states regenerated
- ▶ Energetics depend strongly on absolute mass scale


Invisible

- ▶ Right handed neutrinos
- ▶ ν_L with much lower energies
- ▶ Unparticles, ...

To get *our* model:


- ▶ ν_1 decay is kinematically inaccessible
- ▶ Coupling to ν_1 is much smaller
- ▶ Lifetime estimated by typical $E \simeq 100$ TeV and $z \simeq 1$:
 $\tau_2/m_2 \simeq \tau_3/m_3 \sim 10^2$ s/eV

Track to Cascade Ratio (At Earth)



*the deviation from 1/2 as expected is due to
SM corrections that are accounted for

Invisible ν Decay Constraints and Evidence



PBD, I. Tamborra 1805.05950

S. Hannestad, G. Raffelt [hep-ph/0509278](#)


KamiokaNDE-II PRL 58 1490 (1987)

G. Pagliaroli, et al. 1506.02624

J. Berryman, A. de Gouvea, D. Hernandez 1411.0308

M. Gonzalez-Garcia and M. Maltoni 0802.3699

Invisible ν Decay Constraints and Evidence



PBD, I. Tamborra 1805.05950

S. Hannestad, G. Raffelt [hep-ph/0509278](#)

KamiokaNDE-II PRL 58 1490 (1987)


G. Pagliaroli, et al. 1506.02624

J. Berryman, A. de Gouvea, D. Hernandez 1411.0308

M. Gonzalez-Garcia and M. Maltoni 0802.3699

ν_2, ν_3 decay leads to 16%
reduction in $\bar{\nu}_e$ flux:
SN1987A doesn't apply

Invisible ν Decay Constraints and Evidence



PBD, I. Tamborra [1805.05950](#)

S. Hannestad, G. Raffelt [hep-ph/0509278](#)

KamiokaNDE-II PRL 58 1490 (1987)

G. Pagliaroli, et al. [1506.02624](#)

J. Berryman, A. de Gouvea, D. Hernandez [1411.0308](#)

M. Gonzalez-Garcia and M. Maltoni [0802.3699](#)


N. Bell, E. Pierpaoli, K. Sigurdson [astro-ph/0511410](#)

ν_2, ν_3 decay leads to 16%
reduction in $\bar{\nu}_e$ flux:

SN1987A doesn't apply

CMB constraints assume
all flavors decay,
< 3 decaying is allowed...

Invisible ν Decay Constraints and Evidence



PBD, I. Tamborra [1805.05950](#)

S. Hannestad, G. Raffelt [hep-ph/0509278](#)

KamiokaNDE-II PRL 58 1490 (1987)

G. Pagliaroli, et al. [1506.02624](#)

J. Berryman, A. de Gouvea, D. Hernandez [1411.0308](#)

M. Gonzalez-Garcia and M. Maltoni [0802.3699](#)

N. Bell, E. Pierpaoli, K. Sigurdson [astro-ph/0511410](#)

M. Archidiacono, et al. [1404.5915](#)

ν_2, ν_3 decay leads to 16%
reduction in $\bar{\nu}_e$ flux:

SN1987A doesn't apply

CMB constraints assume

all flavors decay,

< 3 decaying is allowed...


and may be slightly preferred

Deficit(?) of ν_τ Events

- ▶ IceCube can *sometimes* identify ν_τ CC
- ▶ Should have seen 2-3 events, seen none*

IC 1710.01191

ν_τ suppression from decay:



Multiply with efficiency to find
total sensitivity reduced by 59%

* ~ 1 new net event may exist,
new sensitivities will be higher

The Message

- ▶ There seems to be some tension in IceCube's data
- ▶ Inconsistent with standard physics
 - ▶ Multiple sources don't help
 - ▶ Multi-zone type conspiracies could solve this
- ▶ DM is an option, not great
- ▶ Neutrino decay works, favored at 3.4σ

The Message

- ▶ There seems to be some tension in IceCube's data
- ▶ Inconsistent with standard physics
 - ▶ Multiple sources don't help
 - ▶ Multi-zone type conspiracies could solve this
- ▶ DM is an option, not great
- ▶ Neutrino decay works, favored at 3.4σ

It is possible to make strong particle physics statements in
astrophysical environments

The Message

- ▶ There seems to be some tension in IceCube's data
- ▶ Inconsistent with standard physics
 - ▶ Multiple sources don't help
 - ▶ Multi-zone type conspiracies could solve this
- ▶ DM is an option, not great
- ▶ Neutrino decay works, favored at 3.4σ

It is possible to make strong particle physics statements in
astrophysical environments

Looking forward:

- ▶ Play close attention to ν_τ searches
- ▶ Anisotropy + flavor (DM)
- ▶ More flavor + energy dependent fits: BPL

Thank you!

Backups

Neutrino Decay Affects Flavor

The oscillation averaged probability is



$$\bar{P}(\nu_\alpha \rightarrow \nu_\beta) = \sum_{i=1}^3 |U_{\alpha i}|^2 |U_{\beta i}|^2 e^{-\Lambda_i}$$

$$\Lambda_i \equiv \frac{d_H f(z) m_i}{E_\nu \tau_i}$$


$$f(z) = \int_0^z \frac{dz'}{(1+z')^2 \sqrt{(1+z')^3 \Omega_m + \Omega_\Lambda}}$$

We take $\Lambda_2 = \Lambda_3$ for simplicity and $\Lambda_1 = 0$.

IceCube's Tracks and Cascades



IceCube Neutrinos Origin



$< 9.5\%$ galactic fraction at 90% CL

PBD, D. Marfatia, T. Weiler [1703.09721](#)