Document Identifier: **PMC8006653**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: All animal experiments were approved by the local animal ethics committee.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Methods Animals and experimental protocol Male C57Bl6 wild-type mice (n = 42) (Janvier laboratory, Le Genest-Saint-Isle, France) aged 8 to 10 weeks were randomly divided into five groups (Figure 1).
Subject Demographics
Age: Methods Animals and experimental protocol Male C57Bl6 wild-type mice (n = 42) (Janvier laboratory, Le Genest-Saint-Isle, France) aged 8 to 10 weeks were randomly divided into five groups (Figure 1).
Randomization
Methods Animals and experimental protocol Male C57Bl6 wild-type mice (n = 42) (Janvier laboratory, Le Genest-Saint-Isle, France) aged 8 to 10 weeks were randomly divided into five groups (Figure 1).
Blinding
Data were collected and blindly analysed.
Power Analysis
The sample size was calculated using The Cohens d calculation.
Replication
not required.
Data Information
Availability: The final solution 185 mL of 1 mM DHE in 50 mL of phosphate-buffered saline (final concentration: 37 mM) was deposited (1 mL) on each slide in a dark room.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
Methods Animals and experimental protocol Male C57Bl6 wild-type mice (n = 42) (Janvier laboratory, Le Genest-Saint-Isle, France) aged 8 to 10 weeks were randomly divided into five groups (Figure 1).	C57Bl6		
Software and Algorithms			
Images were assembled using Photoshop (Adobe Systems, San Jose, CA).	Photoshop		Suggestion: Adobe Photoshop (RRID:SCR_014199)(link)
21 Statistical analysis was performed using GraphPad Prism 7.	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Interaction between chemotherapies and treatment was performed by a two-way ANOVA analysis followed by a Bonferroni post hoc test.	ANOVA
	Bonferroni post hoc test
Shapiro-Wilk test was used to assess normality.	Shapiro-Wilk test
The Mann-Whitney U test was used to assess statistical differences between two experimental conditions.	Mann-Whitney U test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq