Document Identifier: **PMC8010833**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study was performed in strict accordance with the provisions of the Declaration of Helsinki and its amendments, and was approved by the ethics committee of Qilu Hospital of Shandong University (approval no. KYLL-2019-270).
Consent: All the participants provided their written informed consent.
Inclusion and Exclusion Criteria
The exclusion criteria were severe liver disease (liver cirrhosis or serum aspartate aminotransferase (AST) or alanine aminotransferase (ALT) activities >120 U/L), severe kidney disease (estimated glomerular filtration rate [eGFR] <60 mL/minute/1.73 m2), malignant disease, haematological disease and serum bilirubin concentration greater than two times the upper limit of the normal range.
Attrition
not detected.
Sex as a biological variable
Sarcopenia was defined as a SMI <7.0 kg/m2 in men and <5.4 kg/m2 in women, according to the Asian Working Group for Sarcopenia recommendations.
Subject Demographics
Age: Materials and methods Study participants Sarcopenia is more common in middle-aged and elderly patients, and most of the inpatients with T2DM in our hospital were >50 years old.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
All the statistical analyses were performed using SPSS 23.0 software (IBM Corp., Armonk, NY, USA) and $P < 0.05$ was considered to represent statistical significance.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
19 Statistical analysis The Kolmogorov–Smirnov test was used to determine whether datasets were normally distributed.	Kolmogorov–Smirnov test
Student’s t-test (normally distributed data) or Mann–Whitney U-test (non-normally distributed data) were used to compare differences between the two groups.	Student’s t-test
	Mann–Whitney U-test
The relationships between serum bilirubin concentration and SMI were assessed using Pearson’s correlation analysis and multiple linear regression analysis in both men and women.	Pearson’s correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq