Document Identifier: **PMC8018676**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: All sites had institutional review board (IRB) approval, both locally and at the University of Pittsburgh or University of Iowa, with written informed consent for genomic studies and data sharing.
Consent: All sites had institutional review board (IRB) approval, both locally and at the University of Pittsburgh or University of Iowa, with written informed consent for genomic studies and data sharing.
Inclusion and Exclusion Criteria
Subjects where the specific subtype of cleft was not known were excluded from this study.
Attrition
For the current study, affected individuals were classified as either having a bilateral cleft lip (BCL N = 44), a bilateral cleft lip and palate (BCLP N = 530), a unilateral cleft lip (UCL N = 434), or a unilateral cleft lip and palate (UCLP N = 1,123).
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

Replication cohort To replicate the statistically significant results from our modifier analysis , data from the GENEVA consortium were used , which was described previously . 2 , 4 , 21 Briefly , this cohort recruited affected individual-parent trios , where the affected individual had an oral cleft .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
33 Epigenomic context of results Topologically associated domains (TADs) were defined for significantly associated loci using the H1-ESC cell line in 3D Genome Browser .	H1-ESC		
Software and Algorithms			
Samples were genotyped for approximately 580,000 single-nucleotide polymorphic (SNP) markers from the Illumina HumanCore+Exome array, of which approximately 539,000 SNPs passed quality control filters recommended by the Center for Inherited Disease Research (CIDR) and the Genetics Coordinating Center (GCC) at the University of Washington. 2 These data were then phased with SHAPEIT2 23 and imputed with IMPUTE2 24 to the 1000 Genomes Project phase 3 release (September 2014) reference panel.	1000 Genomes Project		Suggestion: 1000 Genomes Project and AWS (RRID:SCR_008801)(link)
28 , 29 Regional association plots were made with LocusZoom , where the LD blocks and recombination rates were estimated from European populations .	LocusZoom		Suggestion: LOCUSZOOM (RRID:SCR_009257)(link)
The samples were genotyped for approximately 589,000 SNPs using the Illumina Human610-Quadv.1_B BeadChip , phased using SHAPEIT , and imputed to the 1000 Genomes Project phase I (June 2011) reference panel using IMPUTE2 .	1000 Genomes		Suggestion: 1000 Genomes Project and AWS (RRID:SCR_008801)(link)
	IMPUTE2		Suggestion: IMPUTE2 (RRID:SCR_013055)(link)
Imputed genotype probabilities were converted to most-likely genotype calls with GTOOL .	GTOOL		Suggestion: GTOOL (RRID:SCR_009215)(link)
Because the replication cohort did not include related individuals , the modifier analyses (comparing BCL versus UCL and BCLP versus UCLP) were conducted using logistic regression models in PLINK (v1.9) , with sex and the first four PCs as quantitative covariates , instead of the mixed-model approach that adjusts for relatedness implemented in GENESIS.	PLINK		Suggestion: PLINK (RRID:SCR_001757)(link)
	GENESIS		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
A chi-square test was used to determine if the number of SNPs that both had similar CIs and were previously reported in the literature overlapped more than expected by chance.	chi-square test
p values less than a Bonferroni correction for the number of SNPs in the region (0.05the number of SNPs tested) were considered to be evidence of significant replication .	Bonferroni

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq