Document Identifier: **PMC8019962**

# SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

**Table 1: Rigor Adherence Table**

| |
|---|
| Ethics |
| IRB: The animal protocols followed the guidelines of the International Association of the Study of Pain (in accordance with the UK Animal Act) and had approval from the Ethics Committee of Shaoxing Peoples Hospital (Shaoxing Hospital, Zhejiang University School of Medicine) (approval number: 2020-79). |
| Euthanasia Agents: Central cannulation and microinjection Rats underwent anesthesia by sodium pentobarbital injection (45 mgkg, i.p.) and fixation in a stereotaxic apparatus for central cannulation and microinjection 20. |
| Inclusion and Exclusion Criteria |
| not detected. |
| Attrition |
| not detected. |
| Sex as a biological variable |
| Animals Specific-pathogen-free male Sprague-Dawley rats (200-250 g) were provided by Shanghai SLAC Laboratory Animal Center, Co. Ltd. (Shanghai, China). |
| Subject Demographics |
| Weight: Animals Specific-pathogen-free male Sprague-Dawley rats (200-250 g) were provided by Shanghai SLAC Laboratory Animal Center, Co. Ltd. (Shanghai, China). |
| Randomization |
| not detected. |
| Blinding |
| not detected. |
| Power Analysis |
| not detected. |
| Replication |
| not required. |

**Table 2: Key Resources Table**

| Your Sentences | REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|---|
| Antibodies | | | |
| After blocking with 5 skim milk in tris buffered saline with Tween 20 for 1 hour at ambient, the membranes underwent successive incubations with rabbit primary antibodies against IL-1R, IL-6R, and TNFR1 (1:500 Neuromics, Edina, MN or Abcam), and horseradish peroxidase-conjugated anti-rabbit immunoglobulin G (1:250). | IL-1R | | |
| | IL-6R | | |
| | TNFR1 | | |
| | anti-rabbit immunoglobulin G | | |
| In a similar fashion, p-PI3K, p-Akt, and p-mTOR amounts as well as the levels of their non-phosphorylated forms were determined by immunoblot, with primary antibodies from Neuromics or Abcam. 7. | p-PI3K, | | |
| | p-PI3K | | |
| | p-Akt | | |
| Experimental Models: Organisms/Strains | | | |
| Animals Specific-pathogen-free male Sprague-Dawley rats (200-250 g) were provided by Shanghai SLAC Laboratory Animal Center, Co. Ltd. (Shanghai, China). | Sprague-Dawley | | Suggestion: RRID:RGD_70508( link) |
| Software and Algorithms | | | |
| Autoradiography was employed for detection and the NIH Scion image software (NIH, Bethesda, MA) was utilized for quantifying immunoreactive bands by densitometry. | Scion image | | |
| Statistical analyses GraphPad Prism 8 (GraphPad Software, San Diego, CA) was employed for data analysis. | GraphPad | | Suggestion: GraphPad Prism, RRID:SCR_002798( link) |

**Other Entities Detected**

| Your Sentences | Recognized Entity |
|---|---|
| Statistical Tests | |
| One- or two-way analysis of variance and post-hoc Bonferroni, Tukeys, or Dunnetts tests were performed for comparisons. | analysis of variance |
| | Bonferroni |
| | Dunnetts tests |

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq