Document Identifier: **PMC8030014**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: 19 Informed consent was obtained from all patients.
Inclusion and Exclusion Criteria
Exclusion criteria were as follows: Known significant arterial abnormalities in the upper extremities, known cardiac arrhythmia, upper arm circumference below 15 cm or above 42 cm, or possible extreme uncontrolled hypertension (ie, resting office BP > 250140 mmHg).
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
This reanalysis was performed by a SOMNOmedics programmer blinded to the corresponding OscBP measurements.
Power Analysis
The study was designed to have at least 80 power to detect a 5-mmHg systolic BP difference between the devices.
Replication
A small questionnaire on device-related pain andor discomfort as well as sleeping habits was given to all participants. 2.3 Estimation of markers of arterial stiffness To study the potential impact of aortic stiffness

as well as peripheral wave reflection on the measurement of the PTT device, carotid-femoral pulse wave velocity (cfPWV) as well as radially obtained pulse wave analysis (PWA) was performed in duplicate with the SphygmoCor system (version 8.2, Atcor Medical, Sydney, Australia).

Number: A small questionnaire on device-related pain andor discomfort as well as sleeping habits was given to all participants. 2.3 Estimation of markers of arterial stiffness To study the potential impact of aortic stiffness as well as peripheral wave reflection on the measurement of the PTT device, carotid-femoral pulse wave velocity (cfPWV) as well as radially obtained pulse wave analysis (PWA) was performed in duplicate with the SphygmoCor system (version 8.2, Atcor Medical, Sydney, Australia).

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
With the patient sitting in an upright position with back support, a conventional oscillometric BP device (OscBP, Spacelabs 90 217, Spacelabs Healthcare, Hawthorne, USA) 21 was fitted on the right arm, while the SomBP was fitted on the left arm.	Spacelabs Healthcare		
25 All statistical analyses were performed using STATA software (v. 16.0, Statacorp, Texas, USA).	STATA		Suggestion: Stata (RRID:SCR_012763)(link)
	Statacorp		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Means were compared using paired t test.	t test
Pearsons correlation was performed to test for association between different BP measurement modalities.	Pearsons correlation
Categorical variables were compared using chi-squared tests.	chi-squared tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq