Document Identifier: **PMC8031359**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The protocol conformed to the Declaration of Helsinki and regional regulatory guidelines, and the study was approved by regional institutional review boards (IRBs).
Consent: Each patient signed an IRB-approved consent form before any study procedures were initiated.
Inclusion and Exclusion Criteria
Exclusion criteria included known or suspected secondary hypertension or severe diastolic hypertension (seated diastolic DBP >114 mm Hg) severe renal impairment (estimated glomerular filtration rate <30 mLmin1.73 m2) history of a major cardiovascular event in the previous 6 months type 1 or poorly controlled type 2 diabetes mellitus (glycated hemoglobin >8) poor compliance with study medication during the placebo run-in period or hyperkalemia (serum potassium concentration>upper limit of normal for the reference laboratory).
Attrition
not detected.
Sex as a biological variable
In addition, night-shift workers, pregnant or nursing women, and women of childbearing potential not using approved means of contraception were excluded from participation.
Subject Demographics
Age: At randomization, each patient was required to be 18 years or older, with a clinic SBP >=150 and <=180 mm Hg and a 24-hour mean SBP >=130 and <=170 mm Hg.
Randomization
2 Methods 2.1 Patients and study design This was a 6-week, randomized, double-blind, multicenter, placebo-controlled study comparing the antihypertensive effects and safety and tolerability of AZL-M, 40 or 80 mg once daily, with placebo in AA patients with stages 1 or 2 systolic hypertension.
Blinding
not detected.
Power Analysis

Given the assumptions of a standard deviation of 13 mm Hg and a 20 dropout rate, the sample size of this study provided at least 90 power to detect a difference of 6 mm Hg between each of the AZL-M treatment groups and placebo for the primary end point.
Replication
Clinic BP was measured in triplicate in the nondominant arm using an automated BP monitor (Omron HEM 705-CP, Lake Forest, IL, USA) after the patient was seated for 5 minutes.
Number: Clinic BP was measured in triplicate in the nondominant arm using an automated BP monitor (Omron HEM 705-CP, Lake Forest, IL, USA) after the patient was seated for 5 minutes.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq