Document Identifier: **PMC8039634**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This study was approved by the Research Ethics Committee of Royal Hospital.
Inclusion and Exclusion Criteria
Patient with incomplete information regarding disease status during the follow-up period were excluded from the study.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: All Omani patients aged 18 years and above who were diagnosed with DTC and followed up at the NDEC thyroid oncology clinic during this period were included in the study.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Gender, tumor characteristics, lymph node status, distant metastases, biochemical profile (e.g., thyroglobulin (Tg), thyroglobulin antibody (TgAb), and thyroid-stimulating hormone (TSH) levels), and imaging results were recorded at subsequent follow-up visits.	thyroglobulin (Tg), thyroglobulin		
Recombinant DNA			
Those with pT1 or pT2 tumors at N0M0 stage without aggressive histology were categorized as low risk, while those with aggressive histology or pT3, or N1 tumors were categorized as intermediate risk.	pT1		
	pT2		Suggestion: RRID:Addgene_26557(link)
	pT3		Suggestion: RRID:Addgene_69123(link)
Patients with pT4 tumors or tumors at any T or N stage with M1 progression were considered high risk.	pT4		Suggestion: RRID:Addgene_108352(link)
Software and Algorithms			
Data were analyzed using the (IBM SPSS Statistics for Windows, Version 20.0.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Associations were tested using univariate or multivariate tests.	multivariate tests
For the univariate analysis, chi-square or independent t-tests were performed according to the nature of the variables.	chi-square
	t-tests
For the multivariate analysis, binary logistic regression was used to adjust for potential confounders.	binary logistic regression
Kaplan-Meier curves were plotted, and log-rank tests conducted to compare prognostic factors for DFS.	log-rank tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq