Document Identifier: **PMC8045161**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Number: Statistical analysis All the data were obtained from three independent experiments and presented as the mean +- standard deviation , with between-group differences analyzed by multiple t-tests followed by the Holm-Sidak correction test .
Cell Line Authentication

not detected.	
Data Information	
Identifiers: The gene expression datasets (GSE4342 and GSE38302) were downloaded from the GEO database (20,21).	GSE4342
	GSE38302

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The membranes were then incubated with the following specific monoclonal or polyclonal primary antibodies overnight at 4degC: Anti-CNTN1 (1:200 cat. no. sc-136133 Santa Cruz Biotechnology , Inc .)	Anti-CNTN1		
, anti-E-cadherin (1:200 cat. no. sc-8426 Santa Cruz Biotechnology , Inc .)	anti-E-cadherin		
, anti-N-cadherin (1:200 cat. no. sc-7939 Santa Cruz Biotechnology , Inc .) , anti-vimentin (1:200 cat. no. sc-6260 Santa Cruz Biotechnology , Inc .) and anti-Actin (1:5,000 cat. no. A5441 Sigma-Aldrich Merck KGaA) .	anti-N-cadherin		
	anti-Actin		
Subsequently , the membranes were incubated with the corresponding horseradish peroxidase-conjugated anti-mouse IgG (1:10,000 cat. no. 7076 Cell Signaling Technology , Inc .) or anti-rabbit IgG antibodies (1:10,000 cat. no. 7074 Cell Signaling Technology , Inc .) for 1 h at room temperature.	anti-mouse IgG		
	anti-rabbit IgG		
An anti-a-tubulin (cat. no. ab24610 Abcam) antibody was added to the slides at a dilution of 1:200 and incubated overnight at 4degC .	anti-a-tubulin		
The samples were then treated with an Alexa Fluor-conjugated anti-mouse antibody (1:1,000 cat. no. R37115 Thermo Fisher Scientific , Inc .) for 1 h protected from light at room temperature.	anti-mouse		
Experimental Models: Cell Lines			
The differential expression of genes between gefitinib-sensitive cells (HCC827 and Calu3) and intrinsically resistant cells (A549, H460 and H1975) in GSE4342, and between acquired gefitinib resistant cells (PC9GR) and parental cells (PC9) in GSE38302 was determined using GEO2R (http:www.ncbi.nlm.nih.govgeogeo2r).	HCC827		Suggestion: RRID:CVCL_2063(link)
	A549		Suggestion: RRID:CVCL_0023(link)
	H460		Suggestion: RRID:CVCL_0459(link)
Virus packaging and transfection Briefly, the 293T cells were purchased from the American	293T		Suggestion: RRID:CVCL_0063(link)

Type Culture Collection and were cultured in high-glucose DMEM, supplemented with 10 FBS.			
Cell cycle and apoptosis assay For the cell cycle analysis , 2x105 A549-shLuc and A549-shCNTN1-1 cells were seeded in a 6-well plate.	A549-shLuc		
	A549-shCNTN1-1		
Recombinant DNA			
The cells were cultured to 50-60 confluency and co-transfected with either 10 ug control (shLuc) or shRNA-CNTN1, with a packaging plasmid 7.5 ug psPAX2 (Invitrogen Thermo Fisher Scientific Inc.) and an envelope plasmid 2.5 ug pMD2.G (Invitrogen Thermo Fisher Scientific Inc.) using 40 ug PEI reagent (Sigma-Aldrich Merck KGaA).	psPAX2		
	pMD2.G		
Software and Algorithms			
Materials and methods Public databases Gene Expression Omnibus (GEO http:www.ncbi.nlm.nih.govgds) was used for the analysis of CNTN1 mRNA expression levels between gefitinib-sensitive and -insensitive NSCLC.	Gene Expression Omnibus		Suggestion: Gene Expression Omnibus (GEO) (RRID:SCR_005012)(link)
The differential expression of genes between gefitinib-sensitive cells (HCC827 and Calu3) and intrinsically resistant cells (A549, H460 and H1975) in GSE4342, and between acquired gefitinib resistant cells (PC9GR) and parental cells (PC9) in GSE38302 was determined using GEO2R (http:www.ncbi.nlm.nih.govgeogeo2r).	GEO2R		Suggestion: GEO2R (RRID:SCR_016569)(link)
The Kaplan-Meier plot was used to determine overall survival (OS) time using Kaplan-Meier Plotter database (https:kmplot.com) (22) and a log rank test.	Kaplan-Meier Plotter		Suggestion: Kaplan Meier Plotter (RRID:SCR_018753)(link)
The number of cells that had migrated to the lower side of the membrane were counted using ImageJ software (version 1.52a NIH) .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
All statistical analyses were performed using GraphPad Prism v7.0 (	GraphPad Prism		
(GraphPad Software, Inc.).	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The Kaplan-Meier plot was used to determine overall survival (OS) time using Kaplan-Meier Plotter database (https:kmplot.com) (22) and a log rank test.	log rank test
Statistical analysis All the data were obtained from three independent experiments and presented as the mean +- standard deviation , with between-group differences analyzed by multiple t-tests followed by the Holm-Sidak correction test .	multiple t-tests
	Holm-Sidak correction test
Unresolved RRIDs	
The cells were cultured to 50-60 confluency and co-transfected with either 10 ug control (shLuc) or shRNA-CNTN1, with a packaging plasmid 7.5 ug psPAX2 (Invitrogen Thermo Fisher Scientific Inc.) and an envelope plasmid 2.5 ug pMD2.G (Invitrogen Thermo Fisher Scientific Inc.) using 40 ug PEI reagent (Sigma-Aldrich Merck KGaA).	plasmid 7.
	plasmid 2.

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq