Document Identifier: **PMC8048835**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2 PATIENTS AND METHODS 2.1 Ethical considerations This study is part of the registry on epidemiology and management of patients with UVFP, which has been approved by the Ethics Commission of the Medical University of Vienna (EK 1507/2019).
Consent: According to the Austrian law and the guidelines of the research ethics committee, written informed consent was obtained from all patients.
Inclusion and Exclusion Criteria
Patients with BVFP were excluded from this study.
Attrition
To the other 72 patients with persisting UVFP, either VT or SES was recommended. 26/72 patients opted for VT (Group 1) and 25/72 patients for SES (Group 2), whereas 21 patients did not remain in our medical care due to other medical reasons, distance to the hospital, or lack of therapy motivation.
Sex as a biological variable
Of the 51/72 UVFP patients remaining in therapy, 35 females and 16 males with mean age (+-SD) age of 54 (+-14) years. 33/51 patients had a left-sided and 18/51 patients a right-sided UVFP (Table 1).
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
2.5 Statistics Data were analysed using IBM SPSS 21 (Armonk).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The primary outcome parameter "vocal fold mobility" at time point "2-week post-operative" was compared to "3 months post-operative" within the two groups using the McNemar test.	McNemar test
The outcome parameters "R", "B", and "H," "vocal fold position" and "glottal closure" at time point "2-week post-operative" were compared to "3 months post-operative" within the two groups using the Wilcoxon-test.	Wilcoxon-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq