Document Identifier: **PMC8050764**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Methods Institutional review board approval was obtained before we proceeded with this study.
Consent: All patients read and signed the informed consent.
Inclusion and Exclusion Criteria
Patients were excluded from this study if MRI had been performed more than 4 weeks after injury or if they had previous trauma or surgery in the affected knee, associated ligament injuries, or poor-quality MRI scans.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Clinical notes, operative reports, and MRI scans of adult patients younger than 40 years of age who had sustained noncontact unilateral ACL rupture during soccer (soccer group) or basketball (basketball group) were reviewed to determine the detailed injury information.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
For patients with bone bruises, the bone bruise borders were traced using the freehand region-of-interest (ROI) function of ImageJ software (National Institutes of Health).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
All statistical analyses were performed using the SPSS statistical package (Version 16 SPSS).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical Analysis The bone bruise pattern and meniscal injury pattern between soccer and basketball players were compared using chi-square analysis.	chi-square
A 2 x 2 analysis of variance (ANOVA) with a mixed design was performed to compare the specific bone bruise locations in the lateral-medial direction in patients with the LF+LT pattern, between groups (soccer and basketball groups) and between bones (femur and tibia), for the tibia and the femur.	analysis of variance
	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq