Materials Design Analysis Reporting (MDAR) Checklist for Authors

The MDAR framework establishes a minimum set of requirements in transparent reporting applicable to studies in the life sciences (see Statement of Task: doi:10.31222/osf.oio/9sm4x.). The MDAR checklist is a tool for authors, editors and others seeking to adopt the MDAR framework for transparent reporting in manuscripts and other outputs. Please refer to the MDAR Elaboration Document for additional context for the MDAR framework.

Materials

Antibodies	Yes (indicate where provided: page no/section/legend)	n/a
For commercial reagents, provide supplier name, catalogue number and RRID, if available	Yes, 1 antibody detected, 0 RRID provided : AQP4 Please add identifiers for all resources where possible	

Cell Materials	Yes (indicate where provided: page no/section/legend)	n/a
Cell lines: Provide species information, strain. Provide accession number in repository OR supplier name, catalog number, clone number, OR RRID	No cell lines detected Please add identifiers for all resources where possible	
Primary cultures: Provide species, strain, sex of origin, genetic modification status.	Not currently checked by SciScore	

Experimental Animals	Yes (indicate where provided: page no/section/legend)	n/a
Laboratory animals: Provide species, strain, sex, age, genetic modification status. Provide accession number in repository OR supplier name, catalog number, clone number, OR RRID	No organisms detected Please add identifiers for all resources where possible	
Animal observed in or captured from the field: Provide species, sex and age where possible	Not currently checked by SciScore	
Model organisms: Provide Accession number in repository (where relevant) OR RRID	See laboratory animals section for information.	

Plants and microbes	Yes (indicate where provided: page no/section/legend)	n/a
Plants: provide species and strain, unique accession number if available, and source (including location for collected wild specimens)	Not currently checked by SciScore	
Microbes: provide species and strain, unique accession number if available, and source	Not currently checked by SciScore	

Human research participants	Yes (indicate where provided: page no/section/legend)	n/a
Identify authority granting ethics approval (IRB or equivalent committee(s), provide reference number for approval.	Standard Protocol Approvals, Registrations, and Patient Consents The Institutional Review Board of the NCC approved the study, and written informed consent was obtained from all participants.	
Provide statement confirming informed consent obtained from study participants.	Standard Protocol Approvals, Registrations, and Patient Consents The Institutional Review Board of the NCC approved the study, and written informed consent was obtained from all participants.	
Report on age and sex for all study participants.	**Age:**Pretreatment samples were collected within a median of 2 months (range 1-3 months) from prior attacks. **Sex:**not detected.	

Design

Study protocol	Yes (indicate where provided: page no/section/legend)	n/a
For clinical trials, provide the trial registration number OR cite DOI in manuscript.	Not detected.	

Laboratory protocol	Yes (indicate where provided: page no/section/legend)	n/a
Provide DOI or other citation details if detailed step-by-step protocols are available.	Not detected.	

Experimental study design (statistics details)	Yes (indicate where provided: page no/section/legend)	n/a
State whether and how the following have been done, or if they were not carried out		
Sample size determination	not detected.	
Randomization	4,5 Twenty participants newly treated with rituximab according to Kim protocol 6 and underwent at least a 24-month follow-up were randomly selected from the NMOSD cohort of the National Cancer Center (NCC) to minimize interindividual variability of treatment 14 participants without clinical attacks (stable) during follow-up were included for focused investigation of interattack periods, and 6 participants with clinical attacks (unstable) were included for attack-related measurements.	
Blinding	sGFAP and sNfL concentrations were measured using a single-molecule array assay (Simoa, Quanterix, MA). sGFAP and sNfL levels were independently evaluated by a blinded examiner in duplicate, and mean intra-assay and interassay coefficients of variation were below 10.	
inclusion/exclusion criteria	not detected.	

Sample definition and in-laboratory replication	Yes (indicate where provided: page no/section/legend)	n/a
State number of times the experiment was replicated in laboratory	sGFAP and sNfL concentrations were measured using a single-molecule array assay (Simoa, Quanterix, MA). sGFAP and sNfL levels were independently evaluated by a blinded examiner in duplicate, and mean intra-assay and interassay coefficients of variation were below 10.	
Define whether data describe technical or biological replicates	Not detected.	

Ethics	Yes (indicate where provided: page no/section/legend)	n/a
Studies involving human participants: State details of authority granting ethics approval (IRB or equivalent committee(s), provide reference number for approval.	Standard Protocol Approvals, Registrations, and Patient Consents The Institutional Review Board of the NCC approved the study, and written informed consent was obtained from all participants.	
Studies involving experimental animals: State details of authority granting ethics approval (IRB or equivalent committee(s), provide reference number for approval.	Not detected.	

Ethics	Yes (indicate where provided: page no/section/legend)	n/a
Studies involving specimen and field samples: State if relevant permits obtained, provide details of authority approving study; if none were required, explain why.	Not detected.	

Dual Use Research of Concern (DURC)	Yes (indicate where provided: page no/section/legend)	n/a
If study is subject to dual use research of concern, state the authority granting approval and reference number for the regulatory approval	Not currently checked by SciScore	

Analysis

Attrition	Yes (indicate where provided: page no/section/legend)	n/a
State if sample or data point from the analysis is excluded, and whether the criteria for exclusion were determined and specified in advance.	not detected.	

Statistics	Yes (indicate where provided: page no/section/legend)	n/a
Describe statistical tests used and justify choice of tests.	Statistical Analysis A Fisher exact test was applied to determine the statistical significance of the cutoff value-based positive or negative statuses of sGFAP and sNfL levels, measured at individual time points, were associated with clinical relapse or relapse-free status.	

Data availability	Yes (indicate where provided: page no/section/legend)	n/a
State whether newly created datasets are available, including protocols for access or restriction on access.	Data Availability Anonymized data not published within this article will be made available on request from any appropriately qualified investigator.	
If data are publicly available, provide accession number in repository or DOI or URL.	Not detected.	
If publicly available data are reused, provide accession number in repository or DOI or URL, where possible.	Not detected.	

Code availability	Yes (indicate where provided: page no/section/legend)	n/a
For all newly generated code and software essential for replicating the main findings of the study:		
State whether the code or software is available.	Not detected.	
If code is publicly available, provide accession number in repository, or DOI or URL.	Not detected.	

Analysis

Adherence to community standards	Yes (indicate where provided: page no/section/legend)	n/a
MDAR framework recommends adoption of discipline-specific guidelines, established and endorsed through community initiatives. Journals have their own policy about requiring specific guidelines and recommendations to complement MDAR.		
State if relevant guidelines (eg., ICMJE, MIBBI, ARRIVE) have been followed, and whether a checklist (eg., CONSORT, PRISMA, ARRIVE) is provided with the manuscript.	Not currently checked by SciScore	