Document Identifier: **PMC8054958**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Standard Protocol Approvals, Registrations, and Patient Consents The Institutional Review Board of the NCC approved the study, and written informed consent was obtained from all participants.
Consent: Standard Protocol Approvals, Registrations, and Patient Consents The Institutional Review Board of the NCC approved the study, and written informed consent was obtained from all participants.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Pretreatment samples were collected within a median of 2 months (range 1-3 months) from prior attacks.
Randomization
4,5 Twenty participants newly treated with rituximab according to Kim protocol 6 and underwent at least a 24-month follow-up were randomly selected from the NMOSD cohort of the National Cancer Center (NCC) to minimize interindividual variability of treatment 14 participants without clinical attacks (stable) during follow-up were included for focused investigation of interattack periods, and 6 participants with clinical attacks (unstable) were included for attack-related measurements.
Blinding
sGFAP and sNfL concentrations were measured using a single-molecule array assay (Simoa, Quanterix, MA). sGFAP and sNfL levels were independently evaluated by a blinded examiner in duplicate, and mean intra-assay and interassay coefficients of variation were below 10.
Power Analysis
not detected.

Replication
sGFAP and sNfL concentrations were measured using a single-molecule array assay (Simoa, Quanterix, MA). sGFAP and sNfL levels were independently evaluated by a blinded examiner in duplicate, and mean intra-assay and interassay coefficients of variation were below 10.
Number: sGFAP and sNfL concentrations were measured using a single-molecule array assay (Simoa, Quanterix, MA). sGFAP and sNfL levels were independently evaluated by a blinded examiner in duplicate, and mean intra-assay and interassay coefficients of variation were below 10.
Data Information
Availability: Data Availability Anonymized data not published within this article will be made available on request from any appropriately qualified investigator.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
All individuals satisfied the 2015 diagnostic criteria for NMOSD and were positive for AQP4 antibodies tested by live cell-based assays.	AQP4		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical Analysis A Fisher exact test was applied to determine the statistical significance of the cutoff value-based positive or negative statuses of sGFAP and sNfL levels, measured at individual time points, were associated with clinical relapse or relapse-free status.	Fisher exact test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq