Document Identifier: **PMC8075316**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
Materials and Methods Bacterial Strains and Plasmids A. hydrophila SS332 was isolated from a fecal specimen from a 76-year-old male patient in Lishui Central Hospital, Zhejiang, China in 2013.
Subject Demographics
Age: Materials and Methods Bacterial Strains and Plasmids A. hydrophila SS332 was isolated from a fecal specimen from a 76-year-old male patient in Lishui Central Hospital, Zhejiang, China in 2013.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Number: The maximum likelihood phylogenetic tree of AFM-1 was constructed by MEGA-X software with 1000 bootstrap replications .

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Recombinant DNA			
pUCP20 and pUCP24 were used as vectors in cloning experiments .	pUCP20		
	pUCP24		
Software and Algorithms			
It was identified by a microorganism autoanalysis system (VITEK(r)2, BioMerieux, France), and homologous comparisons of the 16S rRNA gene sequence to the GenBank database were performed by using the BLAST program (www.ncbi.nlm.nih.govBLAST	BLAST		Suggestion: WU-BLAST (RRID:SCR_011824)(link)
The initial complete genome was assembled from PacBio long reads by Canu v1.7 and then corrected by Pilon with Illumina short reads .	Canu		Suggestion: Canu (RRID:SCR_015880)(link)
	Pilon		Suggestion: Pilon (RRID:SCR_014731)(link)
Potential open reading frames (ORFs) were predicted using Prodigal and annotated against the UniProtSwiss-Prot and nonredundant protein databases using the BLASTX program .	Prodigal		Suggestion: Prodigal (RRID:SCR_021246)(link)
	BLASTX		Suggestion: BLASTX (RRID:SCR_001653)(link)
Annotation of ARGs and mobile genetic elements was performed using online databases , including the Comprehensive Antibiotic Resistance Database (CARD) (https:card.mcmaster . ca) and ISfinder (https:www-is . biotoul . fr) .	ISfinder		Suggestion: ISFinder (RRID:SCR_003020)(link)
The maximum likelihood phylogenetic tree of AFM-1 was constructed by MEGA-X software with 1000 bootstrap replications .	MEGA-X		
The molecular weight and pI value of AFM-1 were predicted using ProtParam2 .	ProtParam2		
The putative signal peptide cleavage site of bla AFM-1 was identified by SignalP 5.0 .	SignalP		
Comparisons of the nucleotide sequences were performed using BLASTN .	BLASTN		
Other bioinformatics tools were written using Python and Biopython .	Python		Suggestion: python-bx (RRID:SCR_024202)(link)
	Biopython		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq