Document Identifier: **PMC8088975**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: 2.6 Xenograft The study was approved by the Animal Care Committee of Iwate Medical University (
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
Authentication: 2 MATERIALS AND METHODS 2.1 Cell culture and reagents BaF3 cells 39 were provided by RIKEN BioResource Research Center and maintained in RPMI-1640 medium

containing 10 FBS, 10 ngmL recombinant mouse IL-3 (PeproTech), and penicillinstreptomycin. pBABE-EGFR L858R (11012), pBABE-EGFR del exon19 (32062), pBABE-EGFR L858RT790M (32073), and pBABE-EGFR del ex19T790M (32072) were obtained from Addgene. pBABE-EGFR L858RT790MC797S and pBABE-EGFR del ex19T790MC797S were constructed by site-directed mutagenesis with C797S-1 (GACATAGTCCAGGAGGCTGCCGAAGGGCATGAG) and C797S-2 (GACATAGTCCAGGAGTCAGCCGAAGGGCATGAG) primer set using pBABE-EGFR L858RT790M or pBABE-EGFR del ex19T790M as templates. pBABE-EGFR WT was also constructed by site-directed mutagenesis with R858L-1 (ACAGATTTTGGGCTGGCCAAACTGCTGGGT) and R858L-2 (ACCCAGCAGTTTGGCCAGCCCAAAATCTGT) primer set using pBABE-EGFR L858R as a template.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The following antibodies were used: rabbit anti-phospho-EGFR (Y1068) (1:3000 Cell Signaling 2237) , mouse anti-EGFR (1:3000 Cell Signaling 2239) , rabbit anti-phospho-AKT (S473) (1:3000 Cell Signaling 9271) , rabbit anti-AKT (1:3000 Cell Signaling 9272) , rabbit anti-phospho-ERK12 (T202Y204) (1:3000 Cell Signaling 9101) , rabbit anti-ERK12 (1:10 000 Santa Cruz sc-93) , mouse anti-actin (1:1000 Sigma C7207)	anti-phospho-EGFR (Y1068		
	anti-EGFR		
	anti-phospho-AKT (S473		
	anti-AKT		
	anti-phospho-ERK12		
	anti-ERK12		
	anti-actin		
Experimental Models: Cell Lines			
PC-9 cells were maintained in RPMI-1640 medium containing 10 FBS and penicillinstreptomycin .	PC-9		Suggestion: RRID:CVCL_B260(link)
2.4 Cell viability assay BaF3 (3000 cellswell) , PC-9 (3000 cellswell) , or A549 cells (7500 cellswell) were seeded into 96-well plates .	A549		Suggestion: RRID:CVCL_0023(link)
PC-9-EGFR-del exon19T790MC797S cells (5 x 106 cells) were suspended in 100 mL of 1:1 Matrigel and subcutaneously implanted into BALBc nunu mice (Charles River) using a 21G syringe.	PC-9-EGFR-del exon19T790MC797S		
Recombinant DNA			
2 MATERIALS AND METHODS 2.1 Cell culture and reagents BaF3 cells 39 were provided by RIKEN BioResource Research Center and maintained in RPMI-1640 medium containing 10 FBS, 10 ngmL recombinant mouse IL-3 (PeproTech), and penicillinstreptomycin. pBABE-EGFR L858R (11012), pBABE-EGFR del exon19 (32062), pBABE-EGFR L858RT790M (32073), and pBABE-EGFR del ex19T790M (32072) were obtained from Addgene. pBABE-EGFR L858RT790MC797S and pBABE-EGFR del ex19T790MC797S were constructed by site-directed mutagenesis with C797S-1 (GACATAGTCCAGGAGGCTGCCGA	pBABE-EGFR L858RT790MC797S		
	pBABE-EGFR		
	pBABE-EGFR L858RT790M		

AGGGCATGAG) and C797S-2 (GACATAGTCCAGGAGTCAGCCGA AGGGCATGAG) primer set using pBABE-EGFR L858RT790M or pBABE-EGFR del ex19T790M as templates. pBABE-EGFR WT was also constructed by site-directed mutagenesis with R858L-1 (ACAGATTTTGGGCTGGCCAAACT GCTGGGT) and R858L-2 (ACCCAGCAGTTTGGCCAGCCCAA AATCTGT) primer set using pBABE-EGFR L858R as a template.			
Software and Algorithms			
2 MATERIALS AND METHODS 2.1 Cell culture and reagents BaF3 cells 39 were provided by RIKEN BioResource Research Center and maintained in RPMI-1640 medium containing 10 FBS, 10 ngmL recombinant mouse IL-3 (PeproTech), and penicillinstreptomycin. pBABE-EGFR L858R (11012), pBABE-EGFR del exon19 (32062), pBABE-EGFR L858RT790M (32073), and pBABE-EGFR del ex19T790M (32072) were obtained from Addgene. pBABE-EGFR L858RT790MC797S and pBABE-EGFR del ex19T790MC797S were constructed by site-directed mutagenesis with C797S-1 (GACATAGTCCAGGAGGCTGCCGA AGGGCATGAG) and C797S-2 (GACATAGTCCAGGAGTCAGCCGA AGGGCATGAG) primer set using pBABE-EGFR L858RT790M or pBABE-EGFR del ex19T790M as templates. pBABE-EGFR WT was also constructed by site-directed mutagenesis with R858L-1 (ACAGATTTTGGGCTGGCCAAACT GCTGGGT) and R858L-2 (ACCCAGCAGTTTGGCCAGCCCAA AATCTGT) primer set using pBABE-EGFR L858R as a template.	RIKEN BioResource Addgene.		Suggestion: RIKEN BioResource Center (RRID:SCR_003250)(link) Suggestion: Addgene (RRID:SCR_002037)(link)
2.2 JFCR39 panel assays and COMPARE analysis The JFCR39 panel assays were developed to evaluate the antitumor profile of test compounds across a panel of 39 human cancer cell lines (termed JFCR39) , as previously described.	COMPARE		Suggestion: COMPARE (RRID:SCR_000855)(link)

Images were analyzed using ImageJ.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Fingerprints are presented in the graphic profiles of relative values of logarithmic GI50 concentrations across the JFCR39 . 40 COMPARE analysis was carried out to evaluate the mechanism of action by calculating the Pearson correlation coefficients between the fingerprint of a test compound and those of reference compounds , as previously described.	Pearson correlation
43 For the analysis , we extracted and used the signature probe sets whose expression changes after each compound treatment were significantly up- or downregulated using Students t test (fold change greater than 2 and a P value less than .05) .	t test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq