Document Identifier: **PMC8097289**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: All procedures were performed with the approval of the Animal Care and Use Committee of the Institute of Hydrobiology, Chinese Academy of Sciences. Cloning and Sequence Analysis According to the genome sequences of gibel carp clone F (CI01000363_00413629_00414546 and ENSONIP00000026093-D2), foxl2a and foxl2b cDNAs were amplified from gibel carp ovary cDNA library by 3 and 5 rapid amplification of cDNA ends (supplementary table S1, Supplementary Material online).
Inclusion and Exclusion Criteria
not detected.
Attrition
All of F0 individuals of Cgfoxl2a-B mutants grew to fertile females, of which two individuals were chosen to proliferate by gynogenesis to construct Cgfoxl2a-B mutant lines.
Sex as a biological variable
All of F0 individuals of Cgfoxl2a-B mutants grew to fertile females, of which two individuals were chosen to proliferate by gynogenesis to construct Cgfoxl2a-B mutant lines.
Subject Demographics
Age: Unfertilized eggs, 50 epiboly, hatching fry, gonads from WT, and foxl2 mutants at different stages from 10 dph to 1 year old were randomly sampled.
Randomization
Unfertilized eggs, 50 epiboly, hatching fry, gonads from WT, and foxl2 mutants at different stages from 10 dph to 1 year old were randomly sampled.
Blinding
not detected.
Power Analysis
not detected.
Replication

All the samples were analyzed in triplicates.
Number: All the samples were analyzed in triplicates.
Cell Line Authentication
not detected.
Data Information
Availability: The complete cDNA sequences of 12 foxl2 transcripts were deposited in GenBank (accession numbers from MT468195 to MT468206).

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
One microgram plasmids were transfected into HEK293T cells.	HEK293T		Suggestion: RRID:CVCL_0063(link)
Recombinant DNA			
The plasmid pCS2-Cas9 was digested with XbaI (NEB) and then was transcribed using the T7 mMessage mMachine Kit (Ambion) (Liu et al. 2014).	pCS2-Cas9		
Subcellular Localization WT Cgfoxl2a-B2, Cgfoxl2a-B2D54 , Cgfoxl2a-B1D55 , and Cgfoxl2a-B3D25 + 8 were subcloned into XhoIKpnI sites of pEGFP-N3 vector (BD Biosciences Clontech).	pEGFP-N3		
Plasmid Constructs For luciferase assays, expression plasmids were generated by cloning the full length ORF of WT Cgfoxl2a-B2, Cgfoxl2a-B2D54 , Cgfoxl2a-B1D55 , and Cgfoxl2a-B3D25 + 8 were subcloned into KpnIXhoI sites of pcDNA3.1(+) vector.	pcDNA3.1(+)		
For promoter activity analysis, 1,329 and 1,140 bp 5-flanking regulatory region of Cgcyp19a1a-A promoter (GenBank accession number MW160449) and Cgcyp19a1a-B promoter (GenBank accession number MW160450) were subcloned into KpnIXhoI sites of pGL3-basic luciferase reporter vector (Cgcyp19a1a-Apro-luc and Cgcyp19a1a-Bpro-luc), respectively.	pGL3-basic		
Transfection and Luciferase Activity Assays CAB and EPC were seeded overnight in 24-well plates and transfected using FuGENE HD Transfection Reagent (Promega) with various plasmids at a ratio of 10:10:1 (200 ng expression plasmid: 200 ng Cgcyp19a1a-A pro-luc or Cgcyp19a1a-B pro-luc plasmid: 20 ng Renilla luciferase plasmid pRL-TK).	pRL-TK		
Empty vector (pcDNA3.1) was adjusted to ensure DNA concentration in total among wells.	pcDNA3.1		Suggestion: RRID:Addgene_100544(link)
Software and Algorithms			

Multiple amino acid sequence alignment and phylogenetic analysis were performed with DNAMAN version 7.0 and Molecular Evolutionary Genetics Analysis (MEGA 7) (Kumar et al. 2016).	MEGA		Suggestion: MEGA (RRID:SCR_023017)(link)
All the amino acid sequences used in this study were obtained from GenBank (http:www.ncbi.nlm.nih.gov, last accessed January 23, 2021).	http:www.ncbi.nlm.nih.gov		
The latter information was obtained from the Ensembl (http:www.ensembl.org, last accessed January 23, 2021).	Ensembl		Suggestion: Ensembl (RRID:SCR_002344)(link)
	http:www.ensembl.org		
All experiments were performed in triplicate and the significant differences were calculated by SPSS software (SPSS Inc.).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
For statistical analysis, Tukeys test was calculated with SPSS software (SPSS Inc.).	Tukeys test
Unresolved RRIDs	
Transfection and Luciferase Activity Assays CAB and EPC were seeded overnight in 24-well plates and transfected using FuGENE HD Transfection Reagent (Promega) with various plasmids at a ratio of 10:10:1 (200 ng expression plasmid: 200 ng Cgcyp19a1a-A pro-luc or Cgcyp19a1a-B pro-luc plasmid: 20 ng Renilla luciferase plasmid pRL-TK).	plasmid: 200
	plasmid: 20

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq