Document Identifier: **PMC8111093**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The test was carried out according to a specific protocol approved by the Bioethics Committee of Medical University of Bialystok (R-l-002/4/2019) .
Inclusion and Exclusion Criteria
Table 1 Group Tissue sample source No. of samples No. of force curves analyzed I Squamous cell hyperplasia with low grade dysplasia (leukoplakia) 6 n = 2026 II Oral squamous cell carcinoma 4 n = 3458 III Healthy buccal mucosa (control) 3 n = 564 IV Free margin of the squamous cell hyperplasia with low grade dysplasia (leukoplakia) 6 n = 1589 Inclusion criteria for this study were age 18–72 years and a clinical diagnosis of keratosis with simultaneous overgrowth folds of the mucous membrane outside the lesion requiring a cut or with a clinically healthy mucous membrane of the mouth but an overgrown frenulum lip cheek .
Attrition
not detected.
Sex as a biological variable
Group I consisted of 6 patients (3 women and 3 men) diagnosed with squamous cell hyperplasia with low grade dysplasia (leukoplakia), group II included 4 patients (2 women and 2 men) diagnosed with OSCC, and group III (control group) consisted of 3 patients (1 woman and 2 men) with healthy mouth mucosa.
Subject Demographics
Age: Table 1 Group Tissue sample source No. of samples No. of force curves analyzed I Squamous cell hyperplasia with low grade dysplasia (leukoplakia) 6 n = 2026 II Oral squamous cell carcinoma 4 n = 3458 III Healthy buccal mucosa (control) 3 n = 564 IV Free margin of the squamous cell hyperplasia with low grade dysplasia (leukoplakia) 6 n = 1589 Inclusion criteria for this study were age 18–72 years and a clinical diagnosis of keratosis with simultaneous overgrowth folds of the mucous membrane outside the lesion requiring a cut or with a clinically healthy mucous membrane of the mouth but an overgrown frenulum lip cheek .
Randomization
not detected.
Blinding
not detected.

Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Subsequently , the following antibodies were used: Bcl-2 (DAKO M0887	Bcl-2		
mouse monoclonal antibody , 1:100 dilution) , P16 CINtec (Histology Kit , Ventana Roche , clone E6H4 , mouse monoclonal) , p53 (DAKO mouse monoclonal antibody M 7001 , clone DO-7) and collagen IV (mouse monoclonal antibody COL4A1 (5E10): sc-517572 Santa Cruz Biotechnology , 1:50	p53		
For the blood vessels assessment , we used CD31 antibody (CD31 , Endothelial Cell (Dako Omnis Clone JC70A)	CD31		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq