Document Identifier: **PMC8139582**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Euthanasia Agents: For lentiviral Zeb1 shRNA or control shRNA treatment , K-RasLA1 mice at 1 month of age were anesthetized by intraperitoneal injection of a combination of ketamine (100 mg/kg) and xylazine (10 mg/kg) .
IACUC: The treated mice were then housed in a biosafety room and closely monitored daily in accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) .
Inclusion and Exclusion Criteria
Data inclusion/exclusion criteria: No data were excluded.
Attrition
not detected.
Sex as a biological variable
On the basis of our previous studies, we did not detect any differences with regard to sex, and thus, male and female mice were chosen at random.
Subject Demographics
Age: For lentiviral Zeb1 shRNA or control shRNA treatment , K-RasLA1 mice at 1 month of age were anesthetized by intraperitoneal injection of a combination of ketamine (100 mg/kg) and xylazine (10 mg/kg) .
Randomization
On the basis of our previous studies, we did not detect any differences with regard to sex, and thus, male and female mice were chosen at random.
Blinding
Tumor pathology was evaluated blindly by an experienced pathologist (M. Cu).
Power Analysis
not detected.
Replication

Replicates: The number of experimental replications is provided in the figure legends, and P values are provided.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Cells were cocultured in the absence or presence of neutralizing anti-CD47 monoclonal antibody (mAb, 10 μg/ml) or control mAb (BioXCell) for 48 hours and then immunostained as described.	anti-CD47		
Immunoprecipitation with histone 3 antibody (H3) included in an EpiTect ChIP antibody kit (Qiagen , catalog no. GAH-2206) was used as a positive control .	histone 3		
Experimental Models: Cell Lines			
Briefly , 293T cells were transfected with the lentiviral vector and packaging plasmids , and the supernatants , containing recombinant pseudolentiviral particles , were collected from culture dishes on the second and third days after transfection .	293T		Suggestion: RRID:CVCL_0063(link)
Experimental Models: Organisms/Strains			
K-rasLA1 mice (36) in a C57BL/6 background were obtained from the Jackson laboratory.	K-rasLA1		
These mice were crossed with Zeb1 (+/–) mice (55) also in a C57BL/6 background to yield K-RasLA1, Zeb1 (+/+) and K-RasLA1, Zeb1 (+/–) mice.	C57BL/6		Suggestion: RRID:IMSR_JAX:000664(link)
Recombinant DNA			
To generate a lentivirus expressing Zeb1 shRNA , a primer containing a Zeb1 shRNA sequence (5#-CTGTCTAGACAAAAAAAGACA ACGTGAAAGACAATCTCTTGAATT GTCAAACACGTTGTCTTGGGGATC TGTGGTCTCATACA-3#) was used with a T3 primer to amplify a 500–base pair fragment containing the H1 promoter using the pSuper vector as a template.	pSuper		Suggestion: RRID:Addgene_51598(link)
Software and Algorithms			
Images shown are maximum projections of confocal stacks , adjusted for contrast and brightness	Adobe Photoshop		Suggestion: Adobe Photoshop, RRID:SCR_014199(link)

with Adobe Photoshop Elements v9.0.2			

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Significance was determined by the Student's t test.	t test
Oligonucleotides	
To generate a lentivirus expressing Zeb1 shRNA , a primer containing a Zeb1 shRNA sequence (5#-CTGTCTAGACAAAAAAAGACA ACGTGAAAGACAATCTCTTGAATT GTCAAACACGTTGTCTTGGGGATC TGTGGTCTCATACA-3#) was used with a T3 primer to amplify a 500–base pair fragment containing the H1 promoter using the pSuper vector as a template.	5#-CTGTCTAGACAAAAAAAGACA ACGTGAAAGACAATCTCTTGAATT GTCAAACACGTTGTCTTGGGGATC TGTGGTCTCATACA-3#
The long oligonucleotide was used , together with a sp6 oligonucleotide (5#-ATTTAGGTGACACTATAGAAT -3#) , to PCR-amplify a fragment containing the entire U6 promoter plus shRNA sequences; the resultant product was digested with XbaI and SpeI ligated into the NheI of the lentivirus vector , and the insert was sequenced to ensure that no errors occurred during the PCR or cloning steps .	5#-ATTTAGGTGACACTATAGAAT -3#
Lentivirus with a scrambled shRNA sequence 5#-CAACAAGATGAAGAGCACCAA TCTCTTGAAT TGGTGCTCTTCATCTTGTTG-3# was used as a control—this control sequence was blasted against all mouse RNA sequences to insure that it did not target an mRNA .	5#-CAACAAGATGAAGAGCACCAA TCTCTTGAAT TGGTGCTCTTCATCTTGTTG-3#

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq